

Table 1: All cancers (ICD-10 C00–C97 excluding skin cancers other than melanoma C44)

Australia 2000

Age group	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
0–4	153	23.3	108	17.3	261	20.4	17	2.6	13	2.1	30	2.3
5–9	65	9.4	58	8.8	123	9.1	25	3.6	16	2.4	41	3.0
10–14	90	13.1	86	13.2	176	13.2	14	2.0	22	3.4	36	2.7
15–19	165	24.3	139	21.4	304	22.9	29	4.3	19	2.9	48	3.6
20–24	269	41.1	251	39.5	520	40.3	37	5.6	34	5.3	71	5.5
25–29	425	58.8	443	60.9	868	59.9	41	5.7	40	5.5	81	5.6
30–34	570	80.5	766	106.6	1,336	93.6	89	12.6	96	13.4	185	13.0
35–39	804	107.4	1,278	169.0	2,082	138.4	136	18.2	194	25.6	330	21.9
40–44	1,166	162.0	1,944	266.7	3,110	214.7	269	37.4	342	46.9	611	42.2
45–49	1,739	260.7	2,746	407.3	4,485	334.4	473	70.9	560	83.1	1,033	77.0
50–54	2,911	458.8	3,592	576.4	6,503	517.1	947	149.3	861	138.2	1,808	143.8
55–59	4,059	828.0	3,678	776.8	7,737	802.9	1,372	279.9	1,067	225.4	2,439	253.1
60–64	5,215	1,301.2	3,887	979.5	9,102	1,141.1	1,834	457.6	1,216	306.4	3,050	382.4
65–69	6,470	1,948.6	4,030	1,167.8	10,500	1,550.7	2,597	782.1	1,561	452.4	4,158	614.1
70–74	7,709	2,573.2	4,700	1,408.7	12,409	1,959.6	3,582	1,195.6	2,171	650.7	5,753	908.5
75–79	6,974	3,175.9	4,752	1,651.5	11,726	2,311.3	3,663	1,668.1	2,456	853.5	6,119	1,206.1
80–84	4,203	3,532.9	3,429	1,804.7	7,632	2,470.2	2,618	2,200.6	2,089	1,099.5	4,707	1,523.5
85 and over	2,948	3,803.0	3,409	1,946.3	6,357	2,515.9	2,295	2,960.6	2,671	1,525.0	4,966	1,965.4
Total	45,935		39,296		85,231		20,038		15,428		35,466	

Rates per 100,000 with 95% confidence intervals (95% CI)

Crude rate	483.3	407.3	445.0	210.8	159.9	185.2
95% CI	478.8 – 487.7	403.3 – 411.3	442.0 – 448.0	207.9 – 213.7	157.4 – 162.4	183.2 – 187.1
AS rate (Aust. 2001)	535.7	390.4	450.9	245.0	147.5	188.3
95% CI	530.8 – 540.7	386.5 – 394.3	447.8 – 453.9	241.6 – 248.5	145.1 – 149.8	186.3 – 190.3
AS rate (new WHO World)	382.6	297.4	333.7	161.2	101.6	127.4
95% CI	379.1 – 386.1	294.4 – 300.5	331.4 – 336.0	159.0 – 163.5	99.9 – 103.3	126.1 – 128.8
Lifetime risk (0–74)	1 in 3	1 in 4	1 in 3	1 in 7	1 in 11	1 in 9
PYLL (0–74)				136,660	116,425	253,085
Per cent of all cancers	100.0	100.0	100.0	100.0	100.0	100.0

Average annual numbers and rates by State and Territory, 1996–2000

	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate
NSW	15,223	536.8	12,634	377.9	27,857	443.5	6,636	245.9	5,141	147.5	11,777	187.8
Vic	11,098	543.9	9,685	393.2	20,784	454.0	5,025	256.5	4,030	157.2	9,055	197.9
Qld	8,386	574.3	6,879	411.4	15,266	481.4	3,586	259.9	2,539	149.9	6,126	197.2
WA	3,757	518.2	3,144	367.2	6,900	430.2	1,717	254.1	1,303	151.7	3,019	194.5
SA	3,962	556.0	3,308	390.1	7,270	457.9	1,734	250.3	1,374	152.1	3,107	192.2
Tas	1,151	533.5	1,001	394.3	2,152	450.4	561	270.1	450	168.8	1,011	210.8
ACT	558	559.0	485	381.1	1,042	454.9	232	263.8	217	185.8	450	217.2
NT	229	480.0	171	367.1	400	427.1	93	244.4	70	199.7	163	223.3

Notes

1. AS rates use Australian 2001 Standard Population unless new WHO World Standard Population is indicated. All rates are expressed per 100,000 population.

2. Incidence numbers are based on state of usual residence at time of diagnosis; mortality numbers on state of registration of death.

Source: *Cancer in Australia 2000*, AIHW & AACR, 2003.

Table 2: Cancer of the lip (ICD-10 C00)

Australia 2000

Age group	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
0-4	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
5-9	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
10-14	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
15-19	1	0.1	0	0.0	1	0.1	0	0.0	0	0.0	0	0.0
20-24	8	1.2	2	0.3	10	0.8	0	0.0	0	0.0	0	0.0
25-29	18	2.5	1	0.1	19	1.3	0	0.0	0	0.0	0	0.0
30-34	23	3.2	5	0.7	28	2.0	0	0.0	0	0.0	0	0.0
35-39	46	6.1	10	1.3	56	3.7	0	0.0	0	0.0	0	0.0
40-44	52	7.2	16	2.2	68	4.7	0	0.0	0	0.0	0	0.0
45-49	53	7.9	8	1.2	61	4.5	0	0.0	0	0.0	0	0.0
50-54	70	11.0	19	3.0	89	7.1	2	0.3	0	0.0	2	0.2
55-59	66	13.5	23	4.9	89	9.2	0	0.0	0	0.0	0	0.0
60-64	71	17.7	16	4.0	87	10.9	1	0.2	0	0.0	1	0.1
65-69	42	12.6	25	7.2	67	9.9	1	0.3	0	0.0	1	0.1
70-74	62	20.7	32	9.6	94	14.8	3	1.0	0	0.0	3	0.5
75-79	76	34.6	38	13.2	114	22.5	3	1.4	2	0.7	5	1.0
80-84	42	35.3	27	14.2	69	22.3	0	0.0	0	0.0	0	0.0
85 and over	26	33.5	36	20.6	62	24.5	3	3.9	4	2.3	7	2.8
Total	656		258		914		13		6		19	

Rates per 100,000 with 95% confidence intervals (95% CI)

Crude rate	6.9	2.7	4.8	0.1	0.1	0.1
95% CI	6.4 – 7.4	2.3 – 3.0	4.5 – 5.1	0.1 – 0.2	0.0 – 0.1	0.1 – 0.1
AS rate (Aust. 2001)	7.4	2.5	4.8	0.2	0.0	0.1
95% CI	6.8 – 8.0	2.2 – 2.8	4.5 – 5.1	0.1 – 0.3	0.0 – 0.1	0.1 – 0.1
AS rate (new WHO World)	5.7	1.8	3.7	0.1	0.0	0.1
95% CI	5.3 – 6.1	1.6 – 2.0	3.4 – 3.9	0.0 – 0.2	0.0 – 0.0	0.0 – 0.1
Lifetime risk (0-74)	1 in 193	1 in 578	1 in 290	1 in 10,711	#DIV/0!	1 in 22,079
PYLL (0-74)				73	0	73
Per cent of all cancers	1.4	0.7	1.1	0.1	0.0	0.1

Average annual numbers and rates by State and Territory, 1996-2000

	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate
NSW	202	7.1	75	2.2	277	4.4	3	0.1	2	0.1	5	0.1
Vic	175	8.4	80	3.1	256	5.6	2	0.1	1	0.0	3	0.1
Qld	192	12.5	67	4.0	259	8.1	2	0.1	1	0.0	3	0.1
WA	68	8.5	26	3.0	94	5.7	1	0.1	0	0.0	1	0.1
SA	90	12.6	33	3.8	123	7.8	1	0.1	0	0.0	1	0.1
Tas	31	14.0	11	4.3	42	8.8	1	0.5	0	0.1	1	0.3
ACT	4	2.8	3	2.2	6	2.5	0	0.1	0	0.0	0	0.1
NT	5	7.6	1	3.6	6	5.9	0	0.5	0	0.0	0	0.3

Notes

1. AS rates use Australian 2001 Standard Population unless new WHO World Standard Population is indicated. All rates are expressed per 100,000 population.

2. Incidence numbers are based on state of usual residence at time of diagnosis; mortality numbers on state of registration of death.

Source: *Cancer in Australia 2000*, AIHW & AACR, 2003.

Table 3: Cancer of the tongue (ICD-10 C01–C02)

Australia 2000

Age group	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
0–4	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
5–9	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
10–14	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
15–19	0	0.0	1	0.2	1	0.1	1	0.1	0	0.0	1	0.1
20–24	3	0.5	1	0.2	4	0.3	0	0.0	1	0.2	1	0.1
25–29	0	0.0	3	0.4	3	0.2	0	0.0	0	0.0	0	0.0
30–34	3	0.4	5	0.7	8	0.6	1	0.1	0	0.0	1	0.1
35–39	12	1.6	3	0.4	15	1.0	1	0.1	2	0.3	3	0.2
40–44	20	2.8	7	1.0	27	1.9	1	0.1	4	0.5	5	0.3
45–49	33	4.9	9	1.3	42	3.1	6	0.9	4	0.6	10	0.7
50–54	32	5.0	19	3.0	51	4.1	8	1.3	7	1.1	15	1.2
55–59	37	7.5	16	3.4	53	5.5	18	3.7	4	0.8	22	2.3
60–64	46	11.5	10	2.5	56	7.0	6	1.5	2	0.5	8	1.0
65–69	41	12.3	16	4.6	57	8.4	12	3.6	9	2.6	21	3.1
70–74	30	10.0	19	5.7	49	7.7	22	7.3	4	1.2	26	4.1
75–79	18	8.2	14	4.9	32	6.3	9	4.1	4	1.4	13	2.6
80–84	11	9.2	9	4.7	20	6.5	7	5.9	9	4.7	16	5.2
85 and over	8	10.3	18	10.3	26	10.3	10	12.9	14	8.0	24	9.5
Total	294		150		444		102		64		166	

Rates per 100,000 with 95% confidence intervals (95% CI)

Crude rate	3.1	1.6	2.3	1.1	0.7	0.9
95% CI	2.7 – 3.4	1.3 – 1.8	2.1 – 2.5	0.9 – 1.3	0.5 – 0.8	0.7 – 1.0
AS rate (Aust. 2001)	3.2	1.5	2.3	1.2	0.6	0.9
95% CI	2.9 – 3.6	1.2 – 1.7	2.1 – 2.6	1.0 – 1.4	0.5 – 0.8	0.7 – 1.0
AS rate (new WHO World)	2.6	1.1	1.8	0.9	0.4	0.6
95% CI	2.3 – 2.9	0.9 – 1.3	1.7 – 2.0	0.7 – 1.0	0.3 – 0.6	0.5 – 0.7
Lifetime risk (0–74)	1 in 354	1 in 856	1 in 502	1 in 1,062	1 in 2,551	1 in 1,516
PYLL (0–74)				1,050	698	1,748
Per cent of all cancers	0.6	0.4	0.5	0.5	0.4	0.5

Average annual numbers and rates by State and Territory, 1996–2000

	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate
NSW	103	3.5	52	1.5	155	2.5	37	1.3	19	0.6	56	0.9
Vic	64	3.0	35	1.4	100	2.2	29	1.4	14	0.5	43	0.9
Qld	60	3.8	34	2.0	94	2.9	22	1.5	9	0.5	31	1.0
WA	25	3.2	11	1.3	35	2.2	11	1.5	5	0.6	16	1.0
SA	17	2.3	12	1.4	29	1.8	8	1.1	5	0.5	13	0.8
Tas	7	2.9	3	1.1	10	2.0	2	1.0	1	0.3	3	0.6
ACT	3	3.4	2	1.6	5	2.4	1	0.8	1	0.4	1	0.6
NT	3	5.5	1	1.8	4	3.8	1	1.8	0	0.8	1	1.4

Notes

- AS rates use Australian 2001 Standard Population unless new WHO World Standard Population is indicated. All rates are expressed per 100,000 population.
 - Incidence numbers are based on state of usual residence at time of diagnosis; mortality numbers on state of registration of death.
- Source: *Cancer in Australia 2000*, AIHW & AACR, 2003.

Table 4: Cancer of the salivary gland (ICD-10 C07–C08)

Australia 2000

Age group	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
0–4	1	0.2	0	0.0	1	0.1	0	0.0	0	0.0	0	0.0
5–9	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
10–14	1	0.1	1	0.2	2	0.1	0	0.0	0	0.0	0	0.0
15–19	2	0.3	0	0.0	2	0.2	0	0.0	0	0.0	0	0.0
20–24	4	0.6	4	0.6	8	0.6	0	0.0	0	0.0	0	0.0
25–29	1	0.1	5	0.7	6	0.4	0	0.0	0	0.0	0	0.0
30–34	1	0.1	4	0.6	5	0.4	1	0.1	0	0.0	1	0.1
35–39	4	0.5	3	0.4	7	0.5	1	0.1	0	0.0	1	0.1
40–44	5	0.7	9	1.2	14	1.0	1	0.1	0	0.0	1	0.1
45–49	2	0.3	6	0.9	8	0.6	0	0.0	0	0.0	0	0.0
50–54	6	0.9	9	1.4	15	1.2	3	0.5	1	0.2	4	0.3
55–59	15	3.1	1	0.2	16	1.7	1	0.2	1	0.2	2	0.2
60–64	17	4.2	7	1.8	24	3.0	4	1.0	1	0.3	5	0.6
65–69	16	4.8	3	0.9	19	2.8	5	1.5	2	0.6	7	1.0
70–74	25	8.3	7	2.1	32	5.1	7	2.3	3	0.9	10	1.6
75–79	19	8.7	12	4.2	31	6.1	7	3.2	1	0.3	8	1.6
80–84	17	14.3	6	3.2	23	7.4	5	4.2	4	2.1	9	2.9
85 and over	11	14.2	11	6.3	22	8.7	5	6.5	4	2.3	9	3.6
Total	147		88		235		40		17		57	

Rates per 100,000 with 95% confidence intervals (95% CI)

Crude rate	1.5	0.9	1.2	0.4	0.2	0.3
95% CI	1.3 – 1.8	0.7 – 1.1	1.1 – 1.4	0.3 – 0.6	0.1 – 0.3	0.2 – 0.4
AS rate (Aust. 2001)	1.7	0.9	1.2	0.5	0.2	0.3
95% CI	1.4 – 2.0	0.7 – 1.0	1.1 – 1.4	0.3 – 0.6	0.1 – 0.2	0.2 – 0.4
AS rate (new WHO World)	1.2	0.7	0.9	0.3	0.1	0.2
95% CI	1.0 – 1.4	0.5 – 0.8	0.8 – 1.1	0.2 – 0.4	0.1 – 0.2	0.1 – 0.3
Lifetime risk (0–74)	1 in 819	1 in 1,829	1 in 1,143	1 in 3,373	1 in 9,513	1 in 5,037
PYLL (0–74)				303	75	378
Per cent of all cancers	0.3	0.2	0.3	0.2	0.1	0.2

Average annual numbers and rates by State and Territory, 1996–2000

	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate
NSW	45	1.6	29	0.9	74	1.2	12	0.5	7	0.2	20	0.3
Vic	25	1.3	20	0.8	46	1.0	13	0.7	5	0.2	18	0.4
Qld	30	2.0	14	0.8	44	1.4	5	0.4	4	0.2	9	0.3
WA	12	1.7	8	0.9	20	1.2	5	0.8	1	0.1	6	0.4
SA	11	1.6	7	0.8	19	1.2	4	0.6	2	0.2	6	0.4
Tas	3	1.4	2	0.9	5	1.1	2	0.8	0	0.0	2	0.3
ACT	4	3.7	1	0.7	5	2.0	0	0.5	0	0.4	1	0.4
NT	1	1.2	0	0.7	1	0.9	0	0.4	0	0.5	1	0.4

Notes

1. AS rates use Australian 2001 Standard Population unless new WHO World Standard Population is indicated. All rates are expressed per 100,000 population.

2. Incidence numbers are based on state of usual residence at time of diagnosis; mortality numbers on state of registration of death.

Source: *Cancer in Australia 2000*, AIHW & AACR, 2003.

Table 5: Cancer of the mouth (ICD-10 C03–C06)

Australia 2000

Age group	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
0–4	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
5–9	1	0.1	0	0.0	1	0.1	0	0.0	0	0.0	0	0.0
10–14	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
15–19	1	0.1	1	0.2	2	0.2	0	0.0	0	0.0	0	0.0
20–24	1	0.2	1	0.2	2	0.2	0	0.0	0	0.0	0	0.0
25–29	2	0.3	1	0.1	3	0.2	0	0.0	1	0.1	1	0.1
30–34	1	0.1	6	0.8	7	0.5	0	0.0	0	0.0	0	0.0
35–39	4	0.5	7	0.9	11	0.7	1	0.1	1	0.1	2	0.1
40–44	9	1.3	13	1.8	22	1.5	2	0.3	0	0.0	2	0.1
45–49	21	3.1	7	1.0	28	2.1	2	0.3	1	0.1	3	0.2
50–54	29	4.6	15	2.4	44	3.5	6	0.9	3	0.5	9	0.7
55–59	39	8.0	13	2.7	52	5.4	8	1.6	3	0.6	11	1.1
60–64	44	11.0	19	4.8	63	7.9	14	3.5	5	1.3	19	2.4
65–69	41	12.3	20	5.8	61	9.0	18	5.4	5	1.4	23	3.4
70–74	41	13.7	18	5.4	59	9.3	22	7.3	6	1.8	28	4.4
75–79	27	12.3	26	9.0	53	10.4	8	3.6	8	2.8	16	3.2
80–84	17	14.3	29	15.3	46	14.9	3	2.5	9	4.7	12	3.9
85 and over	8	10.3	19	10.8	27	10.7	6	7.7	8	4.6	14	5.5
Total	286		195		481		90		50		140	

Rates per 100,000 with 95% confidence intervals (95% CI)

Crude rate	3.0	2.0	2.5	0.9	0.5	0.7
95% CI	2.7 – 3.4	1.7 – 2.3	2.3 – 2.7	0.8 – 1.1	0.4 – 0.7	0.6 – 0.9
AS rate (Aust. 2001)	3.2	1.9	2.5	1.0	0.5	0.7
95% CI	2.8 – 3.6	1.6 – 2.2	2.3 – 2.8	0.8 – 1.2	0.3 – 0.6	0.6 – 0.9
AS rate (new WHO World)	2.4	1.4	1.9	0.7	0.3	0.5
95% CI	2.2 – 2.7	1.2 – 1.6	1.7 – 2.1	0.6 – 0.9	0.2 – 0.4	0.4 – 0.6
Lifetime risk (0–74)	1 in 362	1 in 765	1 in 494	1 in 1,024	1 in 3,312	1 in 1,585
PYLL (0–74)				798	348	1,145
Per cent of all cancers	0.6	0.5	0.6	0.4	0.3	0.4

Average annual numbers and rates by State and Territory, 1996–2000

	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate
NSW	105	3.6	61	1.8	166	2.6	28	0.9	17	0.5	45	0.7
Vic	64	3.0	49	2.0	113	2.5	15	0.7	13	0.5	28	0.6
Qld	55	3.5	32	1.9	87	2.7	12	0.8	8	0.5	20	0.6
WA	29	3.7	15	1.7	44	2.7	9	1.2	5	0.6	14	0.9
SA	20	2.8	14	1.6	35	2.2	8	1.1	5	0.5	12	0.8
Tas	8	3.6	5	2.1	13	2.8	3	1.1	1	0.5	4	0.8
ACT	4	3.2	2	1.5	6	2.4	2	1.4	1	0.6	2	1.0
NT	3	3.8	1	1.7	4	2.9	1	1.0	0	0.6	1	0.8

Notes

1. AS rates use Australian 2001 Standard Population unless new WHO World Standard Population is indicated. All rates are expressed per 100,000 population.

2. Incidence numbers are based on state of usual residence at time of diagnosis; mortality numbers on state of registration of death.

Source: *Cancer in Australia 2000*, AIHW & AACR, 2003.

Table 6: Cancer of the gum (ICD-10 C03)

Australia 2000

Age group	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
0-4	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
5-9	1	0.1	0	0.0	1	0.1	0	0.0	0	0.0	0	0.0
10-14	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
15-19	1	0.1	0	0.0	1	0.1	0	0.0	0	0.0	0	0.0
20-24	1	0.2	0	0.0	1	0.1	0	0.0	0	0.0	0	0.0
25-29	0	0.0	1	0.1	1	0.1	0	0.0	0	0.0	0	0.0
30-34	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
35-39	0	0.0	0	0.0	0	0.0	0	0.0	1	0.1	1	0.1
40-44	2	0.3	1	0.1	3	0.2	0	0.0	0	0.0	0	0.0
45-49	5	0.7	1	0.1	6	0.4	0	0.0	0	0.0	0	0.0
50-54	5	0.8	2	0.3	7	0.6	1	0.2	0	0.0	1	0.1
55-59	3	0.6	1	0.2	4	0.4	0	0.0	0	0.0	0	0.0
60-64	5	1.2	5	1.3	10	1.3	1	0.2	1	0.3	2	0.3
65-69	6	1.8	5	1.4	11	1.6	1	0.3	1	0.3	2	0.3
70-74	2	0.7	0	0.0	2	0.3	0	0.0	1	0.3	1	0.2
75-79	3	1.4	5	1.7	8	1.6	1	0.5	3	1.0	4	0.8
80-84	4	3.4	8	4.2	12	3.9	1	0.8	1	0.5	2	0.6
85 and over	0	0.0	3	1.7	3	1.2	1	1.3	1	0.6	2	0.8
Total	38		32		70		6		9		15	

Rates per 100,000 with 95% confidence intervals (95% CI)

Crude rate	0.4	0.3	0.4	0.1	0.1	0.1
95% CI	0.3 – 0.5	0.2 – 0.4	0.3 – 0.5	0.0 – 0.1	0.0 – 0.2	0.0 – 0.1
AS rate (Aust. 2001)	0.4	0.3	0.4	0.1	0.1	0.1
95% CI	0.3 – 0.6	0.2 – 0.4	0.3 – 0.5	0.0 – 0.1	0.0 – 0.1	0.0 – 0.1
AS rate (new WHO World)	0.3	0.2	0.3	0.0	0.1	0.1
95% CI	0.2 – 0.4	0.1 – 0.3	0.2 – 0.3	0.0 – 0.1	0.0 – 0.1	0.0 – 0.1
Lifetime risk (0-74)	1 in 3,034	1 in 5,459	1 in 3,910	1 in 28,237	1 in 20,541	1 in 23,530
PYLL (0-74)				43	60	103
Per cent of all cancers	0.1	0.1	0.1	0.0	0.1	0.0

Average annual numbers and rates by State and Territory, 1996-2000

	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate
NSW	15	0.5	12	0.4	27	0.4	4	0.1	3	0.1	6	0.1
Vic	9	0.4	7	0.3	16	0.4	2	0.1	1	0.1	3	0.1
Qld	6	0.4	5	0.3	11	0.3	1	0.1	1	0.1	3	0.1
WA	4	0.5	3	0.3	7	0.4	0	0.1	1	0.1	1	0.1
SA	1	0.1	2	0.2	3	0.2	1	0.1	1	0.1	2	0.1
Tas	0	0.1	0	0.1	0	0.1	0	0.1	0	0.0	0	0.0
ACT	0	0.0	0	0.0	0	0.0	0	0.2	0	0.0	0	0.1
NT	0	0.2	0	0.0	0	0.1	0	0.5	0	0.0	0	0.3

Notes

1. AS rates use Australian 2001 Standard Population unless new WHO World Standard Population is indicated. All rates are expressed per 100,000 population.
 2. Incidence numbers are based on state of usual residence at time of diagnosis; mortality numbers on state of registration of death.
- Source: *Cancer in Australia 2000*, AIHW & AACR, 2003.

Table 7: Cancer of the floor of mouth (ICD-10 C04)

Australia 2000

Age group	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
0-4	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
5-9	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
10-14	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
15-19	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
20-24	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
25-29	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
30-34	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
35-39	2	0.3	1	0.1	3	0.2	1	0.1	0	0.0	1	0.1
40-44	3	0.4	3	0.4	6	0.4	0	0.0	0	0.0	0	0.0
45-49	11	1.6	4	0.6	15	1.1	2	0.3	1	0.1	3	0.2
50-54	14	2.2	6	1.0	20	1.6	2	0.3	1	0.2	3	0.2
55-59	20	4.1	6	1.3	26	2.7	6	1.2	1	0.2	7	0.7
60-64	18	4.5	8	2.0	26	3.3	7	1.7	2	0.5	9	1.1
65-69	14	4.2	4	1.2	18	2.7	6	1.8	3	0.9	9	1.3
70-74	18	6.0	6	1.8	24	3.8	12	4.0	3	0.9	15	2.4
75-79	16	7.3	7	2.4	23	4.5	3	1.4	2	0.7	5	1.0
80-84	6	5.0	5	2.6	11	3.6	2	1.7	1	0.5	3	1.0
85 and over	2	2.6	5	2.9	7	2.8	2	2.6	1	0.6	3	1.2
Total	124		55		179		43		15		58	

Rates per 100,000 with 95% confidence intervals (95% CI)

Crude rate	1.3	0.6	0.9	0.5	0.2	0.3
95% CI	1.1 – 1.5	0.4 – 0.7	0.8 – 1.1	0.3 – 0.6	0.1 – 0.2	0.2 – 0.4
AS rate (Aust. 2001)	1.4	0.5	0.9	0.5	0.1	0.3
95% CI	1.1 – 1.6	0.4 – 0.7	0.8 – 1.1	0.3 – 0.6	0.1 – 0.2	0.2 – 0.4
AS rate (new WHO World)	1.0	0.4	0.7	0.4	0.1	0.2
95% CI	0.9 – 1.2	0.3 – 0.5	0.6 – 0.8	0.3 – 0.5	0.1 – 0.2	0.2 – 0.3
Lifetime risk (0-74)	1 in 858	1 in 2,398	1 in 1,272	1 in 2,099	1 in 7,163	1 in 3,289
PYLL (0-74)				405	123	528
Per cent of all cancers	0.3	0.1	0.2	0.2	0.1	0.2

Average annual numbers and rates by State and Territory, 1996-2000

	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate
NSW	45	1.5	16	0.5	61	1.0	12	0.4	5	0.1	17	0.3
Vic	26	1.2	14	0.6	40	0.9	6	0.3	5	0.2	11	0.2
Qld	24	1.5	10	0.6	34	1.1	7	0.5	3	0.2	10	0.3
WA	11	1.3	4	0.5	15	0.9	3	0.3	1	0.1	3	0.2
SA	10	1.4	3	0.4	13	0.9	3	0.5	1	0.1	4	0.3
Tas	4	1.8	1	0.5	5	1.1	2	0.7	0	0.1	2	0.4
ACT	2	1.3	0	0.0	2	0.6	1	1.0	0	0.0	1	0.5
NT	1	1.7	0	0.7	2	1.2	0	0.4	0	0.3	0	0.3

Notes

1. AS rates use Australian 2001 Standard Population unless new WHO World Standard Population is indicated. All rates are expressed per 100,000 population.

2. Incidence numbers are based on state of usual residence at time of diagnosis; mortality numbers on state of registration of death.

Source: *Cancer in Australia 2000*, AIHW & AACR, 2003.

Table 8: Cancer of palate and other and unspecified parts of mouth (ICD-10 C05–C06)

Australia 2000

Age group	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
0–4	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
5–9	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
10–14	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
15–19	0	0.0	1	0.2	1	0.1	0	0.0	0	0.0	0	0.0
20–24	0	0.0	1	0.2	1	0.1	0	0.0	0	0.0	0	0.0
25–29	2	0.3	0	0.0	2	0.1	0	0.0	1	0.1	1	0.1
30–34	1	0.1	6	0.8	7	0.5	0	0.0	0	0.0	0	0.0
35–39	2	0.3	6	0.8	8	0.5	0	0.0	0	0.0	0	0.0
40–44	4	0.6	9	1.2	13	0.9	2	0.3	0	0.0	2	0.1
45–49	5	0.7	2	0.3	7	0.5	0	0.0	0	0.0	0	0.0
50–54	10	1.6	7	1.1	17	1.4	3	0.5	2	0.3	5	0.4
55–59	16	3.3	6	1.3	22	2.3	2	0.4	2	0.4	4	0.4
60–64	21	5.2	6	1.5	27	3.4	6	1.5	2	0.5	8	1.0
65–69	21	6.3	11	3.2	32	4.7	11	3.3	1	0.3	12	1.8
70–74	21	7.0	12	3.6	33	5.2	10	3.3	2	0.6	12	1.9
75–79	8	3.6	14	4.9	22	4.3	4	1.8	3	1.0	7	1.4
80–84	7	5.9	16	8.4	23	7.4	0	0.0	7	3.7	7	2.3
85 and over	6	7.7	11	6.3	17	6.7	3	3.9	6	3.4	9	3.6
Total	124		108		232		41		26		67	

Rates per 100,000 with 95% confidence intervals (95% CI)

Crude rate	1.3	1.1	1.2	0.4	0.3	0.3
95% CI	1.1 – 1.5	0.9 – 1.3	1.1 – 1.4	0.3 – 0.6	0.2 – 0.4	0.3 – 0.4
AS rate (Aust. 2001)	1.4	1.1	1.2	0.5	0.2	0.4
95% CI	1.2 – 1.6	0.9 – 1.3	1.1 – 1.4	0.3 – 0.6	0.1 – 0.3	0.3 – 0.4
AS rate (new WHO World)	1.1	0.8	0.9	0.3	0.2	0.2
95% CI	0.9 – 1.2	0.6 – 0.9	0.8 – 1.0	0.2 – 0.4	0.1 – 0.2	0.2 – 0.3
Lifetime risk (0–74)	1 in 788	1 in 1,413	1 in 1,016	1 in 2,150	1 in 8,795	1 in 3,515
PYLL (0–74)				350	165	515
Per cent of all cancers	0.3	0.3	0.3	0.2	0.2	0.2

Average annual numbers and rates by State and Territory, 1996–2000

	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate
NSW	45	1.5	33	1.0	78	1.2	12	0.4	10	0.3	21	0.3
Vic	29	1.4	27	1.1	57	1.2	7	0.3	6	0.2	13	0.3
Qld	25	1.6	17	1.0	42	1.3	4	0.3	4	0.2	8	0.3
WA	15	1.9	8	0.9	22	1.4	6	0.8	4	0.4	10	0.6
SA	9	1.3	9	1.0	18	1.2	4	0.5	3	0.3	6	0.4
Tas	4	1.7	4	1.6	8	1.7	1	0.4	1	0.5	2	0.4
ACT	2	1.9	2	1.5	4	1.7	0	0.2	1	0.6	1	0.4
NT	2	2.0	1	1.0	2	1.6	0	0.2	0	0.3	0	0.2

Notes

1. AS rates use Australian 2001 Standard Population unless new WHO World Standard Population is indicated. All rates are expressed per 100,000 population.

2. Incidence numbers are based on state of usual residence at time of diagnosis; mortality numbers on state of registration of death.

Source: *Cancer in Australia 2000*, AIHW & AACR, 2003.

Table 9: Cancer of the tonsil (ICD-10 C09)

Australia 2000

Age group	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
0-4	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
5-9	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
10-14	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
15-19	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
20-24	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
25-29	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
30-34	0	0.0	1	0.1	1	0.1	0	0.0	0	0.0	0	0.0
35-39	1	0.1	1	0.1	2	0.1	1	0.1	0	0.0	1	0.1
40-44	8	1.1	6	0.8	14	1.0	3	0.4	2	0.3	5	0.3
45-49	18	2.7	9	1.3	27	2.0	2	0.3	1	0.1	3	0.2
50-54	26	4.1	4	0.6	30	2.4	5	0.8	0	0.0	5	0.4
55-59	21	4.3	3	0.6	24	2.5	11	2.2	2	0.4	13	1.3
60-64	20	5.0	8	2.0	28	3.5	5	1.2	2	0.5	7	0.9
65-69	22	6.6	5	1.4	27	4.0	4	1.2	1	0.3	5	0.7
70-74	20	6.7	4	1.2	24	3.8	3	1.0	2	0.6	5	0.8
75-79	7	3.2	5	1.7	12	2.4	4	1.8	0	0.0	4	0.8
80-84	3	2.5	0	0.0	3	1.0	1	0.8	0	0.0	1	0.3
85 and over	1	1.3	2	1.1	3	1.2	1	1.3	0	0.0	1	0.4
Total	147		48		195		40		10		50	

Rates per 100,000 with 95% confidence intervals (95% CI)

Crude rate	1.5	0.5	1.0	0.4	0.1	0.3
95% CI	1.3 – 1.8	0.4 – 0.6	0.9 – 1.2	0.3 – 0.6	0.0 – 0.2	0.2 – 0.3
AS rate (Aust. 2001)	1.6	0.5	1.0	0.4	0.1	0.3
95% CI	1.3 – 1.8	0.4 – 0.6	0.9 – 1.2	0.3 – 0.6	0.0 – 0.2	0.2 – 0.3
AS rate (new WHO World)	1.3	0.4	0.8	0.3	0.1	0.2
95% CI	1.1 – 1.5	0.3 – 0.5	0.7 – 0.9	0.2 – 0.5	0.0 – 0.1	0.2 – 0.3
Lifetime risk (0-74)	1 in 654	1 in 2,390	1 in 1,034	1 in 2,727	1 in 8,936	1 in 4,178
PYLL (0-74)				595	165	760
Per cent of all cancers	0.3	0.1	0.2	0.2	0.1	0.1

Average annual numbers and rates by State and Territory, 1996-2000

	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate
NSW	45	1.5	15	0.5	61	1.0	13	0.4	3	0.1	16	0.3
Vic	37	1.7	12	0.5	48	1.1	9	0.4	3	0.1	12	0.3
Qld	27	1.7	8	0.5	35	1.1	8	0.5	2	0.1	10	0.3
WA	13	1.6	3	0.4	16	1.0	5	0.6	0	0.1	5	0.3
SA	11	1.5	5	0.6	16	1.0	3	0.4	2	0.2	4	0.3
Tas	3	1.4	1	0.3	4	0.8	1	0.5	0	0.0	1	0.3
ACT	1	1.0	0	0.3	2	0.6	1	0.7	0	0.0	1	0.3
NT	5	8.8	1	1.0	6	5.1	2	3.5	0	0.4	2	2.0

Notes

1. AS rates use Australian 2001 Standard Population unless new WHO World Standard Population is indicated. All rates are expressed per 100,000 population.
 2. Incidence numbers are based on state of usual residence at time of diagnosis; mortality numbers on state of registration of death.
- Source: *Cancer in Australia 2000*, AIHW & AACR, 2003.

Table 10: Cancer of the oropharynx (ICD-10 C10)

Australia 2000

Age group	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
0-4	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
5-9	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
10-14	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
15-19	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
20-24	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
25-29	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
30-34	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
35-39	4	0.5	0	0.0	4	0.3	0	0.0	0	0.0	0	0.0
40-44	1	0.1	0	0.0	1	0.1	1	0.1	1	0.1	2	0.1
45-49	8	1.2	1	0.1	9	0.7	3	0.4	1	0.1	4	0.3
50-54	11	1.7	0	0.0	11	0.9	9	1.4	1	0.2	10	0.8
55-59	5	1.0	1	0.2	6	0.6	6	1.2	1	0.2	7	0.7
60-64	13	3.2	0	0.0	13	1.6	8	2.0	1	0.3	9	1.1
65-69	10	3.0	1	0.3	11	1.6	15	4.5	1	0.3	16	2.4
70-74	7	2.3	3	0.9	10	1.6	6	2.0	7	2.1	13	2.1
75-79	2	0.9	4	1.4	6	1.2	5	2.3	2	0.7	7	1.4
80-84	0	0.0	4	2.1	4	1.3	5	4.2	1	0.5	6	1.9
85 and over	1	1.3	3	1.7	4	1.6	2	2.6	3	1.7	5	2.0
Total	62		17		79		60		19		79	

Rates per 100,000 with 95% confidence intervals (95% CI)

Crude rate	0.7	0.2	0.4	0.6	0.2	0.4
95% CI	0.5 – 0.8	0.1 – 0.3	0.3 – 0.5	0.5 – 0.8	0.1 – 0.3	0.3 – 0.5
AS rate (Aust. 2001)	0.7	0.2	0.4	0.7	0.2	0.4
95% CI	0.5 – 0.8	0.1 – 0.2	0.3 – 0.5	0.5 – 0.9	0.1 – 0.3	0.3 – 0.5
AS rate (new WHO World)	0.5	0.1	0.3	0.5	0.1	0.3
95% CI	0.4 – 0.7	0.0 – 0.1	0.3 – 0.4	0.4 – 0.6	0.1 – 0.2	0.2 – 0.4
Lifetime risk (0-74)	1 in 1,514	1 in 12,916	1 in 2,727	1 in 1,703	1 in 6,067	1 in 2,666
PYLL (0-74)				650	138	788
Per cent of all cancers	0.1	0.0	0.1	0.3	0.1	0.2

Average annual numbers and rates by State and Territory, 1996-2000

	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate
NSW	19	0.6	7	0.2	26	0.4	18	0.6	7	0.2	26	0.4
Vic	13	0.6	3	0.1	16	0.4	14	0.7	3	0.1	17	0.4
Qld	14	0.9	2	0.1	15	0.5	10	0.7	5	0.3	15	0.5
WA	4	0.6	1	0.1	5	0.3	4	0.6	1	0.1	5	0.3
SA	2	0.3	1	0.2	4	0.2	4	0.5	1	0.1	5	0.3
Tas	2	0.9	0	0.1	2	0.5	1	0.3	0	0.1	1	0.2
ACT	0	0.3	0	0.2	1	0.3	0	0.5	0	0.0	0	0.1
NT	1	1.5	0	0.7	1	1.1	1	0.9	0	0.0	1	0.5

Notes

1. AS rates use Australian 2001 Standard Population unless new WHO World Standard Population is indicated. All rates are expressed per 100,000 population.
 2. Incidence numbers are based on state of usual residence at time of diagnosis; mortality numbers on state of registration of death.
- Source: *Cancer in Australia 2000*, AIHW & AACR, 2003.

Table 11: Cancer of the tonsil and oropharynx (ICD-10 C09–C10)

Australia 2000

Age group	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
0–4	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
5–9	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
10–14	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
15–19	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
20–24	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
25–29	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
30–34	0	0.0	1	0.1	1	0.1	0	0.0	0	0.0	0	0.0
35–39	5	0.7	1	0.1	6	0.4	1	0.1	0	0.0	1	0.1
40–44	9	1.3	6	0.8	15	1.0	4	0.6	3	0.4	7	0.5
45–49	26	3.9	10	1.5	36	2.7	5	0.7	2	0.3	7	0.5
50–54	37	5.8	4	0.6	41	3.3	14	2.2	1	0.2	15	1.2
55–59	26	5.3	4	0.8	30	3.1	17	3.5	3	0.6	20	2.1
60–64	33	8.2	8	2.0	41	5.1	13	3.2	3	0.8	16	2.0
65–69	32	9.6	6	1.7	38	5.6	19	5.7	2	0.6	21	3.1
70–74	27	9.0	7	2.1	34	5.4	9	3.0	9	2.7	18	2.8
75–79	9	4.1	9	3.1	18	3.5	9	4.1	2	0.7	11	2.2
80–84	3	2.5	4	2.1	7	2.3	6	5.0	1	0.5	7	2.3
85 and over	2	2.6	5	2.9	7	2.8	3	3.9	3	1.7	6	2.4
Total	209		65		274		100		29		129	

Rates per 100,000 with 95% confidence intervals (95% CI)

Crude rate	2.2	0.7	1.4	1.1	0.3	0.7
95% CI	1.9 – 2.5	0.5 – 0.8	1.3 – 1.6	0.8 – 1.3	0.2 – 0.4	0.6 – 0.8
AS rate (Aust. 2001)	2.3	0.6	1.4	1.1	0.3	0.7
95% CI	2.0 – 2.6	0.5 – 0.8	1.3 – 1.6	0.9 – 1.3	0.2 – 0.4	0.6 – 0.8
AS rate (new WHO World)	1.8	0.5	1.1	0.9	0.2	0.5
95% CI	1.6 – 2.1	0.4 – 0.6	1.0 – 1.3	0.7 – 1.0	0.1 – 0.3	0.4 – 0.6
Lifetime risk (0–74)	1 in 457	1 in 2,017	1 in 750	1 in 1,049	1 in 3,614	1 in 1,628
PYLL (0–74)				1,245	303	1,548
Per cent of all cancers	0.5	0.2	0.3	0.5	0.2	0.4

Average annual numbers and rates by State and Territory, 1996–2000

	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate
NSW	65	2.2	22	0.7	86	1.4	31	1.1	11	0.3	42	0.7
Vic	50	2.3	15	0.6	65	1.4	23	1.1	6	0.2	29	0.6
Qld	41	2.6	10	0.6	50	1.6	18	1.2	7	0.4	25	0.8
WA	18	2.2	4	0.5	22	1.3	9	1.2	1	0.2	10	0.6
SA	14	1.9	6	0.8	20	1.3	7	0.9	3	0.3	9	0.6
Tas	5	2.3	1	0.4	6	1.3	2	0.8	0	0.1	2	0.4
ACT	2	1.3	1	0.4	2	0.9	1	1.2	0	0.0	1	0.5
NT	6	10.3	1	1.7	7	6.2	2	4.4	0	0.4	3	2.4

Notes

1. AS rates use Australian 2001 Standard Population unless new WHO World Standard Population is indicated. All rates are expressed per 100,000 population.
 2. Incidence numbers are based on state of usual residence at time of diagnosis; mortality numbers on state of registration of death.
- Source: *Cancer in Australia 2000*, AIHW & AACR, 2003.

Table 12: Cancer of nasopharynx (ICD-10 C11)

Australia 2000

Age group	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
0-4	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
5-9	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
10-14	1	0.1	0	0.0	1	0.1	0	0.0	0	0.0	0	0.0
15-19	2	0.3	0	0.0	2	0.2	0	0.0	1	0.2	1	0.1
20-24	2	0.3	1	0.2	3	0.2	0	0.0	0	0.0	0	0.0
25-29	0	0.0	1	0.1	1	0.1	0	0.0	0	0.0	0	0.0
30-34	4	0.6	3	0.4	7	0.5	0	0.0	1	0.1	1	0.1
35-39	6	0.8	2	0.3	8	0.5	0	0.0	0	0.0	0	0.0
40-44	6	0.8	0	0.0	6	0.4	0	0.0	0	0.0	0	0.0
45-49	7	1.0	2	0.3	9	0.7	10	1.5	2	0.3	12	0.9
50-54	11	1.7	9	1.4	20	1.6	6	0.9	2	0.3	8	0.6
55-59	9	1.8	2	0.4	11	1.1	6	1.2	2	0.4	8	0.8
60-64	9	2.2	5	1.3	14	1.8	3	0.7	0	0.0	3	0.4
65-69	5	1.5	4	1.2	9	1.3	5	1.5	3	0.9	8	1.2
70-74	7	2.3	4	1.2	11	1.7	3	1.0	5	1.5	8	1.3
75-79	5	2.3	1	0.3	6	1.2	6	2.7	2	0.7	8	1.6
80-84	2	1.7	2	1.1	4	1.3	4	3.4	6	3.2	10	3.2
85 and over	0	0.0	0	0.0	0	0.0	2	2.6	0	0.0	2	0.8
Total	76		36		112		45		24		69	

Rates per 100,000 with 95% confidence intervals (95% CI)

Crude rate	0.8	0.4	0.6	0.5	0.2	0.4
95% CI	0.6 – 1.0	0.3 – 0.5	0.5 – 0.7	0.3 – 0.6	0.1 – 0.3	0.3 – 0.4
AS rate (Aust. 2001)	0.8	0.4	0.6	0.5	0.2	0.4
95% CI	0.6 – 1.0	0.3 – 0.5	0.5 – 0.7	0.4 – 0.7	0.1 – 0.3	0.3 – 0.5
AS rate (new WHO World)	0.7	0.3	0.5	0.4	0.2	0.3
95% CI	0.5 – 0.8	0.2 – 0.4	0.4 – 0.6	0.3 – 0.5	0.1 – 0.2	0.2 – 0.3
Lifetime risk (0-74)	1 in 1,465	1 in 2,960	1 in 1,964	1 in 2,889	1 in 5,404	1 in 3,755
PYLL (0-74)				598	270	868
Per cent of all cancers	0.2	0.1	0.1	0.2	0.2	0.2

Average annual numbers and rates by State and Territory, 1996-2000

	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate
NSW	34	1.1	15	0.5	49	0.8	18	0.6	8	0.2	26	0.4
Vic	19	0.9	9	0.4	28	0.6	10	0.5	4	0.2	14	0.3
Qld	12	0.7	4	0.2	16	0.5	9	0.6	4	0.2	13	0.4
WA	5	0.6	2	0.2	7	0.4	4	0.5	2	0.2	6	0.4
SA	4	0.5	1	0.2	5	0.3	3	0.4	1	0.1	4	0.2
Tas	2	0.7	1	0.2	2	0.5	0	0.2	1	0.2	1	0.2
ACT	1	1.0	1	0.4	2	0.6	0	0.3	0	0.2	1	0.2
NT	1	2.5	0	1.0	2	1.7	1	0.9	0	0.5	1	0.7

Notes

1. AS rates use Australian 2001 Standard Population unless new WHO World Standard Population is indicated. All rates are expressed per 100,000 population.
 2. Incidence numbers are based on state of usual residence at time of diagnosis; mortality numbers on state of registration of death.
- Source: *Cancer in Australia 2000*, AIHW & AACR, 2003.

Table 13: Cancers of the hypopharynx (ICD-10 C12–C13)

Australia 2000

Age group	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
0–4	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
5–9	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
10–14	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
15–19	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
20–24	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
25–29	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
30–34	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
35–39	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
40–44	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
45–49	8	1.2	1	0.1	9	0.7	3	0.4	0	0.0	3	0.2
50–54	16	2.5	6	1.0	22	1.7	3	0.5	1	0.2	4	0.3
55–59	26	5.3	4	0.8	30	3.1	4	0.8	0	0.0	4	0.4
60–64	21	5.2	2	0.5	23	2.9	9	2.2	1	0.3	10	1.3
65–69	20	6.0	4	1.2	24	3.5	9	2.7	0	0.0	9	1.3
70–74	26	8.7	4	1.2	30	4.7	9	3.0	1	0.3	10	1.6
75–79	19	8.7	4	1.4	23	4.5	4	1.8	1	0.3	5	1.0
80–84	6	5.0	2	1.1	8	2.6	4	3.4	0	0.0	4	1.3
85 and over	6	7.7	3	1.7	9	3.6	3	3.9	0	0.0	3	1.2
Total	148		30		178		48		4		52	

Rates per 100,000 with 95% confidence intervals (95% CI)

Crude rate	1.6	0.3	0.9	0.5	0.0	0.3
95% CI	1.3 – 1.8	0.2 – 0.4	0.8 – 1.1	0.4 – 0.6	0.0 – 0.1	0.2 – 0.3
AS rate (Aust. 2001)	1.7	0.3	0.9	0.6	0.0	0.3
95% CI	1.4 – 1.9	0.2 – 0.4	0.8 – 1.1	0.4 – 0.7	0.0 – 0.1	0.2 – 0.3
AS rate (new WHO World)	1.2	0.2	0.7	0.4	0.0	0.2
95% CI	1.0 – 1.4	0.1 – 0.3	0.6 – 0.8	0.3 – 0.5	0.0 – 0.1	0.1 – 0.3
Lifetime risk (0–74)	1 in 691	1 in 4,152	1 in 1,198	1 in 2,063	1 in 28,083	1 in 3,908
PYLL (0–74)				423	38	460
Per cent of all cancers	0.3	0.1	0.2	0.2	0.0	0.1

Average annual numbers and rates by State and Territory, 1996–2000

	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate
NSW	52	1.8	10	0.3	62	1.0	19	0.6	4	0.1	23	0.4
Vic	33	1.6	7	0.3	40	0.9	12	0.6	3	0.1	15	0.3
Qld	31	2.1	5	0.3	36	1.1	11	0.7	1	0.1	12	0.4
WA	14	1.8	2	0.2	15	0.9	7	0.9	1	0.1	7	0.5
SA	7	1.0	1	0.1	8	0.5	5	0.6	1	0.1	5	0.3
Tas	3	1.2	1	0.5	4	0.8	2	0.7	0	0.2	2	0.4
ACT	2	1.3	0	0.1	2	0.7	1	0.7	0	0.0	1	0.3
NT	3	5.5	0	0.0	3	3.0	1	2.7	0	0.0	1	1.3

Notes

1. AS rates use Australian 2001 Standard Population unless new WHO World Standard Population is indicated. All rates are expressed per 100,000 population.

2. Incidence numbers are based on state of usual residence at time of diagnosis; mortality numbers on state of registration of death.

Source: *Cancer in Australia 2000*, AIHW & AACR, 2003.

Table 14: Cancer of other and ill-defined sites in the lip, oral cavity and pharynx (ICD-10 C14)

Australia 2000

Age group	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
0-4	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
5-9	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
10-14	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
15-19	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
20-24	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
25-29	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
30-34	0	0.0	0	0.0	0	0.0	0	0.0	1	0.1	1	0.1
35-39	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
40-44	4	0.6	0	0.0	4	0.3	0	0.0	0	0.0	0	0.0
45-49	3	0.4	2	0.3	5	0.4	3	0.4	0	0.0	3	0.2
50-54	6	0.9	4	0.6	10	0.8	3	0.5	1	0.2	4	0.3
55-59	10	2.0	0	0.0	10	1.0	2	0.4	1	0.2	3	0.3
60-64	13	3.2	1	0.3	14	1.8	4	1.0	1	0.3	5	0.6
65-69	9	2.7	3	0.9	12	1.8	2	0.6	2	0.6	4	0.6
70-74	3	1.0	2	0.6	5	0.8	3	1.0	1	0.3	4	0.6
75-79	11	5.0	5	1.7	16	3.2	8	3.6	1	0.3	9	1.8
80-84	3	2.5	1	0.5	4	1.3	0	0.0	0	0.0	0	0.0
85 and over	0	0.0	2	1.1	2	0.8	4	5.2	1	0.6	5	2.0
Total	62		20		82		29		9		38	

Rates per 100,000 with 95% confidence intervals (95% CI)

Crude rate	0.7	0.2	0.4	0.3	0.1	0.2
95% CI	0.5 – 0.8	0.1 – 0.3	0.3 – 0.5	0.2 – 0.4	0.0 – 0.2	0.1 – 0.3
AS rate (Aust. 2001)	0.7	0.2	0.4	0.3	0.1	0.2
95% CI	0.5 – 0.9	0.1 – 0.3	0.3 – 0.5	0.2 – 0.5	0.0 – 0.1	0.1 – 0.3
AS rate (new WHO World)	0.5	0.1	0.3	0.2	0.1	0.1
95% CI	0.4 – 0.7	0.1 – 0.2	0.3 – 0.4	0.1 – 0.3	0.0 – 0.1	0.1 – 0.2
Lifetime risk (0-74)	1 in 1,828	1 in 7,521	1 in 2,942	1 in 5,087	1 in 12,179	1 in 7,214
PYLL (0-74)				258	113	370
Per cent of all cancers	0.1	0.1	0.1	0.1	0.1	0.1

Average annual numbers and rates by State and Territory, 1996-2000

	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate
NSW	17	0.6	4	0.1	22	0.3	9	0.3	2	0.1	11	0.2
Vic	13	0.6	4	0.1	16	0.4	7	0.3	3	0.1	9	0.2
Qld	13	0.9	5	0.3	18	0.6	8	0.5	2	0.1	9	0.3
WA	6	0.9	1	0.1	7	0.4	3	0.5	1	0.1	4	0.3
SA	2	0.2	1	0.1	2	0.1	2	0.3	1	0.1	3	0.2
Tas	1	0.4	1	0.5	2	0.5	1	0.4	1	0.5	2	0.5
ACT	0	0.5	0	0.2	1	0.3	0	1.0	0	0.2	1	0.4
NT	0	1.3	0	0.0	0	0.7	0	0.6	0	0.0	0	0.4

Notes

- AS rates use Australian 2001 Standard Population unless new WHO World Standard Population is indicated. All rates are expressed per 100,000 population.
 - Incidence numbers are based on state of usual residence at time of diagnosis; mortality numbers on state of registration of death.
- Source: *Cancer in Australia 2000*, AIHW & AACR, 2003.

Table 15: Cancer of the head and neck (ICD-10 C01–C14)

Australia 2000

Age group	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
0–4	1	0.2	0	0.0	1	0.1	0	0.0	0	0.0	0	0.0
5–9	1	0.1	0	0.0	1	0.1	0	0.0	0	0.0	0	0.0
10–14	2	0.3	1	0.2	3	0.2	0	0.0	0	0.0	0	0.0
15–19	5	0.7	2	0.3	7	0.5	1	0.1	1	0.2	2	0.2
20–24	10	1.5	7	1.1	17	1.3	0	0.0	1	0.2	1	0.1
25–29	3	0.4	10	1.4	13	0.9	0	0.0	1	0.1	1	0.1
30–34	9	1.3	19	2.6	28	2.0	2	0.3	2	0.3	4	0.3
35–39	31	4.1	16	2.1	47	3.1	4	0.5	3	0.4	7	0.5
40–44	53	7.4	35	4.8	88	6.1	8	1.1	7	1.0	15	1.0
45–49	100	15.0	37	5.5	137	10.2	29	4.3	9	1.3	38	2.8
50–54	137	21.6	66	10.6	203	16.1	43	6.8	16	2.6	59	4.7
55–59	162	33.0	40	8.4	202	21.0	56	11.4	14	3.0	70	7.3
60–64	183	45.7	52	13.1	235	29.5	53	13.2	13	3.3	66	8.3
65–69	164	49.4	56	16.2	220	32.5	70	21.1	23	6.7	93	13.7
70–74	159	53.1	61	18.3	220	34.7	75	25.0	29	8.7	104	16.4
75–79	108	49.2	71	24.7	179	35.3	51	23.2	19	6.6	70	13.8
80–84	59	49.6	53	27.9	112	36.2	29	24.4	29	15.3	58	18.8
85 and over	35	45.2	58	33.1	93	36.8	33	42.6	30	17.1	63	24.9
Total	1,222		584		1,806		454		197		651	

Rates per 100,000 with 95% confidence intervals (95% CI)

Crude rate	12.9	6.1	9.4	4.8	2.0	3.4
95% CI	12.1 – 13.6	5.6 – 6.5	9.0 – 9.9	4.3 – 5.2	1.8 – 2.3	3.1 – 3.7
AS rate (Aust. 2001)	13.6	5.8	9.5	5.3	1.9	3.4
95% CI	12.8 – 14.4	5.3 – 6.3	9.1 – 10.0	4.8 – 5.7	1.6 – 2.2	3.2 – 3.7
AS rate (new WHO World)	10.5	4.4	7.4	3.8	1.4	2.5
95% CI	9.9 – 11.1	4.0 – 4.8	7.0 – 7.7	3.4 – 4.1	1.2 – 1.6	2.3 – 2.7
Lifetime risk (0–74)	1 in 86	1 in 237	1 in 127	1 in 239	1 in 726	1 in 362
PYLL (0–74)				4,673	1,843	6,515
Per cent of all cancers	2.7	1.5	2.1	2.3	1.3	1.8

Average annual numbers and rates by State and Territory, 1996–2000

	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate
NSW	420	14.3	193	5.7	613	9.8	152	5.3	67	2.0	220	3.6
Vic	269	12.6	139	5.7	408	8.9	107	5.2	46	1.8	153	3.4
Qld	242	15.6	103	6.2	345	10.8	84	5.8	34	2.0	118	3.8
WA	109	14.1	42	4.9	151	9.3	48	6.7	16	1.9	63	4.1
SA	75	10.4	43	5.0	118	7.5	36	5.1	16	1.8	52	3.3
Tas	28	12.5	15	5.8	43	9.0	11	5.1	5	1.9	16	3.4
ACT	16	14.4	7	4.9	22	9.2	5	5.8	2	1.7	7	3.4
NT	17	30.0	4	6.9	21	19.3	6	11.8	2	2.8	8	7.5

Notes

1. AS rates use Australian 2001 Standard Population unless new WHO World Standard Population is indicated. All rates are expressed per 100,000 population.

2. Incidence numbers are based on state of usual residence at time of diagnosis; mortality numbers on state of registration of death.

Source: *Cancer in Australia 2000*, AIHW & AACR, 2003.

Table 16: Cancer of the oesophagus (ICD-10 C15)

Australia 2000

Age group	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
0-4	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
5-9	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
10-14	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
15-19	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
20-24	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
25-29	1	0.1	0	0.0	1	0.1	2	0.3	1	0.1	3	0.2
30-34	0	0.0	1	0.1	1	0.1	0	0.0	1	0.1	1	0.1
35-39	8	1.1	3	0.4	11	0.7	4	0.5	2	0.3	6	0.4
40-44	19	2.6	1	0.1	20	1.4	13	1.8	2	0.3	15	1.0
45-49	27	4.0	8	1.2	35	2.6	20	3.0	3	0.4	23	1.7
50-54	65	10.2	11	1.8	76	6.0	47	7.4	7	1.1	54	4.3
55-59	65	13.3	21	4.4	86	8.9	54	11.0	15	3.2	69	7.2
60-64	85	21.2	24	6.0	109	13.7	80	20.0	16	4.0	96	12.0
65-69	99	29.8	33	9.6	132	19.5	98	29.5	22	6.4	120	17.7
70-74	120	40.1	50	15.0	170	26.8	108	36.0	35	10.5	143	22.6
75-79	109	49.6	80	27.8	189	37.3	127	57.8	49	17.0	176	34.7
80-84	67	56.3	48	25.3	115	37.2	64	53.8	46	24.2	110	35.6
85 and over	46	59.3	74	42.2	120	47.5	66	85.1	89	50.8	155	61.3
Total	711		354		1,065		683		288		971	

Rates per 100,000 with 95% confidence intervals (95% CI)

Crude rate	7.5	3.7	5.6	7.2	3.0	5.1
95% CI	6.9 – 8.0	3.3 – 4.1	5.2 – 5.9	6.6 – 7.7	2.6 – 3.3	4.8 – 5.4
AS rate (Aust. 2001)	8.3	3.3	5.6	8.2	2.6	5.2
95% CI	7.7 – 8.9	3.0 – 3.7	5.3 – 6.0	7.5 – 8.8	2.3 – 2.9	4.8 – 5.5
AS rate (new WHO World)	5.8	2.2	3.9	5.5	1.6	3.5
95% CI	5.4 – 6.3	1.9 – 2.4	3.7 – 4.1	5.1 – 6.0	1.4 – 1.8	3.2 – 3.7
Lifetime risk (0-74)	1 in 164	1 in 518	1 in 251	1 in 183	1 in 757	1 in 298
PYLL (0-74)				5,225	1,185	6,410
Per cent of all cancers	1.5	0.9	1.2	3.4	1.9	2.7

Average annual numbers and rates by State and Territory, 1996-2000

	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate
NSW	208	7.4	124	3.5	332	5.3	213	7.7	107	2.9	320	5.1
Vic	168	8.4	100	3.8	268	5.8	173	8.8	94	3.5	267	5.8
Qld	126	8.8	58	3.4	183	5.9	123	8.7	45	2.6	167	5.4
WA	63	9.0	30	3.5	93	6.0	52	7.5	24	2.7	75	4.9
SA	62	8.8	32	3.5	94	5.9	55	7.8	23	2.5	79	4.9
Tas	23	10.3	12	4.4	35	7.2	20	9.8	8	3.0	28	6.0
ACT	8	8.8	4	3.3	11	5.5	9	9.6	3	2.9	12	5.7
NT	5	12.9	1	3.0	6	7.9	3	7.9	1	2.5	3	5.0

Notes

- AS rates use Australian 2001 Standard Population unless new WHO World Standard Population is indicated. All rates are expressed per 100,000 population.
 - Incidence numbers are based on state of usual residence at time of diagnosis; mortality numbers on state of registration of death.
- Source: *Cancer in Australia 2000*, AIHW & AACR, 2003.

Table 17: Cancer of the stomach (ICD-10 C16)

Australia 2000

Age group	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
0-4	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
5-9	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
10-14	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
15-19	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
20-24	1	0.2	0	0.0	1	0.1	2	0.3	2	0.3	4	0.3
25-29	4	0.6	2	0.3	6	0.4	2	0.3	0	0.0	2	0.1
30-34	8	1.1	7	1.0	15	1.1	0	0.0	1	0.1	1	0.1
35-39	10	1.3	9	1.2	19	1.3	3	0.4	5	0.7	8	0.5
40-44	24	3.3	20	2.7	44	3.0	16	2.2	11	1.5	27	1.9
45-49	46	6.9	34	5.0	80	6.0	20	3.0	9	1.3	29	2.2
50-54	84	13.2	43	6.9	127	10.1	43	6.8	24	3.9	67	5.3
55-59	99	20.2	44	9.3	143	14.8	47	9.6	17	3.6	64	6.6
60-64	130	32.4	59	14.9	189	23.7	58	14.5	25	6.3	83	10.4
65-69	168	50.6	74	21.4	242	35.7	104	31.3	37	10.7	141	20.8
70-74	221	73.8	85	25.5	306	48.3	145	48.4	44	13.2	189	29.8
75-79	225	102.5	134	46.6	359	70.8	134	61.0	88	30.6	222	43.8
80-84	134	112.6	103	54.2	237	76.7	106	89.1	59	31.1	165	53.4
85 and over	113	145.8	99	56.5	212	83.9	83	107.1	104	59.4	187	74.0
Total	1,267		713		1,980		763		426		1,189	

Rates per 100,000 with 95% confidence intervals (95% CI)

Crude rate	13.3	7.4	10.3	8.0	4.4	6.2
95% CI	12.6 – 14.1	6.8 – 7.9	9.9 – 10.8	7.5 – 8.6	4.0 – 4.8	5.9 – 6.6
AS rate (Aust. 2001)	15.1	6.9	10.5	9.3	4.0	6.3
95% CI	14.3 – 16.0	6.3 – 7.4	10.0 – 11.0	8.6 – 10.0	3.6 – 4.3	6.0 – 6.7
AS rate (new WHO World)	10.3	4.7	7.3	6.1	2.6	4.2
95% CI	9.7 – 10.9	4.4 – 5.1	6.9 – 7.6	5.7 – 6.5	2.3 – 2.8	3.9 – 4.4
Lifetime risk (0-74)	1 in 99	1 in 227	1 in 139	1 in 172	1 in 481	1 in 257
PYLL (0-74)				5,040	2,478	7,518
Per cent of all cancers	2.8	1.8	2.3	3.8	2.8	3.4

Average annual numbers and rates by State and Territory, 1996-2000

	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate
NSW	418	15.1	231	6.6	649	10.4	239	8.9	138	3.9	377	6.1
Vic	345	17.0	194	7.5	538	11.7	213	10.8	128	4.9	341	7.5
Qld	213	15.3	111	6.6	324	10.5	127	9.5	68	3.9	195	6.4
WA	104	14.8	61	7.0	164	10.5	71	10.5	44	5.1	115	7.5
SA	112	15.8	63	6.9	175	10.8	74	10.6	49	5.4	124	7.6
Tas	31	14.9	18	6.9	50	10.4	24	11.6	14	5.2	38	7.9
ACT	13	13.8	9	8.0	22	10.6	9	10.0	8	7.3	17	8.7
NT	4	12.5	1	3.8	6	8.2	2	5.5	1	2.4	3	4.0

Notes

1. AS rates use Australian 2001 Standard Population unless new WHO World Standard Population is indicated. All rates are expressed per 100,000 population.

2. Incidence numbers are based on state of usual residence at time of diagnosis; mortality numbers on state of registration of death.

Source: *Cancer in Australia 2000*, AIHW & AACR, 2003.

Table 18: Cancer of the small intestine (ICD-10 C17)

Australia 2000

Age group	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
0-4	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
5-9	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
10-14	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
15-19	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
20-24	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
25-29	1	0.1	1	0.1	2	0.1	1	0.1	0	0.0	1	0.1
30-34	1	0.1	1	0.1	2	0.1	1	0.1	1	0.1	2	0.1
35-39	4	0.5	2	0.3	6	0.4	0	0.0	1	0.1	1	0.1
40-44	8	1.1	3	0.4	11	0.8	2	0.3	1	0.1	3	0.2
45-49	12	1.8	9	1.3	21	1.6	2	0.3	1	0.1	3	0.2
50-54	13	2.0	15	2.4	28	2.2	3	0.5	2	0.3	5	0.4
55-59	13	2.7	11	2.3	24	2.5	6	1.2	0	0.0	6	0.6
60-64	14	3.5	8	2.0	22	2.8	5	1.2	3	0.8	8	1.0
65-69	26	7.8	15	4.3	41	6.1	11	3.3	4	1.2	15	2.2
70-74	22	7.3	15	4.5	37	5.8	10	3.3	5	1.5	15	2.4
75-79	16	7.3	18	6.3	34	6.7	9	4.1	6	2.1	15	3.0
80-84	4	3.4	6	3.2	10	3.2	1	0.8	6	3.2	7	2.3
85 and over	8	10.3	12	6.9	20	7.9	6	7.7	6	3.4	12	4.7
Total	142		116		258		57		36		93	

Rates per 100,000 with 95% confidence intervals (95% CI)

Crude rate	1.5	1.2	1.3	0.6	0.4	0.5
95% CI	1.2 – 1.7	1.0 – 1.4	1.2 – 1.5	0.4 – 0.8	0.3 – 0.5	0.4 – 0.6
AS rate (Aust. 2001)	1.6	1.1	1.4	0.7	0.3	0.5
95% CI	1.3 – 1.9	0.9 – 1.4	1.2 – 1.5	0.5 – 0.8	0.2 – 0.5	0.4 – 0.6
AS rate (new WHO World)	1.2	0.9	1.0	0.5	0.2	0.3
95% CI	1.0 – 1.4	0.7 – 1.0	0.9 – 1.1	0.4 – 0.6	0.2 – 0.3	0.3 – 0.4
Lifetime risk (0-74)	1 in 739	1 in 1,119	1 in 894	1 in 1,914	1 in 4,661	1 in 2,735
PYLL (0-74)				553	265	818
Per cent of all cancers	0.3	0.3	0.3	0.3	0.2	0.3

Average annual numbers and rates by State and Territory, 1996-2000

	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate
NSW	44	1.5	34	1.0	78	1.2	17	0.6	12	0.3	29	0.5
Vic	28	1.4	27	1.1	55	1.2	10	0.5	7	0.3	17	0.4
Qld	29	1.9	20	1.2	49	1.5	9	0.6	8	0.5	17	0.6
WA	13	1.7	10	1.2	23	1.5	5	0.6	4	0.4	9	0.5
SA	11	1.5	7	0.8	17	1.1	6	0.8	2	0.3	8	0.5
Tas	4	1.6	3	1.2	7	1.4	1	0.6	2	0.7	3	0.7
ACT	1	0.9	1	1.1	3	1.0	1	0.9	0	0.2	1	0.5
NT	0	0.0	1	1.1	1	0.5	0	0.4	0	0.0	0	0.2

Notes

- AS rates use Australian 2001 Standard Population unless new WHO World Standard Population is indicated. All rates are expressed per 100,000 population.
 - Incidence numbers are based on state of usual residence at time of diagnosis; mortality numbers on state of registration of death.
- Source: *Cancer in Australia 2000*, AIHW & AACR, 2003.

Table 19: Cancer of the colon (ICD-10 C18)

Australia 2000

Age group	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
0-4	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
5-9	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
10-14	2	0.3	1	0.2	3	0.2	0	0.0	0	0.0	0	0.0
15-19	1	0.1	4	0.6	5	0.4	0	0.0	0	0.0	0	0.0
20-24	3	0.5	6	0.9	9	0.7	1	0.2	3	0.5	4	0.3
25-29	10	1.4	9	1.2	19	1.3	1	0.1	0	0.0	1	0.1
30-34	17	2.4	15	2.1	32	2.2	8	1.1	7	1.0	15	1.1
35-39	26	3.5	38	5.0	64	4.3	11	1.5	11	1.5	22	1.5
40-44	79	11.0	66	9.1	145	10.0	23	3.2	15	2.1	38	2.6
45-49	118	17.7	119	17.7	237	17.7	42	6.3	45	6.7	87	6.5
50-54	208	32.8	210	33.7	418	33.2	81	12.8	49	7.9	130	10.3
55-59	360	73.4	286	60.4	646	67.0	145	29.6	101	21.3	246	25.5
60-64	519	129.5	387	97.5	906	113.6	209	52.1	135	34.0	344	43.1
65-69	615	185.2	455	131.9	1,070	158.0	246	74.1	159	46.1	405	59.8
70-74	770	257.0	599	179.5	1,369	216.2	311	103.8	222	66.5	533	84.2
75-79	666	303.3	637	221.4	1,303	256.8	293	133.4	285	99.0	578	113.9
80-84	427	358.9	483	254.2	910	294.5	231	194.2	244	128.4	475	153.7
85 and over	320	412.8	515	294.0	835	330.5	184	237.4	373	213.0	557	220.4
Total	4,141		3,830		7,971		1,786		1,649		3,435	

Rates per 100,000 with 95% confidence intervals (95% CI)

Crude rate	43.6	39.7	41.6	18.8	17.1	17.9
95% CI	42.2 – 44.9	38.4 – 41.0	40.7 – 42.5	17.9 – 19.7	16.3 – 17.9	17.3 – 18.5
AS rate (Aust. 2001)	49.0	37.0	42.3	21.6	15.5	18.3
95% CI	47.5 – 50.5	35.8 – 38.2	41.3 – 43.2	20.6 – 22.7	14.7 – 16.2	17.7 – 18.9
AS rate (new WHO World)	33.8	25.9	29.5	14.5	10.2	12.2
95% CI	32.8 – 34.8	25.0 – 26.8	28.8 – 30.1	13.8 – 15.1	9.7 – 10.7	11.7 – 12.6
Lifetime risk (0-74)	1 in 28	1 in 38	1 in 33	1 in 71	1 in 107	1 in 86
PYLL (0-74)				12,350	8,898	21,248
Per cent of all cancers	9.0	9.7	9.4	8.9	10.7	9.7

Average annual numbers and rates by State and Territory, 1996-2000

	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate
NSW	1,305	46.6	1,215	35.1	2,519	40.2	616	22.8	548	15.5	1,164	18.6
Vic	979	48.4	958	37.6	1,938	42.3	480	24.2	450	17.3	930	20.4
Qld	711	49.6	680	40.4	1,391	44.6	336	24.3	287	16.9	623	20.2
WA	316	45.2	284	33.4	600	38.5	144	21.1	132	15.3	276	17.9
SA	339	48.0	349	38.7	687	42.7	145	20.8	147	16.0	292	18.1
Tas	96	44.1	107	40.4	203	42.2	48	22.9	54	19.8	102	21.4
ACT	48	50.3	43	36.4	91	42.4	25	28.1	23	20.2	48	24.0
NT	15	33.5	10	33.1	25	34.1	4	11.9	5	15.3	9	13.9

Notes

- AS rates use Australian 2001 Standard Population unless new WHO World Standard Population is indicated. All rates are expressed per 100,000 population.
 - Incidence numbers are based on state of usual residence at time of diagnosis; mortality numbers on state of registration of death.
- Source: *Cancer in Australia 2000*, AIHW & AACR, 2003.

Table 20: Cancer of the rectum (ICD-10 C19–C20)

Australia 2000

Age group	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
0–4	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
5–9	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
10–14	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
15–19	1	0.1	0	0.0	1	0.1	0	0.0	0	0.0	0	0.0
20–24	0	0.0	1	0.2	1	0.1	0	0.0	0	0.0	0	0.0
25–29	6	0.8	3	0.4	9	0.6	1	0.1	1	0.1	2	0.1
30–34	15	2.1	13	1.8	28	2.0	2	0.3	2	0.3	4	0.3
35–39	29	3.9	17	2.2	46	3.1	10	1.3	2	0.3	12	0.8
40–44	48	6.7	43	5.9	91	6.3	10	1.4	4	0.5	14	1.0
45–49	113	16.9	87	12.9	200	14.9	19	2.8	13	1.9	32	2.4
50–54	218	34.4	126	20.2	344	27.4	45	7.1	27	4.3	72	5.7
55–59	269	54.9	134	28.3	403	41.8	50	10.2	37	7.8	87	9.0
60–64	344	85.8	165	41.6	509	63.8	95	23.7	40	10.1	135	16.9
65–69	380	114.4	182	52.7	562	83.0	111	33.4	44	12.8	155	22.9
70–74	473	157.9	235	70.4	708	111.8	135	45.1	64	19.2	199	31.4
75–79	410	186.7	246	85.5	656	129.3	131	59.7	74	25.7	205	40.4
80–84	198	166.4	185	97.4	383	124.0	81	68.1	63	33.2	144	46.6
85 and over	114	147.1	159	90.8	273	108.0	67	86.4	96	54.8	163	64.5
Total	2,618		1,596		4,214		757		467		1,224	

Rates per 100,000 with 95% confidence intervals (95% CI)

Crude rate	27.5	16.5	22.0	8.0	4.8	6.4
95% CI	26.5 – 28.6	15.7 – 17.4	21.3 – 22.7	7.4 – 8.5	4.4 – 5.3	6.0 – 6.7
AS rate (Aust. 2001)	30.1	15.6	22.3	9.0	4.4	6.5
95% CI	28.9 – 31.2	14.9 – 16.4	21.6 – 23.0	8.4 – 9.7	4.0 – 4.8	6.1 – 6.9
AS rate (new WHO World)	21.6	11.3	16.2	6.1	3.0	4.4
95% CI	20.8 – 22.5	10.7 – 11.9	15.7 – 16.7	5.7 – 6.6	2.7 – 3.3	4.2 – 4.7
Lifetime risk (0–74)	1 in 42	1 in 85	1 in 57	1 in 160	1 in 349	1 in 221
PYLL (0–74)				5,600	2,940	8,540
Per cent of all cancers	5.7	4.1	4.9	3.8	3.0	3.5

Average annual numbers and rates by State and Territory, 1996–2000

	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate
NSW	824	28.7	510	15.1	1,334	21.3	214	7.8	139	4.0	353	5.6
Vic	637	30.7	415	16.5	1,052	23.0	181	9.2	128	4.9	310	6.8
Qld	419	28.1	250	15.0	669	21.2	109	7.7	68	4.0	177	5.7
WA	220	30.4	125	14.8	345	21.8	84	12.0	47	5.4	130	8.4
SA	221	30.6	148	16.8	369	23.1	71	10.2	52	5.7	123	7.6
Tas	63	28.8	44	17.1	107	22.4	22	10.6	20	7.7	42	8.9
ACT	26	23.8	17	14.2	42	19.1	5	4.8	5	4.2	10	4.7
NT	9	19.9	5	9.4	14	14.8	4	9.4	1	3.1	5	6.5

Notes

1. AS rates use Australian 2001 Standard Population unless new WHO World Standard Population is indicated. All rates are expressed per 100,000 population.

2. Incidence numbers are based on state of usual residence at time of diagnosis; mortality numbers on state of registration of death.

Source: *Cancer in Australia 2000*, AIHW & AACR, 2003.

Table 21: Cancer of the anus (ICD-10 C21)

Australia 2000

Age group	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
0-4	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
5-9	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
10-14	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
15-19	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
20-24	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
25-29	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
30-34	1	0.1	0	0.0	1	0.1	0	0.0	0	0.0	0	0.0
35-39	3	0.4	0	0.0	3	0.2	1	0.1	0	0.0	1	0.1
40-44	7	1.0	6	0.8	13	0.9	0	0.0	1	0.1	1	0.1
45-49	6	0.9	10	1.5	16	1.2	0	0.0	3	0.4	3	0.2
50-54	6	0.9	13	2.1	19	1.5	2	0.3	2	0.3	4	0.3
55-59	13	2.7	9	1.9	22	2.3	2	0.4	4	0.8	6	0.6
60-64	21	5.2	12	3.0	33	4.1	2	0.5	0	0.0	2	0.3
65-69	7	2.1	16	4.6	23	3.4	4	1.2	2	0.6	6	0.9
70-74	20	6.7	18	5.4	38	6.0	5	1.7	3	0.9	8	1.3
75-79	11	5.0	12	4.2	23	4.5	5	2.3	5	1.7	10	2.0
80-84	5	4.2	11	5.8	16	5.2	4	3.4	3	1.6	7	2.3
85 and over	4	5.2	9	5.1	13	5.1	1	1.3	10	5.7	11	4.4
Total	104		116		220		26		33		59	

Rates per 100,000 with 95% confidence intervals (95% CI)

Crude rate	1.1	1.2	1.1	0.3	0.3	0.3
95% CI	0.9 – 1.3	1.0 – 1.4	1.0 – 1.3	0.2 – 0.4	0.2 – 0.5	0.2 – 0.4
AS rate (Aust. 2001)	1.2	1.2	1.2	0.3	0.3	0.3
95% CI	0.9 – 1.4	0.9 – 1.4	1.0 – 1.3	0.2 – 0.4	0.2 – 0.4	0.2 – 0.4
AS rate (new WHO World)	0.9	0.9	0.9	0.2	0.2	0.2
95% CI	0.7 – 1.0	0.7 – 1.0	0.8 – 1.0	0.1 – 0.3	0.1 – 0.3	0.2 – 0.3
Lifetime risk (0-74)	1 in 999	1 in 1,034	1 in 1,016	1 in 4,729	1 in 6,199	1 in 5,406
PYLL (0-74)				185	253	438
Per cent of all cancers	0.2	0.3	0.3	0.1	0.2	0.2

Average annual numbers and rates by State and Territory, 1996-2000

	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate
NSW	45	1.6	55	1.6	100	1.6	8	0.3	11	0.3	20	0.3
Vic	17	0.8	17	0.7	34	0.7	5	0.2	7	0.3	12	0.3
Qld	18	1.2	21	1.2	39	1.2	3	0.2	3	0.2	6	0.2
WA	9	1.2	10	1.2	20	1.2	2	0.2	3	0.3	5	0.3
SA	4	0.5	10	1.1	13	0.8	1	0.1	2	0.2	3	0.2
Tas	4	1.6	3	1.2	7	1.4	1	0.5	1	0.3	2	0.4
ACT	1	1.4	2	1.5	3	1.5	1	0.6	0	0.0	1	0.3
NT	1	1.3	0	0.0	1	0.7	0	0.1	0	0.0	0	0.1

Notes

1. AS rates use Australian 2001 Standard Population unless new WHO World Standard Population is indicated. All rates are expressed per 100,000 population.
 2. Incidence numbers are based on state of usual residence at time of diagnosis; mortality numbers on state of registration of death.
- Source: *Cancer in Australia 2000*, AIHW & AACR, 2003.

Table 22: Cancer of the colon and rectum (including anus) (ICD-10 C18–C21)

Australia 2000

Age group	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
0–4	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
5–9	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
10–14	2	0.3	1	0.2	3	0.2	0	0.0	0	0.0	0	0.0
15–19	2	0.3	4	0.6	6	0.5	0	0.0	0	0.0	0	0.0
20–24	3	0.5	7	1.1	10	0.8	1	0.2	3	0.5	4	0.3
25–29	16	2.2	12	1.7	28	1.9	2	0.3	1	0.1	3	0.2
30–34	33	4.7	28	3.9	61	4.3	10	1.4	9	1.3	19	1.3
35–39	58	7.8	55	7.3	113	7.5	22	2.9	13	1.7	35	2.3
40–44	134	18.6	115	15.8	249	17.2	33	4.6	20	2.7	53	3.7
45–49	237	35.5	216	32.0	453	33.8	61	9.1	61	9.0	122	9.1
50–54	432	68.1	349	56.0	781	62.1	128	20.2	78	12.5	206	16.4
55–59	642	131.0	429	90.6	1,071	111.1	197	40.2	142	30.0	339	35.2
60–64	884	220.6	564	142.1	1,448	181.5	306	76.3	175	44.1	481	60.3
65–69	1,002	301.8	653	189.2	1,655	244.4	361	108.7	205	59.4	566	83.6
70–74	1,263	421.6	852	255.4	2,115	334.0	451	150.5	289	86.6	740	116.9
75–79	1,087	495.0	895	311.0	1,982	390.7	429	195.4	364	126.5	793	156.3
80–84	630	529.5	679	357.4	1,309	423.7	316	265.6	310	163.2	626	202.6
85 and over	438	565.0	683	389.9	1,121	443.7	252	325.1	479	273.5	731	289.3
Total	6,863		5,542		12,405		2,569		2,149		4,718	

Rates per 100,000 with 95% confidence intervals (95% CI)

Crude rate	72.2	57.4	64.8	27.0	22.3	24.6
95% CI	70.5 – 73.9	55.9 – 59.0	63.6 – 65.9	26.0 – 28.1	21.3 – 23.2	23.9 – 25.3
AS rate (Aust. 2001)	80.2	53.8	65.7	31.0	20.2	25.1
95% CI	78.3 – 82.2	52.4 – 55.2	64.6 – 66.9	29.8 – 32.2	19.3 – 21.1	24.4 – 25.8
AS rate (new WHO World)	56.3	38.0	46.5	20.8	13.4	16.8
95% CI	55.0 – 57.6	37.0 – 39.1	45.7 – 47.3	20.0 – 21.6	12.8 – 14.0	16.3 – 17.3
Lifetime risk (0–74)	1 in 17	1 in 26	1 in 21	1 in 49	1 in 81	1 in 61
PYLL (0–74)				18,135	12,090	30,225
Per cent of all cancers	14.9	14.1	14.6	12.8	13.9	13.3

Average annual numbers and rates by State and Territory, 1996–2000

	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate
NSW	2,173	76.9	1,780	51.9	3,953	63.0	834	30.9	693	19.8	1,527	24.6
Vic	1,632	79.9	1,391	54.7	3,023	66.0	662	33.6	581	22.5	1,244	27.4
Qld	1,148	78.9	951	56.7	2,099	67.1	446	32.1	356	21.1	802	26.1
WA	544	76.8	420	49.4	964	61.5	228	33.4	181	21.1	408	26.6
SA	563	79.1	506	56.6	1,070	66.7	216	31.1	200	21.9	416	25.9
Tas	162	74.6	155	58.7	317	66.0	71	34.1	74	27.8	145	30.6
ACT	75	75.5	61	52.2	137	63.0	31	33.5	27	24.4	58	29.0
NT	25	54.7	15	42.4	39	49.7	8	21.5	6	18.4	14	20.5

Notes

1. AS rates use Australian 2001 Standard Population unless new WHO World Standard Population is indicated. All rates are expressed per 100,000 population.

2. Incidence numbers are based on state of usual residence at time of diagnosis; mortality numbers on state of registration of death.

Source: *Cancer in Australia 2000*, AIHW & AACR, 2003.

Table 23: Cancer of the colon and rectum (excluding anus) (ICD-10 C18–C20)

Australia 2000

Age group	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
0–4	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
5–9	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
10–14	2	0.3	1	0.2	3	0.2	0	0.0	0	0.0	0	0.0
15–19	2	0.3	4	0.6	6	0.5	0	0.0	0	0.0	0	0.0
20–24	3	0.5	7	1.1	10	0.8	1	0.2	3	0.5	4	0.3
25–29	16	2.2	12	1.7	28	1.9	2	0.3	1	0.1	3	0.2
30–34	32	4.5	28	3.9	60	4.2	10	1.4	9	1.3	19	1.3
35–39	55	7.3	55	7.3	110	7.3	21	2.8	13	1.7	34	2.3
40–44	127	17.6	109	15.0	236	16.3	33	4.6	19	2.6	52	3.6
45–49	231	34.6	206	30.6	437	32.6	61	9.1	58	8.6	119	8.9
50–54	426	67.1	336	53.9	762	60.6	126	19.9	76	12.2	202	16.1
55–59	629	128.3	420	88.7	1,049	108.9	195	39.8	138	29.1	333	34.6
60–64	863	215.3	552	139.1	1,415	177.4	304	75.8	175	44.1	479	60.1
65–69	995	299.7	637	184.6	1,632	241.0	357	107.5	203	58.8	560	82.7
70–74	1,243	414.9	834	250.0	2,077	328.0	446	148.9	286	85.7	732	115.6
75–79	1,076	490.0	883	306.9	1,959	386.1	424	193.1	359	124.8	783	154.3
80–84	625	525.3	668	351.6	1,293	418.5	312	262.3	307	161.6	619	200.3
85 and over	434	559.9	674	384.8	1,108	438.5	251	323.8	469	267.8	720	285.0
Total	6,759		5,426		12,185		2,543		2,116		4,659	

Rates per 100,000 with 95% confidence intervals (95% CI)

Crude rate	71.1	56.2	63.6	26.8	21.9	24.3
95% CI	69.4 – 72.8	54.7 – 57.7	62.5 – 64.7	25.7 – 27.8	21.0 – 22.9	23.6 – 25.0
AS rate (Aust. 2001)	79.1	52.6	64.6	30.7	19.9	24.8
95% CI	77.1 – 81.0	51.2 – 54.1	63.4 – 65.7	29.5 – 31.9	19.0 – 20.8	24.0 – 25.5
AS rate (new WHO World)	55.4	37.2	45.6	20.6	13.2	16.6
95% CI	54.1 – 56.8	36.1 – 38.2	44.8 – 46.5	19.8 – 21.4	12.6 – 13.8	16.1 – 17.1
Lifetime risk (0–74)	1 in 17	1 in 26	1 in 21	1 in 49	1 in 82	1 in 62
PYLL (0–74)				17,950	11,838	29,788
Per cent of all cancers	14.7	13.8	14.3	12.7	13.7	13.1

Average annual numbers and rates by State and Territory, 1996–2000

	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate
NSW	2,128	75.3	1,725	50.2	3,853	61.4	830	30.6	686	19.5	1,517	24.3
Vic	1,616	79.1	1,374	54.1	2,989	65.3	661	33.4	578	22.2	1,240	27.2
Qld	1,129	77.7	931	55.4	2,060	65.9	445	31.9	355	20.9	801	25.9
WA	535	75.6	409	48.2	945	60.3	228	33.1	179	20.8	406	26.3
SA	560	78.6	497	55.5	1,056	65.9	216	31.0	199	21.6	416	25.8
Tas	159	73.0	152	57.5	310	64.6	70	33.6	74	27.5	144	30.2
ACT	74	74.0	60	50.6	134	61.5	30	32.9	28	24.4	58	28.7
NT	24	53.4	15	42.4	39	49.0	8	21.3	6	18.4	14	20.4

Notes

1. AS rates use Australian 2001 Standard Population unless new WHO World Standard Population is indicated. All rates are expressed per 100,000 population.

2. Incidence numbers are based on state of usual residence at time of diagnosis; mortality numbers on state of registration of death.

Source: *Cancer in Australia 2000*, AIHW & AACR, 2003.

Table 24: Cancer of the liver and intrahepatic bile ducts (ICD-10 C22)

Australia 2000

Age group	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
0-4	4	0.6	4	0.6	8	0.6	0	0.0	0	0.0	0	0.0
5-9	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
10-14	0	0.0	2	0.3	2	0.1	2	0.3	0	0.0	2	0.1
15-19	0	0.0	0	0.0	0	0.0	1	0.1	0	0.0	1	0.1
20-24	0	0.0	1	0.2	1	0.1	2	0.3	1	0.2	3	0.2
25-29	0	0.0	2	0.3	2	0.1	1	0.1	1	0.1	2	0.1
30-34	1	0.1	0	0.0	1	0.1	6	0.8	1	0.1	7	0.5
35-39	5	0.7	0	0.0	5	0.3	2	0.3	2	0.3	4	0.3
40-44	16	2.2	2	0.3	18	1.2	9	1.3	4	0.5	13	0.9
45-49	28	4.2	8	1.2	36	2.7	20	3.0	6	0.9	26	1.9
50-54	34	5.4	12	1.9	46	3.7	32	5.0	14	2.2	46	3.7
55-59	55	11.2	16	3.4	71	7.4	36	7.3	13	2.7	49	5.1
60-64	61	15.2	16	4.0	77	9.7	52	13.0	19	4.8	71	8.9
65-69	88	26.5	22	6.4	110	16.2	84	25.3	26	7.5	110	16.2
70-74	101	33.7	36	10.8	137	21.6	100	33.4	26	7.8	126	19.9
75-79	69	31.4	32	11.1	101	19.9	79	36.0	38	13.2	117	23.1
80-84	40	33.6	29	15.3	69	22.3	55	46.2	38	20.0	93	30.1
85 and over	24	31.0	19	10.8	43	17.0	32	41.3	39	22.3	71	28.1
Total	526		201		727		513		228		741	

Rates per 100,000 with 95% confidence intervals (95% CI)

Crude rate	5.5	2.1	3.8	5.4	2.4	3.9
95% CI	5.1 – 6.0	1.8 – 2.4	3.5 – 4.1	4.9 – 5.9	2.1 – 2.7	3.6 – 4.1
AS rate (Aust. 2001)	6.0	2.0	3.8	6.0	2.2	3.9
95% CI	5.5 – 6.5	1.7 – 2.2	3.6 – 4.1	5.5 – 6.5	1.9 – 2.5	3.6 – 4.2
AS rate (new WHO World)	4.4	1.4	2.8	4.2	1.5	2.8
95% CI	4.0 – 4.8	1.2 – 1.6	2.6 – 3.0	3.8 – 4.6	1.3 – 1.7	2.5 – 3.0
Lifetime risk (0-74)	1 in 201	1 in 682	1 in 314	1 in 222	1 in 735	1 in 345
PYLL (0-74)				4,388	1,553	5,940
Per cent of all cancers	1.1	0.5	0.9	2.6	1.5	2.1

Average annual numbers and rates by State and Territory, 1996-2000

	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate
NSW	164	5.6	60	1.7	224	3.5	159	5.5	76	2.2	235	3.7
Vic	142	6.8	46	1.8	188	4.1	138	6.6	59	2.3	197	4.2
Qld	69	4.7	30	1.8	99	3.1	69	4.8	33	1.9	102	3.2
WA	41	5.4	16	1.9	57	3.5	35	4.8	18	2.0	52	3.3
SA	31	4.2	13	1.5	44	2.7	32	4.3	20	2.1	51	3.1
Tas	10	4.4	3	1.3	13	2.7	11	5.2	4	1.6	16	3.3
ACT	6	6.2	3	2.3	9	4.1	5	4.7	4	3.3	8	4.0
NT	5	12.5	1	1.8	6	7.2	4	12.6	1	4.0	5	8.2

Notes

1. AS rates use Australian 2001 Standard Population unless new WHO World Standard Population is indicated. All rates are expressed per 100,000 population.
 2. Incidence numbers are based on state of usual residence at time of diagnosis; mortality numbers on state of registration of death.
- Source: *Cancer in Australia 2000*, AIHW & AACR, 2003.

Table 25: Cancer of the gallbladder and extrahepatic bile ducts (ICD-10 C23–C24)

Australia 2000

Age group	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
0–4	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
5–9	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
10–14	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
15–19	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
20–24	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
25–29	1	0.1	0	0.0	1	0.1	0	0.0	0	0.0	0	0.0
30–34	4	0.6	2	0.3	6	0.4	0	0.0	1	0.1	1	0.1
35–39	3	0.4	2	0.3	5	0.3	1	0.1	1	0.1	2	0.1
40–44	5	0.7	6	0.8	11	0.8	1	0.1	3	0.4	4	0.3
45–49	9	1.3	5	0.7	14	1.0	3	0.4	2	0.3	5	0.4
50–54	18	2.8	18	2.9	36	2.9	0	0.0	10	1.6	10	0.8
55–59	20	4.1	25	5.3	45	4.7	8	1.6	13	2.7	21	2.2
60–64	26	6.5	21	5.3	47	5.9	10	2.5	8	2.0	18	2.3
65–69	26	7.8	32	9.3	58	8.6	16	4.8	24	7.0	40	5.9
70–74	50	16.7	64	19.2	114	18.0	25	8.3	37	11.1	62	9.8
75–79	32	14.6	61	21.2	93	18.3	25	11.4	35	12.2	60	11.8
80–84	26	21.9	52	27.4	78	25.2	19	16.0	36	18.9	55	17.8
85 and over	23	29.7	49	28.0	72	28.5	18	23.2	36	20.6	54	21.4
Total	243		337		580		126		206		332	

Rates per 100,000 with 95% confidence intervals (95% CI)

Crude rate	2.6	3.5	3.0	1.3	2.1	1.7
95% CI	2.2 – 2.9	3.1 – 3.9	2.8 – 3.3	1.1 – 1.6	1.8 – 2.4	1.5 – 1.9
AS rate (Aust. 2001)	2.9	3.2	3.1	1.6	1.9	1.8
95% CI	2.5 – 3.3	2.9 – 3.6	2.8 – 3.3	1.3 – 1.9	1.7 – 2.2	1.6 – 2.0
AS rate (new WHO World)	2.0	2.2	2.1	1.0	1.3	1.1
95% CI	1.7 – 2.2	1.9 – 2.4	1.9 – 2.3	0.8 – 1.2	1.1 – 1.5	1.0 – 1.3
Lifetime risk (0–74)	1 in 487	1 in 455	1 in 470	1 in 1,111	1 in 788	1 in 919
PYLL (0–74)				600	1,058	1,658
Per cent of all cancers	0.5	0.9	0.7	0.6	1.3	0.9

Average annual numbers and rates by State and Territory, 1996–2000

	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate
NSW	87	3.2	126	3.6	213	3.4	34	1.3	66	1.9	100	1.6
Vic	66	3.3	89	3.5	155	3.4	29	1.6	52	2.0	81	1.8
Qld	43	3.0	56	3.3	100	3.2	22	1.7	36	2.1	58	1.9
WA	24	3.5	26	3.0	50	3.2	16	2.4	18	2.1	33	2.2
SA	27	3.8	39	4.2	66	4.0	15	2.3	25	2.7	40	2.5
Tas	7	3.7	10	3.8	18	3.7	5	2.8	7	2.4	12	2.5
ACT	2	2.8	3	2.1	5	2.3	1	1.2	1	1.2	3	1.2
NT	2	6.3	1	1.7	3	3.8	0	1.8	2	7.6	2	4.7

Notes

- AS rates use Australian 2001 Standard Population unless new WHO World Standard Population is indicated. All rates are expressed per 100,000 population.
 - Incidence numbers are based on state of usual residence at time of diagnosis; mortality numbers on state of registration of death.
- Source: *Cancer in Australia 2000*, AIHW & AACR, 2003.

Table 26: Cancer of the pancreas (ICD-10 C25)

Australia 2000

Age group	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
0-4	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
5-9	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
10-14	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
15-19	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
20-24	1	0.2	0	0.0	1	0.1	0	0.0	0	0.0	0	0.0
25-29	2	0.3	1	0.1	3	0.2	0	0.0	0	0.0	0	0.0
30-34	3	0.4	3	0.4	6	0.4	2	0.3	3	0.4	5	0.4
35-39	6	0.8	1	0.1	7	0.5	8	1.1	1	0.1	9	0.6
40-44	15	2.1	16	2.2	31	2.1	10	1.4	6	0.8	16	1.1
45-49	36	5.4	18	2.7	54	4.0	22	3.3	12	1.8	34	2.5
50-54	64	10.1	29	4.7	93	7.4	57	9.0	28	4.5	85	6.8
55-59	67	13.7	48	10.1	115	11.9	74	15.1	41	8.7	115	11.9
60-64	86	21.5	56	14.1	142	17.8	77	19.2	54	13.6	131	16.4
65-69	143	43.1	90	26.1	233	34.4	143	43.1	84	24.3	227	33.5
70-74	150	50.1	141	42.3	291	46.0	142	47.4	151	45.3	293	46.3
75-79	164	74.7	179	62.2	343	67.6	155	70.6	190	66.0	345	68.0
80-84	97	81.5	154	81.1	251	81.2	109	91.6	147	77.4	256	82.9
85 and over	78	100.6	160	91.3	238	94.2	73	94.2	159	90.8	232	91.8
Total	912		896		1,808		872		876		1,748	

Rates per 100,000 with 95% confidence intervals (95% CI)

Crude rate	9.6	9.3	9.4	9.2	9.1	9.1
95% CI	9.0 – 10.2	8.7 – 9.9	9.0 – 9.9	8.6 – 9.8	8.5 – 9.7	8.7 – 9.6
AS rate (Aust. 2001)	10.9	8.4	9.6	10.4	8.2	9.3
95% CI	10.2 – 11.6	7.9 – 9.0	9.2 – 10.0	9.7 – 11.1	7.7 – 8.8	8.8 – 9.7
AS rate (new WHO World)	7.4	5.5	6.4	7.1	5.3	6.2
95% CI	6.9 – 7.9	5.1 – 5.9	6.1 – 6.7	6.6 – 7.5	4.9 – 5.7	5.9 – 6.5
Lifetime risk (0-74)	1 in 136	1 in 195	1 in 161	1 in 144	1 in 201	1 in 168
PYLL (0-74)				6,283	3,720	10,003
Per cent of all cancers	2.0	2.3	2.1	4.4	5.7	4.9

Average annual numbers and rates by State and Territory, 1996-2000

	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate
NSW	293	10.7	305	8.6	598	9.5	277	10.3	292	8.2	569	9.1
Vic	234	11.7	228	8.7	461	10.1	223	11.1	221	8.5	444	9.7
Qld	161	11.3	144	8.4	305	9.8	145	10.4	137	8.0	282	9.2
WA	75	10.6	78	9.2	153	9.9	70	10.1	71	8.3	140	9.2
SA	80	11.4	76	8.1	156	9.6	68	9.8	72	7.8	140	8.6
Tas	21	10.2	30	10.8	51	10.6	21	10.2	26	9.6	47	9.9
ACT	10	10.8	10	9.1	20	9.8	10	10.7	10	9.3	20	10.0
NT	6	9.8	3	12.9	9	11.9	5	9.8	4	14.7	9	13.0

Notes

1. AS rates use Australian 2001 Standard Population unless new WHO World Standard Population is indicated. All rates are expressed per 100,000 population.

2. Incidence numbers are based on state of usual residence at time of diagnosis; mortality numbers on state of registration of death.

Source: *Cancer in Australia 2000*, AIHW & AACR, 2003.

Table 27: Cancer of the nasal cavities, middle ear and accessory sinuses (ICD-10 C30–C31)

Australia 2000

Age group	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
0–4	0	0.0	1	0.2	1	0.1	0	0.0	0	0.0	0	0.0
5–9	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
10–14	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
15–19	0	0.0	1	0.2	1	0.1	0	0.0	0	0.0	0	0.0
20–24	1	0.2	0	0.0	1	0.1	0	0.0	0	0.0	0	0.0
25–29	0	0.0	1	0.1	1	0.1	1	0.1	0	0.0	1	0.1
30–34	1	0.1	1	0.1	2	0.1	0	0.0	0	0.0	0	0.0
35–39	2	0.3	2	0.3	4	0.3	0	0.0	0	0.0	0	0.0
40–44	6	0.8	3	0.4	9	0.6	2	0.3	1	0.1	3	0.2
45–49	6	0.9	3	0.4	9	0.7	0	0.0	3	0.4	3	0.2
50–54	5	0.8	3	0.5	8	0.6	1	0.2	0	0.0	1	0.1
55–59	6	1.2	5	1.1	11	1.1	0	0.0	1	0.2	1	0.1
60–64	10	2.5	5	1.3	15	1.9	5	1.2	0	0.0	5	0.6
65–69	13	3.9	1	0.3	14	2.1	4	1.2	1	0.3	5	0.7
70–74	14	4.7	3	0.9	17	2.7	6	2.0	1	0.3	7	1.1
75–79	11	5.0	13	4.5	24	4.7	2	0.9	0	0.0	2	0.4
80–84	5	4.2	4	2.1	9	2.9	2	1.7	0	0.0	2	0.6
85 and over	8	10.3	6	3.4	14	5.5	2	2.6	4	2.3	6	2.4
Total	88		52		140		25		11		36	

Rates per 100,000 with 95% confidence intervals (95% CI)

Crude rate	0.9	0.5	0.7	0.3	0.1	0.2
95% CI	0.7 – 1.1	0.4 – 0.7	0.6 – 0.9	0.2 – 0.4	0.0 – 0.2	0.1 – 0.2
AS rate (Aust. 2001)	1.0	0.5	0.7	0.3	0.1	0.2
95% CI	0.8 – 1.2	0.4 – 0.6	0.6 – 0.9	0.2 – 0.4	0.0 – 0.2	0.1 – 0.3
AS rate (new WHO World)	0.7	0.4	0.5	0.2	0.1	0.1
95% CI	0.6 – 0.9	0.3 – 0.5	0.5 – 0.6	0.1 – 0.3	0.0 – 0.1	0.1 – 0.2
Lifetime risk (0–74)	1 in 1,300	1 in 3,510	1 in 1,922	1 in 3,978	1 in 14,463	1 in 6,342
PYLL (0–74)				243	143	385
Per cent of all cancers	0.2	0.1	0.2	0.1	0.1	0.1

Average annual numbers and rates by State and Territory, 1996–2000

	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate
NSW	28	1.0	17	0.5	45	0.7	8	0.3	4	0.1	12	0.2
Vic	25	1.2	11	0.4	36	0.8	10	0.5	3	0.1	13	0.3
Qld	12	0.8	6	0.4	18	0.6	4	0.3	1	0.1	5	0.2
WA	8	1.1	4	0.5	12	0.8	2	0.2	1	0.1	3	0.2
SA	6	0.9	4	0.4	10	0.6	2	0.3	2	0.2	4	0.2
Tas	2	1.0	1	0.6	4	0.8	2	1.1	0	0.2	3	0.6
ACT	2	1.2	0	0.3	2	0.8	0	0.3	0	0.3	1	0.3
NT	0	0.0	0	0.5	0	0.2	0	0.0	0	0.0	0	0.0

Notes

1. AS rates use Australian 2001 Standard Population unless new WHO World Standard Population is indicated. All rates are expressed per 100,000 population.

2. Incidence numbers are based on state of usual residence at time of diagnosis; mortality numbers on state of registration of death.

Source: *Cancer in Australia 2000*, AIHW & AACR, 2003.

Table 28: Cancer of the larynx (ICD-10 C32)

Australia 2000

Age group	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
0-4	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
5-9	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
10-14	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
15-19	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
20-24	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
25-29	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
30-34	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
35-39	1	0.1	2	0.3	3	0.2	0	0.0	0	0.0	0	0.0
40-44	12	1.7	0	0.0	12	0.8	3	0.4	2	0.3	5	0.3
45-49	16	2.4	2	0.3	18	1.3	8	1.2	0	0.0	8	0.6
50-54	42	6.6	8	1.3	50	4.0	16	2.5	2	0.3	18	1.4
55-59	60	12.2	8	1.7	68	7.1	18	3.7	2	0.4	20	2.1
60-64	67	16.7	11	2.8	78	9.8	29	7.2	5	1.3	34	4.3
65-69	78	23.5	9	2.6	87	12.8	30	9.0	7	2.0	37	5.5
70-74	76	25.4	13	3.9	89	14.1	40	13.4	6	1.8	46	7.3
75-79	83	37.8	9	3.1	92	18.1	30	13.7	3	1.0	33	6.5
80-84	29	24.4	2	1.1	31	10.0	19	16.0	5	2.6	24	7.8
85 and over	17	21.9	5	2.9	22	8.7	10	12.9	4	2.3	14	5.5
Total	481		69		550		203		36		239	

Rates per 100,000 with 95% confidence intervals (95% CI)

Crude rate	5.1	0.7	2.9	2.1	0.4	1.2
95% CI	4.6 – 5.5	0.5 – 0.9	2.6 – 3.1	1.8 – 2.4	0.3 – 0.5	1.1 – 1.4
AS rate (Aust. 2001)	5.5	0.7	2.9	2.4	0.4	1.3
95% CI	5.0 – 6.0	0.5 – 0.9	2.7 – 3.2	2.0 – 2.7	0.2 – 0.5	1.1 – 1.4
AS rate (new WHO World)	4.0	0.5	2.2	1.7	0.3	0.9
95% CI	3.6 – 4.3	0.4 – 0.7	2.0 – 2.3	1.4 – 1.9	0.2 – 0.3	0.8 – 1.0
Lifetime risk (0-74)	1 in 226	1 in 1,562	1 in 400	1 in 535	1 in 3,277	1 in 933
PYLL (0-74)				1,680	275	1,955
Per cent of all cancers	1.0	0.2	0.6	1.0	0.2	0.7

Average annual numbers and rates by State and Territory, 1996-2000

	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate
NSW	188	6.4	23	0.7	211	3.4	78	2.8	11	0.3	89	1.4
Vic	125	5.9	20	0.8	145	3.2	49	2.5	8	0.3	57	1.3
Qld	93	6.2	10	0.6	103	3.2	34	2.4	4	0.3	38	1.2
WA	46	6.1	6	0.7	52	3.2	18	2.6	4	0.4	22	1.4
SA	37	4.9	4	0.5	41	2.5	18	2.6	1	0.2	19	1.2
Tas	13	5.9	2	0.9	15	3.2	4	1.7	1	0.3	4	0.9
ACT	6	6.1	1	0.7	7	3.1	2	1.5	0	0.3	2	0.9
NT	5	10.5	1	2.3	6	6.8	3	8.1	0	0.9	4	4.7

Notes

1. AS rates use Australian 2001 Standard Population unless new WHO World Standard Population is indicated. All rates are expressed per 100,000 population.
 2. Incidence numbers are based on state of usual residence at time of diagnosis; mortality numbers on state of registration of death.
- Source: *Cancer in Australia 2000*, AIHW & AACR, 2003.

Table 29: Cancer of the trachea, bronchus and lung (ICD-10 C33–C34)

Australia 2000

Age group	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
0–4	0	0.0	1	0.2	1	0.1	0	0.0	0	0.0	0	0.0
5–9	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
10–14	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
15–19	2	0.3	2	0.3	4	0.3	0	0.0	0	0.0	0	0.0
20–24	2	0.3	0	0.0	2	0.2	0	0.0	0	0.0	0	0.0
25–29	2	0.3	2	0.3	4	0.3	2	0.3	0	0.0	2	0.1
30–34	11	1.6	4	0.6	15	1.1	3	0.4	1	0.1	4	0.3
35–39	27	3.6	20	2.6	47	3.1	12	1.6	12	1.6	24	1.6
40–44	54	7.5	47	6.4	101	7.0	37	5.1	26	3.6	63	4.3
45–49	113	16.9	87	12.9	200	14.9	72	10.8	69	10.2	141	10.5
50–54	261	41.1	172	27.6	433	34.4	183	28.8	130	20.9	313	24.9
55–59	444	90.6	262	55.3	706	73.3	350	71.4	181	38.2	531	55.1
60–64	599	149.5	301	75.8	900	112.8	509	127.0	222	55.9	731	91.6
65–69	839	252.7	409	118.5	1,248	184.3	703	211.7	303	87.8	1,006	148.6
70–74	1,086	362.5	508	152.3	1,594	251.7	938	313.1	437	131.0	1,375	217.1
75–79	989	450.4	450	156.4	1,439	283.6	914	416.2	404	140.4	1,318	259.8
80–84	524	440.5	295	155.3	819	265.1	519	436.2	293	154.2	812	262.8
85 and over	325	419.3	222	126.7	547	216.5	352	454.1	239	136.5	591	233.9
Total	5,278		2,782		8,060		4,594		2,317		6,911	

Rates per 100,000 with 95% confidence intervals (95% CI)

Crude rate	55.5	28.8	42.1	48.3	24.0	36.1
95% CI	54.0 – 57.0	27.8 – 29.9	41.2 – 43.0	46.9 – 49.7	23.0 – 25.0	35.2 – 36.9
AS rate (Aust. 2001)	62.1	27.4	42.7	54.8	22.5	36.6
95% CI	60.4 – 63.8	26.3 – 28.4	41.7 – 43.6	53.2 – 56.4	21.6 – 23.4	35.7 – 37.5
AS rate (new WHO World)	42.7	19.7	30.1	36.8	15.8	25.2
95% CI	41.5 – 43.8	19.0 – 20.5	29.5 – 30.8	35.8 – 37.9	15.1 – 16.4	24.6 – 25.8
Lifetime risk (0–74)	1 in 22	1 in 45	1 in 30	1 in 26	1 in 58	1 in 37
PYLL (0–74)				28,078	15,468	43,545
Per cent of all cancers	11.5	7.1	9.5	22.9	15.0	19.5

Average annual numbers and rates by State and Territory, 1996–2000

	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate
NSW	1,825	64.6	876	25.7	2,701	42.7	1,558	56.3	719	21.0	2,277	36.2
Vic	1,332	65.5	687	27.4	2,018	43.9	1,151	57.6	558	22.2	1,709	37.4
Qld	1,014	70.3	432	26.0	1,447	46.1	881	62.2	352	21.2	1,233	39.6
WA	480	68.6	243	28.9	723	46.3	409	60.0	200	23.8	609	39.5
SA	479	67.0	216	24.4	694	42.8	414	58.7	174	19.5	588	36.3
Tas	155	71.3	76	29.7	232	48.1	141	65.4	61	23.6	203	42.2
ACT	43	45.6	32	27.1	75	35.3	46	49.2	31	26.4	76	36.4
NT	34	76.9	14	42.3	48	61.9	29	71.0	12	36.5	41	55.5

Notes

1. AS rates use Australian 2001 Standard Population unless new WHO World Standard Population is indicated. All rates are expressed per 100,000 population.

2. Incidence numbers are based on state of usual residence at time of diagnosis; mortality numbers on state of registration of death.

Source: *Cancer in Australia 2000*, AIHW & AACR, 2003.

Table 30: Cancer of the thymus, heart, mediastinum and pleura (ICD-10 C37–C38)

Australia 2000

Age group	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
0–4	0	0.0	1	0.2	1	0.1	0	0.0	0	0.0	0	0.0
5–9	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
10–14	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
15–19	2	0.3	0	0.0	2	0.2	0	0.0	0	0.0	0	0.0
20–24	6	0.9	1	0.2	7	0.5	2	0.3	0	0.0	2	0.2
25–29	2	0.3	0	0.0	2	0.1	2	0.3	0	0.0	2	0.1
30–34	3	0.4	1	0.1	4	0.3	3	0.4	0	0.0	3	0.2
35–39	2	0.3	1	0.1	3	0.2	1	0.1	1	0.1	2	0.1
40–44	7	1.0	1	0.1	8	0.6	3	0.4	0	0.0	3	0.2
45–49	3	0.4	0	0.0	3	0.2	0	0.0	1	0.1	1	0.1
50–54	7	1.1	2	0.3	9	0.7	4	0.6	1	0.2	5	0.4
55–59	4	0.8	3	0.6	7	0.7	1	0.2	4	0.8	5	0.5
60–64	4	1.0	2	0.5	6	0.8	1	0.2	1	0.3	2	0.3
65–69	6	1.8	1	0.3	7	1.0	3	0.9	2	0.6	5	0.7
70–74	4	1.3	3	0.9	7	1.1	5	1.7	4	1.2	9	1.4
75–79	6	2.7	1	0.3	7	1.4	4	1.8	2	0.7	6	1.2
80–84	4	3.4	3	1.6	7	2.3	2	1.7	3	1.6	5	1.6
85 and over	0	0.0	1	0.6	1	0.4	3	3.9	4	2.3	7	2.8
Total	60		21		81		34		23		57	

Rates per 100,000 with 95% confidence intervals (95% CI)

Crude rate	0.6	0.2	0.4	0.4	0.2	0.3
95% CI	0.5 – 0.8	0.1 – 0.3	0.3 – 0.5	0.2 – 0.5	0.1 – 0.3	0.2 – 0.4
AS rate (Aust. 2001)	0.7	0.2	0.4	0.4	0.2	0.3
95% CI	0.5 – 0.8	0.1 – 0.3	0.3 – 0.5	0.3 – 0.5	0.1 – 0.3	0.2 – 0.4
AS rate (new WHO World)	0.6	0.2	0.4	0.3	0.2	0.2
95% CI	0.4 – 0.7	0.1 – 0.3	0.3 – 0.4	0.2 – 0.4	0.1 – 0.2	0.2 – 0.3
Lifetime risk (0–74)	1 in 2,071	1 in 5,928	1 in 3,078	1 in 3,837	1 in 6,031	1 in 4,712
PYLL (0–74)				618	195	813
Per cent of all cancers	0.1	0.1	0.1	0.2	0.1	0.2

Average annual numbers and rates by State and Territory, 1996–2000

	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate
NSW	18	0.6	11	0.3	29	0.5	23	0.8	12	0.4	35	0.6
Vic	14	0.7	11	0.4	24	0.5	12	0.6	4	0.2	16	0.3
Qld	13	0.8	6	0.4	19	0.6	10	0.7	3	0.2	13	0.4
WA	4	0.5	1	0.1	5	0.3	14	2.0	3	0.4	17	1.1
SA	4	0.5	2	0.3	6	0.4	5	0.7	4	0.4	9	0.6
Tas	1	0.3	1	0.3	1	0.3	2	0.7	1	0.4	3	0.5
ACT	1	0.4	0	0.1	1	0.3	1	0.8	0	0.4	1	0.6
NT	0	0.4	0	0.4	1	0.4	0	0.4	0	0.0	0	0.2

Notes

1. AS rates use Australian 2001 Standard Population unless new WHO World Standard Population is indicated. All rates are expressed per 100,000 population.

2. Incidence numbers are based on state of usual residence at time of diagnosis; mortality numbers on state of registration of death.

Source: *Cancer in Australia 2000*, AIHW & AACR, 2003.

Table 31: Cancer of the bone and articular cartilage (ICD-10 C40–C41)

Australia 2000

Age group	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
0–4	1	0.2	1	0.2	2	0.2	0	0.0	1	0.2	1	0.1
5–9	2	0.3	2	0.3	4	0.3	0	0.0	0	0.0	0	0.0
10–14	8	1.2	9	1.4	17	1.3	0	0.0	4	0.6	4	0.3
15–19	12	1.8	8	1.2	20	1.5	3	0.4	3	0.5	6	0.5
20–24	7	1.1	8	1.3	15	1.2	5	0.8	3	0.5	8	0.6
25–29	5	0.7	5	0.7	10	0.7	2	0.3	0	0.0	2	0.1
30–34	4	0.6	4	0.6	8	0.6	0	0.0	2	0.3	2	0.1
35–39	7	0.9	4	0.5	11	0.7	0	0.0	1	0.1	1	0.1
40–44	3	0.4	4	0.5	7	0.5	1	0.1	1	0.1	2	0.1
45–49	7	1.0	7	1.0	14	1.0	1	0.1	1	0.1	2	0.1
50–54	7	1.1	0	0.0	7	0.6	1	0.2	2	0.3	3	0.2
55–59	5	1.0	1	0.2	6	0.6	2	0.4	1	0.2	3	0.3
60–64	8	2.0	2	0.5	10	1.3	1	0.2	1	0.3	2	0.3
65–69	5	1.5	4	1.2	9	1.3	3	0.9	0	0.0	3	0.4
70–74	6	2.0	5	1.5	11	1.7	3	1.0	1	0.3	4	0.6
75–79	3	1.4	3	1.0	6	1.2	2	0.9	3	1.0	5	1.0
80–84	0	0.0	2	1.1	2	0.6	1	0.8	2	1.1	3	1.0
85 and over	3	3.9	0	0.0	3	1.2	4	5.2	6	3.4	10	4.0
Total	93		69		162		29		32		61	

Rates per 100,000 with 95% confidence intervals (95% CI)

Crude rate	1.0	0.7	0.8	0.3	0.3	0.3
95% CI	0.8 – 1.2	0.5 – 0.9	0.7 – 1.0	0.2 – 0.4	0.2 – 0.4	0.2 – 0.4
AS rate (Aust. 2001)	1.0	0.7	0.8	0.3	0.3	0.3
95% CI	0.8 – 1.2	0.5 – 0.9	0.7 – 1.0	0.2 – 0.5	0.2 – 0.4	0.2 – 0.4
AS rate (new WHO World)	0.9	0.7	0.8	0.3	0.3	0.3
95% CI	0.8 – 1.1	0.5 – 0.9	0.7 – 1.0	0.2 – 0.4	0.2 – 0.4	0.2 – 0.4
Lifetime risk (0–74)	1 in 1,272	1 in 1,808	1 in 1,493	1 in 4,453	1 in 5,736	1 in 5,056
PYLL (0–74)				690	913	1,603
Per cent of all cancers	0.2	0.2	0.2	0.1	0.2	0.2

Average annual numbers and rates by State and Territory, 1996–2000

	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate
NSW	31	1.0	21	0.6	52	0.8	19	0.7	13	0.4	32	0.5
Vic	27	1.2	23	0.9	50	1.1	14	0.6	9	0.4	23	0.5
Qld	18	1.1	15	0.8	33	1.0	11	0.7	7	0.4	18	0.6
WA	8	1.0	9	1.0	17	1.0	3	0.4	3	0.4	7	0.4
SA	8	1.1	6	0.7	14	0.9	4	0.5	3	0.3	6	0.4
Tas	2	1.0	2	0.7	4	0.8	0	0.1	1	0.5	1	0.3
ACT	2	1.1	2	1.3	4	1.2	1	0.5	0	0.2	1	0.4
NT	1	0.6	1	0.9	1	0.7	0	0.2	0	0.4	0	0.3

Notes

1. AS rates use Australian 2001 Standard Population unless new WHO World Standard Population is indicated. All rates are expressed per 100,000 population.

2. Incidence numbers are based on state of usual residence at time of diagnosis; mortality numbers on state of registration of death.

Source: *Cancer in Australia 2000*, AIHW & AACR, 2003.

Table 32: Cancer of the skin—melanoma (ICD-10 C43)

Australia 2000

Age group	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
0-4	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
5-9	0	0.0	1	0.2	1	0.1	0	0.0	0	0.0	0	0.0
10-14	5	0.7	5	0.8	10	0.7	0	0.0	0	0.0	0	0.0
15-19	34	5.0	36	5.5	70	5.3	2	0.3	0	0.0	2	0.2
20-24	70	10.7	96	15.1	166	12.9	2	0.3	4	0.6	6	0.5
25-29	127	17.6	147	20.2	274	18.9	3	0.4	4	0.6	7	0.5
30-34	163	23.0	217	30.2	380	26.6	8	1.1	9	1.3	17	1.2
35-39	226	30.2	296	39.1	522	34.7	16	2.1	21	2.8	37	2.5
40-44	314	43.6	317	43.5	631	43.6	24	3.3	18	2.5	42	2.9
45-49	383	57.4	347	51.5	730	54.4	28	4.2	24	3.6	52	3.9
50-54	467	73.6	357	57.3	824	65.5	43	6.8	21	3.4	64	5.1
55-59	479	97.7	309	65.3	788	81.8	47	9.6	26	5.5	73	7.6
60-64	471	117.5	277	69.8	748	93.8	62	15.5	29	7.3	91	11.4
65-69	469	141.3	303	87.8	772	114.0	61	18.4	25	7.2	86	12.7
70-74	609	203.3	315	94.4	924	145.9	110	36.7	44	13.2	154	24.3
75-79	479	218.1	335	116.4	814	160.4	95	43.3	43	14.9	138	27.2
80-84	286	240.4	199	104.7	485	157.0	73	61.4	43	22.6	116	37.5
85 and over	188	242.5	204	116.5	392	155.1	43	55.5	43	24.6	86	34.0
Total	4,770		3,761		8,531		617		354		971	

Rates per 100,000 with 95% confidence intervals (95% CI)

Crude rate	50.2	39.0	44.5	6.5	3.7	5.1
95% CI	48.8 – 51.6	37.7 – 40.2	43.6 – 45.5	6.0 – 7.0	3.3 – 4.1	4.8 – 5.4
AS rate (Aust. 2001)	53.7	38.0	45.0	7.3	3.5	5.1
95% CI	52.1 – 55.2	36.8 – 39.2	44.0 – 45.9	6.7 – 7.9	3.1 – 3.8	4.8 – 5.5
AS rate (new WHO World)	41.4	31.2	35.9	5.1	2.6	3.7
95% CI	40.2 – 42.6	30.2 – 32.3	35.1 – 36.7	4.7 – 5.5	2.3 – 2.8	3.5 – 3.9
Lifetime risk (0-74)	1 in 25	1 in 35	1 in 29	1 in 203	1 in 419	1 in 276
PYLL (0-74)				6,150	4,403	10,553
Per cent of all cancers	10.4	9.6	10.0	3.1	2.3	2.7

Average annual numbers and rates by State and Territory, 1996-2000

	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate
NSW	1,585	54.3	1,140	34.8	2,724	43.4	231	8.4	119	3.5	349	5.6
Vic	862	40.7	815	33.8	1,677	36.6	127	6.3	79	3.1	206	4.5
Qld	1,149	74.0	866	51.5	2,015	61.8	134	9.3	72	4.2	206	6.5
WA	445	56.3	331	37.9	776	46.2	54	7.5	31	3.6	85	5.4
SA	322	44.7	284	35.2	606	39.2	41	5.7	27	3.2	68	4.3
Tas	92	41.8	99	40.6	191	40.5	10	4.9	8	3.1	18	3.8
ACT	61	49.0	44	32.0	106	40.2	6	5.9	4	3.2	10	4.3
NT	26	37.5	19	27.0	45	32.7	3	6.1	1	3.6	4	4.9

Notes

1. AS rates use Australian 2001 Standard Population unless new WHO World Standard Population is indicated. All rates are expressed per 100,000 population.

2. Incidence numbers are based on state of usual residence at time of diagnosis; mortality numbers on state of registration of death.

Source: *Cancer in Australia 2000*, AIHW & AACR, 2003.

Table 33: Skin cancers other than melanoma (ICD-10 C44)

Australia 2000

Age group	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
0-4							0	0.0	0	0.0	0	0.0
5-9							0	0.0	0	0.0	0	0.0
10-14							0	0.0	0	0.0	0	0.0
15-19							0	0.0	0	0.0	0	0.0
20-24							0	0.0	0	0.0	0	0.0
25-29							0	0.0	0	0.0	0	0.0
30-34			Non-melanocytic skin cancer incidence data are not routinely collected by all State and Territory cancer registries				0	0.0	0	0.0	0	0.0
35-39							1	0.1	0	0.0	1	0.1
40-44							2	0.3	3	0.4	5	0.3
45-49							3	0.4	2	0.3	5	0.4
50-54							8	1.3	1	0.2	9	0.7
55-59							12	2.4	4	0.8	16	1.7
60-64							14	3.5	4	1.0	18	2.3
65-69							25	7.5	6	1.7	31	4.6
70-74							37	12.4	8	2.4	45	7.1
75-79							50	22.8	21	7.3	71	14.0
80-84							46	38.7	22	11.6	68	22.0
85 and over							43	55.5	47	26.8	90	35.6
Total							241		118		359	

Rates per 100,000 with 95% confidence intervals (95% CI)

Crude rate		2.5	1.2	1.9
95% CI		2.2 – 2.9	1.0 – 1.4	1.7 – 2.1
AS rate (Aust. 2001)		3.1	1.0	1.9
95% CI		2.7 – 3.5	0.9 – 1.2	1.7 – 2.1
AS rate (new WHO World)		1.9	0.6	1.2
95% CI		1.6 – 2.1	0.5 – 0.7	1.0 – 1.3
Lifetime risk (0-74)		1 in 716	1 in 2,917	1 in 1,170
PYLL (0-74)		1,030	360	1,390
Per cent of all cancers		1.2	0.8	1.0

Average annual numbers and rates by State and Territory, 1996-2000

	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate
NSW							95	4.0	46	1.3	141	2.4
Vic							46	2.7	23	0.9	68	1.6
Qld							52	4.2	22	1.3	74	2.6
WA							22	3.7	10	1.1	31	2.2
SA							12	1.8	6	0.7	18	1.1
Tas							5	3.0	1	0.5	7	1.4
ACT							2	2.8	1	0.9	3	1.7
NT							2	4.8	0	0.0	2	2.5

Notes

1. AS rates use Australian 2001 Standard Population unless new WHO World Standard Population is indicated. All rates are expressed per 100,000 population.

2. Incidence numbers are based on state of usual residence at time of diagnosis; mortality numbers on state of registration of death.

Source: *Cancer in Australia 2000*, AIHW & AACR, 2003.

Table 34: Mesothelioma (ICD-10 C45)

Australia 2000

Age group	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
0-4	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
5-9	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
10-14	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
15-19	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
20-24	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
25-29	0	0.0	0	0.0	0	0.0	1	0.1	0	0.0	1	0.1
30-34	0	0.0	1	0.1	1	0.1	1	0.1	0	0.0	1	0.1
35-39	1	0.1	3	0.4	4	0.3	0	0.0	4	0.5	4	0.3
40-44	3	0.4	2	0.3	5	0.3	2	0.3	1	0.1	3	0.2
45-49	9	1.3	2	0.3	11	0.8	4	0.6	0	0.0	4	0.3
50-54	23	3.6	7	1.1	30	2.4	21	3.3	3	0.5	24	1.9
55-59	45	9.2	5	1.1	50	5.2	35	7.1	3	0.6	38	3.9
60-64	59	14.7	7	1.8	66	8.3	46	11.5	4	1.0	50	6.3
65-69	63	19.0	5	1.4	68	10.0	59	17.8	3	0.9	62	9.2
70-74	72	24.0	13	3.9	85	13.4	72	24.0	16	4.8	88	13.9
75-79	52	23.7	17	5.9	69	13.6	64	29.1	15	5.2	79	15.6
80-84	34	28.6	5	2.6	39	12.6	41	34.5	6	3.2	47	15.2
85 and over	30	38.7	8	4.6	38	15.0	28	36.1	6	3.4	34	13.5
Total	391		75		466		374		61		435	

Rates per 100,000 with 95% confidence intervals (95% CI)

Crude rate	4.1	0.8	2.4	3.9	0.6	2.3
95% CI	3.7 – 4.5	0.6 – 1.0	2.2 – 2.7	3.5 – 4.3	0.5 – 0.8	2.1 – 2.5
AS rate (Aust. 2001)	4.6	0.7	2.5	4.4	0.6	2.3
95% CI	4.1 – 5.0	0.6 – 0.9	2.2 – 2.7	4.0 – 4.9	0.4 – 0.7	2.1 – 2.5
AS rate (new WHO World)	3.2	0.5	1.8	3.0	0.4	1.6
95% CI	2.9 – 3.5	0.4 – 0.6	1.6 – 1.9	2.7 – 3.3	0.3 – 0.5	1.5 – 1.8
Lifetime risk (0-74)	1 in 277	1 in 1,924	1 in 491	1 in 309	1 in 2,366	1 in 555
PYLL (0-74)				2,548	415	2,963
Per cent of all cancers	0.9	0.2	0.5	1.9	0.4	1.2

Average annual numbers and rates by State and Territory, 1996-2000

	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate
NSW	134	4.7	24	0.7	158	2.5						
Vic	84	4.1	19	0.8	102	2.2						
Qld	69	4.8	10	0.6	79	2.5						
WA	55	7.7	7	0.8	62	3.9						
SA	37	5.1	7	0.8	43	2.7						
Tas	5	2.4	1	0.3	6	1.2						
ACT	4	3.8	1	0.6	4	2.0						
NT	2	4.5	0	0.8	2	2.8						

Notes

- AS rates use Australian 2001 Standard Population unless new WHO World Standard Population is indicated. All rates are expressed per 100,000 population.
 - Incidence numbers are based on state of usual residence at time of diagnosis; mortality numbers on state of registration of death.
- Source: *Cancer in Australia 2000*, AIHW & AACR, 2003.

Table 35: Kaposi's sarcoma (ICD-10 C46)

Australia 2000

Age group	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
0-4	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
5-9	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
10-14	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
15-19	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
20-24	1	0.2	0	0.0	1	0.1	0	0.0	0	0.0	0	0.0
25-29	4	0.6	0	0.0	4	0.3	0	0.0	0	0.0	0	0.0
30-34	3	0.4	0	0.0	3	0.2	0	0.0	0	0.0	0	0.0
35-39	5	0.7	0	0.0	5	0.3	0	0.0	0	0.0	0	0.0
40-44	4	0.6	0	0.0	4	0.3	0	0.0	0	0.0	0	0.0
45-49	11	1.6	1	0.1	12	0.9	0	0.0	0	0.0	0	0.0
50-54	3	0.5	1	0.2	4	0.3	0	0.0	0	0.0	0	0.0
55-59	2	0.4	1	0.2	3	0.3	0	0.0	0	0.0	0	0.0
60-64	5	1.2	0	0.0	5	0.6	0	0.0	0	0.0	0	0.0
65-69	2	0.6	0	0.0	2	0.3	0	0.0	0	0.0	0	0.0
70-74	5	1.7	2	0.6	7	1.1	0	0.0	0	0.0	0	0.0
75-79	10	4.6	1	0.3	11	2.2	0	0.0	0	0.0	0	0.0
80-84	4	3.4	1	0.5	5	1.6	1	0.8	0	0.0	1	0.3
85 and over	2	2.6	0	0.0	2	0.8	2	2.6	1	0.6	3	1.2
Total	61		7		68		3		1		4	

Rates per 100,000 with 95% confidence intervals (95% CI)

Crude rate	0.6	0.1	0.4	0.0	0.0	0.0
95% CI	0.5 - 0.8	0.0 - 0.1	0.3 - 0.4	0.0 - 0.1	0.0 - 0.0	0.0 - 0.0
AS rate (Aust. 2001)	0.7	0.1	0.4	0.0	0.0	0.0
95% CI	0.5 - 0.9	0.0 - 0.1	0.3 - 0.4	0.0 - 0.1	0.0 - 0.0	0.0 - 0.0
AS rate (new WHO World)	0.5	0.1	0.3	0.0	0.0	0.0
95% CI	0.4 - 0.7	0.0 - 0.1	0.2 - 0.3	0.0 - 0.0	0.0 - 0.0	0.0 - 0.0
Lifetime risk (0-74)	1 in 2,381	1 in 17,866	1 in 4,234	#DIV/0!	#DIV/0!	#DIV/0!
PYLL (0-74)				0	0	0
Per cent of all cancers	0.1	0.0	0.1	0.0	0.0	0.0

Average annual numbers and rates by State and Territory, 1996-2000

	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate
NSW	42	1.4	4	0.1	46	0.7						
Vic	20	0.9	4	0.1	24	0.5						
Qld	6	0.4	1	0.0	7	0.2						
WA	4	0.6	1	0.1	5	0.3						
SA	5	0.7	1	0.1	6	0.4						
Tas	0	0.0	0	0.2	0	0.1						
ACT	1	0.7	0	0.3	1	0.5						
NT	0	1.1	0	0.0	0	0.5						

There were no comparable mortality data for the period 1996-2000

Notes

1. AS rates use Australian 2001 Standard Population unless new WHO World Standard Population is indicated. All rates are expressed per 100,000 population.

2. Incidence numbers are based on state of usual residence at time of diagnosis; mortality numbers on state of registration of death.

Source: *Cancer in Australia 2000*, AIHW & AACR, 2003.

Table 36: Cancer of the peripheral nerves and autonomic nervous system (ICD-10 C47)

Australia 2000

Age group	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
0-4	1	0.2	1	0.2	2	0.2	0	0.0	0	0.0	0	0.0
5-9	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
10-14	1	0.1	0	0.0	1	0.1	0	0.0	0	0.0	0	0.0
15-19	0	0.0	2	0.3	2	0.2	0	0.0	0	0.0	0	0.0
20-24	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
25-29	1	0.1	1	0.1	2	0.1	0	0.0	0	0.0	0	0.0
30-34	1	0.1	3	0.4	4	0.3	1	0.1	0	0.0	1	0.1
35-39	0	0.0	1	0.1	1	0.1	0	0.0	1	0.1	1	0.1
40-44	3	0.4	1	0.1	4	0.3	1	0.1	0	0.0	1	0.1
45-49	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
50-54	1	0.2	1	0.2	2	0.2	1	0.2	0	0.0	1	0.1
55-59	1	0.2	1	0.2	2	0.2	0	0.0	1	0.2	1	0.1
60-64	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
65-69	0	0.0	2	0.6	2	0.3	0	0.0	2	0.6	2	0.3
70-74	0	0.0	2	0.6	2	0.3	0	0.0	1	0.3	1	0.2
75-79	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
80-84	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
85 and over	0	0.0	1	0.6	1	0.4	0	0.0	0	0.0	0	0.0
Total	9		16		25		3		5		8	

Rates per 100,000 with 95% confidence intervals (95% CI)

Crude rate	0.1	0.2	0.1	0.0	0.1	0.0
95% CI	0.0 – 0.2	0.1 – 0.2	0.1 – 0.2	0.0 – 0.1	0.0 – 0.1	0.0 – 0.1
AS rate (Aust. 2001)	0.1	0.2	0.1	0.0	0.1	0.0
95% CI	0.0 – 0.2	0.1 – 0.2	0.1 – 0.2	0.0 – 0.1	0.0 – 0.1	0.0 – 0.1
AS rate (new WHO World)	0.1	0.2	0.1	0.0	0.0	0.0
95% CI	0.0 – 0.2	0.1 – 0.2	0.1 – 0.2	0.0 – 0.1	0.0 – 0.1	0.0 – 0.1
Lifetime risk (0-74)	1 in 14,744	1 in 7,033	1 in 9,432	1 in 45,695	1 in 16,358	1 in 23,749
PYLL (0-74)				98	73	170
Per cent of all cancers	0.0	0.0	0.0	0.0	0.0	0.0

Average annual numbers and rates by State and Territory, 1996-2000

	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate
NSW	6	0.2	6	0.2	12	0.2	4	0.2	2	0.1	7	0.1
Vic	2	0.1	3	0.1	5	0.1	3	0.1	2	0.1	5	0.1
Qld	2	0.1	2	0.1	4	0.1	2	0.1	3	0.2	4	0.1
WA	0	0.0	0	0.0	0	0.0	1	0.1	1	0.1	2	0.1
SA	2	0.2	1	0.2	3	0.2	0	0.0	1	0.1	1	0.1
Tas	0	0.0	0	0.0	0	0.0	1	0.5	0	0.2	2	0.3
ACT	0	0.5	0	0.0	0	0.2	0	0.0	0	0.0	0	0.0
NT	0	0.2	0	0.0	0	0.1	0	0.0	0	0.0	0	0.0

Notes

1. AS rates use Australian 2001 Standard Population unless new WHO World Standard Population is indicated. All rates are expressed per 100,000 population.
 2. Incidence numbers are based on state of usual residence at time of diagnosis; mortality numbers on state of registration of death.
- Source: *Cancer in Australia 2000*, AIHW & AACR, 2003.

Table 37: Cancer of the retroperitoneum and peritoneum (ICD-10 C48)

Australia 2000

Age group	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
0-4	3	0.5	1	0.2	4	0.3	0	0.0	0	0.0	0	0.0
5-9	0	0.0	2	0.3	2	0.1	0	0.0	0	0.0	0	0.0
10-14	0	0.0	0	0.0	0	0.0	1	0.1	0	0.0	1	0.1
15-19	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
20-24	0	0.0	1	0.2	1	0.1	0	0.0	0	0.0	0	0.0
25-29	1	0.1	0	0.0	1	0.1	1	0.1	0	0.0	1	0.1
30-34	1	0.1	1	0.1	2	0.1	0	0.0	0	0.0	0	0.0
35-39	1	0.1	1	0.1	2	0.1	1	0.1	1	0.1	2	0.1
40-44	1	0.1	3	0.4	4	0.3	3	0.4	3	0.4	6	0.4
45-49	0	0.0	4	0.6	4	0.3	0	0.0	2	0.3	2	0.1
50-54	1	0.2	8	1.3	9	0.7	1	0.2	2	0.3	3	0.2
55-59	2	0.4	6	1.3	8	0.8	2	0.4	0	0.0	2	0.2
60-64	1	0.2	13	3.3	14	1.8	2	0.5	4	1.0	6	0.8
65-69	2	0.6	13	3.8	15	2.2	6	1.8	6	1.7	12	1.8
70-74	5	1.7	12	3.6	17	2.7	4	1.3	6	1.8	10	1.6
75-79	4	1.8	16	5.6	20	3.9	2	0.9	5	1.7	7	1.4
80-84	0	0.0	11	5.8	11	3.6	1	0.8	9	4.7	10	3.2
85 and over	2	2.6	7	4.0	9	3.6	1	1.3	5	2.9	6	2.4
Total	24		99		123		25		43		68	

Rates per 100,000 with 95% confidence intervals (95% CI)

Crude rate	0.3	1.0	0.6	0.3	0.4	0.4
95% CI	0.2 – 0.4	0.8 – 1.2	0.5 – 0.8	0.2 – 0.4	0.3 – 0.6	0.3 – 0.4
AS rate (Aust. 2001)	0.3	1.0	0.7	0.3	0.4	0.4
95% CI	0.2 – 0.4	0.8 – 1.2	0.5 – 0.8	0.2 – 0.4	0.3 – 0.5	0.3 – 0.4
AS rate (new WHO World)	0.2	0.7	0.5	0.2	0.3	0.3
95% CI	0.1 – 0.3	0.6 – 0.9	0.4 – 0.6	0.1 – 0.3	0.2 – 0.4	0.2 – 0.3
Lifetime risk (0-74)	1 in 4,883	1 in 1,326	1 in 2,072	1 in 3,967	1 in 3,505	1 in 3,711
PYLL (0-74)				383	345	728
Per cent of all cancers	0.1	0.3	0.1	0.1	0.3	0.2

Average annual numbers and rates by State and Territory, 1996-2000

	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate
NSW	12	0.4	38	1.2	51	0.8	7	0.2	17	0.5	25	0.4
Vic	7	0.3	19	0.8	26	0.6	4	0.2	9	0.4	13	0.3
Qld	6	0.4	17	1.0	23	0.7	0	0.3	10	0.6	14	0.5
WA	4	0.5	7	0.8	10	0.6	3	0.4	4	0.4	7	0.4
SA	0	0.0	0	0.0	0	0.0	2	0.3	4	0.5	6	0.4
Tas	0	0.0	0	0.0	0	0.0	0	0.0	0	0.1	0	0.0
ACT	0	0.3	1	0.7	1	0.5	0	0.0	0	0.2	0	0.1
NT	0	0.5	0	0.0	0	0.3	0	1.5	0	0.0	0	0.6

Notes

1. AS rates use Australian 2001 Standard Population unless new WHO World Standard Population is indicated. All rates are expressed per 100,000 population.
 2. Incidence numbers are based on state of usual residence at time of diagnosis; mortality numbers on state of registration of death.
- Source: *Cancer in Australia 2000*, AIHW & AACR, 2003.

Table 38: Cancer of other connective and soft tissue (ICD-10 C49)

Australia 2000												
Age group	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
0-4	1	0.2	2	0.3	3	0.2	0	0.0	0	0.0	0	0.0
5-9	2	0.3	0	0.0	2	0.1	1	0.1	1	0.2	2	0.1
10-14	4	0.6	8	1.2	12	0.9	0	0.0	1	0.2	1	0.1
15-19	4	0.6	8	1.2	12	0.9	3	0.4	1	0.2	4	0.3
20-24	5	0.8	10	1.6	15	1.2	2	0.3	1	0.2	3	0.2
25-29	8	1.1	12	1.7	20	1.4	2	0.3	3	0.4	5	0.3
30-34	11	1.6	4	0.6	15	1.1	3	0.4	1	0.1	4	0.3
35-39	7	0.9	10	1.3	17	1.1	2	0.3	1	0.1	3	0.2
40-44	9	1.3	16	2.2	25	1.7	2	0.3	4	0.5	6	0.4
45-49	14	2.1	10	1.5	24	1.8	3	0.4	4	0.6	7	0.5
50-54	17	2.7	5	0.8	22	1.7	8	1.3	4	0.6	12	1.0
55-59	21	4.3	18	3.8	39	4.0	7	1.4	8	1.7	15	1.6
60-64	26	6.5	15	3.8	41	5.1	11	2.7	8	2.0	19	2.4
65-69	34	10.2	20	5.8	54	8.0	13	3.9	8	2.3	21	3.1
70-74	35	11.7	31	9.3	66	10.4	9	3.0	8	2.4	17	2.7
75-79	19	8.7	23	8.0	42	8.3	9	4.1	12	4.2	21	4.1
80-84	16	13.4	19	10.0	35	11.3	7	5.9	12	6.3	19	6.1
85 and over	17	21.9	25	14.3	42	16.6	10	12.9	14	8.0	24	9.5
Total	250		236		486		92		91		183	

Rates per 100,000 with 95% confidence intervals (95% CI)

Crude rate	2.6	2.4	2.5	1.0	0.9	1.0
95% CI	2.3 - 3.0	2.1 - 2.8	2.3 - 2.8	0.8 - 1.2	0.7 - 1.1	0.8 - 1.1
AS rate (Aust. 2001)	2.9	2.3	2.6	1.1	0.9	1.0
95% CI	2.5 - 3.2	2.0 - 2.6	2.3 - 2.8	0.9 - 1.3	0.7 - 1.1	0.8 - 1.1
AS rate (new WHO World)	2.2	1.9	2.0	0.8	0.7	0.7
95% CI	1.9 - 2.5	1.6 - 2.2	1.9 - 2.2	0.6 - 1.0	0.5 - 0.8	0.6 - 0.8
Lifetime risk (0-74)	1 in 448	1 in 571	1 in 504	1 in 1,339	1 in 1,739	1 in 1,516
PYLL (0-74)				1,350	1,113	2,463
Per cent of all cancers	0.5	0.6	0.6	0.5	0.6	0.5

Average annual numbers and rates by State and Territory, 1996-2000

	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate
NSW	76	2.6	57	1.7	134	2.1	27	1.0	23	0.7	50	0.8
Vic	73	3.5	60	2.5	133	2.9	21	1.1	21	0.9	43	0.9
Qld	61	4.1	0	2.6	105	3.3	11	0.8	10	0.6	21	0.7
WA	0	2.7	12	1.4	34	2.0	7	1.0	4	0.4	11	0.7
SA	37	5.3	31	3.6	67	4.3	6	0.9	6	0.7	12	0.8
Tas	9	4.2	7	2.6	16	3.4	1	0.6	1	0.5	3	0.6
ACT	2	1.5	2	1.4	4	1.5	1	1.3	1	0.7	2	1.0
NT	2	2.1	1	2.4	3	2.3	0	0.4	1	1.2	1	0.8

Notes

- AS rates use Australian 2001 Standard Population unless new WHO World Standard Population is indicated. All rates are expressed per 100,000 population.
- Incidence numbers are based on state of usual residence at time of diagnosis; mortality numbers on state of registration of death.

Source: *Cancer in Australia 2000*, AIHW & AACR, 2003.

Table 39: Cancer of other connective and soft tissue and autonomic nervous system (ICD-10 C47–C49)

Australia 2000

Age group	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
0–4	5	0.8	4	0.6	9	0.7	0	0.0	0	0.0	0	0.0
5–9	2	0.3	2	0.3	4	0.3	1	0.1	1	0.2	2	0.1
10–14	5	0.7	8	1.2	13	1.0	1	0.1	1	0.2	2	0.1
15–19	4	0.6	10	1.5	14	1.1	3	0.4	1	0.2	4	0.3
20–24	5	0.8	11	1.7	16	1.2	2	0.3	1	0.2	3	0.2
25–29	10	1.4	13	1.8	23	1.6	3	0.4	3	0.4	6	0.4
30–34	13	1.8	8	1.1	21	1.5	4	0.6	1	0.1	5	0.4
35–39	8	1.1	12	1.6	20	1.3	3	0.4	3	0.4	6	0.4
40–44	13	1.8	20	2.7	33	2.3	6	0.8	7	1.0	13	0.9
45–49	14	2.1	14	2.1	28	2.1	3	0.4	6	0.9	9	0.7
50–54	19	3.0	14	2.2	33	2.6	10	1.6	6	1.0	16	1.3
55–59	24	4.9	25	5.3	49	5.1	9	1.8	9	1.9	18	1.9
60–64	27	6.7	28	7.1	55	6.9	13	3.2	12	3.0	25	3.1
65–69	36	10.8	35	10.1	71	10.5	19	5.7	16	4.6	35	5.2
70–74	40	13.4	45	13.5	85	13.4	13	4.3	15	4.5	28	4.4
75–79	23	10.5	39	13.6	62	12.2	11	5.0	17	5.9	28	5.5
80–84	16	13.4	30	15.8	46	14.9	8	6.7	21	11.1	29	9.4
85 and over	19	24.5	33	18.8	52	20.6	11	14.2	19	10.8	30	11.9
Total	283		351		634		120		139		259	

Rates per 100,000 with 95% confidence intervals (95% CI)

Crude rate	3.0	3.6	3.3	1.3	1.4	1.4
95% CI	2.6 – 3.3	3.3 – 4.0	3.1 – 3.6	1.0 – 1.5	1.2 – 1.7	1.2 – 1.5
AS rate (Aust. 2001)	3.2	3.5	3.3	1.4	1.3	1.4
95% CI	2.8 – 3.6	3.1 – 3.8	3.1 – 3.6	1.1 – 1.6	1.1 – 1.6	1.2 – 1.5
AS rate (new WHO World)	2.5	2.8	2.7	1.1	1.0	1.0
95% CI	2.2 – 2.8	2.5 – 3.1	2.4 – 2.9	0.9 – 1.2	0.8 – 1.2	0.9 – 1.1
Lifetime risk (0–74)	1 in 399	1 in 378	1 in 389	1 in 980	1 in 1,085	1 in 1,030
PYLL (0–74)				1,830	1,530	3,360
Per cent of all cancers	0.6	0.9	0.7	0.6	0.9	0.7

Average annual numbers and rates by State and Territory, 1996–2000

	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate
NSW	94	3.2	102	3.1	196	3.1	39	1.4	43	1.3	82	1.3
Vic	82	3.9	82	3.4	164	3.6	29	1.4	33	1.3	61	1.3
Qld	69	4.6	64	3.8	132	4.1	17	1.2	23	1.3	40	1.3
WA	25	3.2	19	2.2	44	2.7	12	1.6	8	1.0	20	1.2
SA	38	5.5	32	3.8	70	4.5	9	1.2	11	1.3	20	1.3
Tas	9	4.2	7	2.6	16	3.4	3	1.2	2	0.8	5	1.0
ACT	3	2.2	3	2.1	5	2.3	1	1.3	1	0.9	3	1.1
NT	3	2.8	1	2.4	4	2.7	1	2.0	1	1.2	1	1.4

Notes

1. AS rates use Australian 2001 Standard Population unless new WHO World Standard Population is indicated. All rates are expressed per 100,000 population.

2. Incidence numbers are based on state of usual residence at time of diagnosis; mortality numbers on state of registration of death.

Source: *Cancer in Australia 2000*, AIHW & AACR, 2003.

Table 40: Cancer of the breast (ICD-10 C50)

Australia 2000

Age group	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
0-4	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
5-9	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
10-14	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
15-19	0	0.0	2	0.3	2	0.2	0	0.0	0	0.0	0	0.0
20-24	0	0.0	4	0.6	4	0.3	0	0.0	1	0.2	1	0.1
25-29	1	0.1	54	7.4	55	3.8	0	0.0	5	0.7	5	0.3
30-34	0	0.0	190	26.5	190	13.3	1	0.1	27	3.8	28	2.0
35-39	2	0.3	435	57.5	437	29.0	2	0.3	60	7.9	62	4.1
40-44	3	0.4	840	115.2	843	58.2	0	0.0	126	17.3	126	8.7
45-49	6	0.9	1,270	188.4	1,276	95.1	1	0.1	188	27.9	189	14.1
50-54	7	1.1	1,562	250.7	1,569	124.8	0	0.0	255	40.9	255	20.3
55-59	5	1.0	1,432	302.4	1,437	149.1	3	0.6	246	52.0	249	25.8
60-64	8	2.0	1,338	337.2	1,346	168.7	2	0.5	237	59.7	239	30.0
65-69	9	2.7	1,118	324.0	1,127	166.4	2	0.6	215	62.3	217	32.0
70-74	19	6.3	1,100	329.7	1,119	176.7	1	0.3	294	88.1	295	46.6
75-79	15	6.8	891	309.7	906	178.6	4	1.8	280	97.3	284	56.0
80-84	6	5.0	567	298.4	573	185.5	2	1.7	240	126.3	242	78.3
85 and over	5	6.5	511	291.7	516	204.2	3	3.9	347	198.1	350	138.5
Total	86		11,314		11,400		21		2,521		2,542	

Rates per 100,000 with 95% confidence intervals (95% CI)

Crude rate	0.9	117.3	59.5	0.2	26.1	13.3
95% CI	0.7 – 1.1	115.1 – 119.4	58.4 – 60.6	0.1 – 0.3	25.1 – 27.1	12.8 – 13.8
AS rate (Aust. 2001)	1.0	115.3	60.3	0.3	24.7	13.5
95% CI	0.8 – 1.2	113.2 – 117.5	59.2 – 61.4	0.1 – 0.4	23.8 – 25.7	13.0 – 14.0
AS rate (new WHO World)	0.7	91.7	47.3	0.2	18.1	9.7
95% CI	0.6 – 0.8	90.0 – 93.4	46.4 – 48.2	0.1 – 0.3	17.4 – 18.9	9.3 – 10.0
Lifetime risk (0-74)	1 in 1,343	1 in 11	1 in 21	1 in 7,677	1 in 56	1 in 109
PYLL (0-74)				240	28,305	28,545
Per cent of all cancers	0.2	28.8	13.4	0.1	16.3	7.2

Average annual numbers and rates by State and Territory, 1996-2000

	Incidence						Mortality					
	Males		Females		Persons		Males		Females		Persons	
	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate	Number	AS rate
NSW	31	1.1	3,585	110.9	3,616	58.2	7	0.2	846	25.3	853	13.8
Vic	19	0.9	2,731	114.5	2,750	60.6	7	0.4	702	28.6	709	15.7
Qld	17	1.2	1,873	113.6	1,890	58.9	3	0.3	425	25.5	428	13.7
WA	5	0.7	949	111.5	954	57.8	1	0.1	221	25.7	222	13.9
SA	6	0.9	929	115.4	935	60.9	1	0.1	230	27.0	231	14.8
Tas	2	0.8	248	101.2	250	53.0	1	0.4	68	26.3	68	14.5
ACT	1	1.0	157	118.0	159	62.3	0	0.4	44	36.3	45	20.3
NT	1	1.2	50	93.1	50	44.0	0	0.0	13	32.3	13	15.5

Notes

1. AS rates use Australian 2001 Standard Population unless new WHO World Standard Population is indicated. All rates are expressed per 100,000 population.

2. Incidence numbers are based on state of usual residence at time of diagnosis; mortality numbers on state of registration of death.

Source: *Cancer in Australia 2000*, AIHW & AACR, 2003.