

Appendix B: Tables

This appendix contains 53 detailed tables. The first table shows age-standardised death rates for males and females since 1885. The second table shows ratios of male deaths to female deaths for specific age groups between 1907 and 2000. The next two tables are life expectancy tables for males and females at selected ages. All the following tables show death rates (age-standardised or age-specific) for all the causes described in this report, for males and females and by selected age groupings.

It is important to note that unless otherwise stated:

1. All information in these tables is sourced from the AIHW General Record of Incidence of Mortality (GRIM) Books.
2. All rates are expressed as per 100,000 population.
3. All age-standardised rates were calculated using the Australian population at 30 June 2001.

Table B1: Age-standardised all-cause death rates, 1885–2003

Year	Males	Females
1885 ^(a)	3,146.9	2,227.8
1895 ^(a)	2,606.2	2,069.0
1900 ^(a)	2,369.5	1,956.7
1905 ^(a)	2,301.9	1,900.6
1907	2,234.2	1,844.4
1910	2,106.4	1,735.9
1915	2,255.1	1,768.9
1920	2,182.4	1,745.3
1925	1,985.6	1,576.2
1930	1,757.5	1,429.0
1935	1,859.7	1,472.4
1940	1,836.4	1,394.7
1945	1,688.6	1,313.4
1950	1,717.7	1,254.5
1955	1,661.4	1,148.5
1960	1,620.2	1,091.9
1965	1,658.0	1,069.0
1970	1,719.3	1,084.2
1975	1,481.0	904.0
1980	1,338.7	794.3
1985	1,257.3	773.3
1990	1,095.0	682.3
1995	983.1	622.8
2000	853.3	552.0
2003	791.5	523.0

(a) Estimated from Lancaster (1990).

Table B2: Ratios of male to female age-specific and age-standardised death rates, 1907–2000

Year	0–4	5–14	15–24	25–44	45–64	65–84	85 plus	All ages
1907	1.18	1.09	1.06	1.04	1.39	1.23	1.13	1.21
1910	1.21	1.14	1.07	1.08	1.42	1.21	1.15	1.21
1920	1.23	1.13	1.19	1.09	1.51	1.23	1.15	1.25
1930	1.21	1.36	1.16	1.11	1.42	1.23	1.11	1.23
1940	1.26	1.33	1.52	1.22	1.54	1.32	1.16	1.32
1950	1.28	1.46	2.09	1.34	1.63	1.39	1.10	1.37
1960	1.22	1.46	2.62	1.57	1.80	1.52	1.15	1.48
1970	1.37	1.30	2.97	1.63	1.89	1.55	1.23	1.59
1980	1.29	1.49	2.95	2.01	2.01	1.67	1.24	1.69
1990	1.29	1.48	2.78	2.23	1.83	1.60	1.23	1.60
2000	1.24	1.38	2.50	2.17	1.65	1.56	1.21	1.55

Table B3: Expected age at death (life expectancy), males, 1881–2001

Year^(a) or mid-point of range of years	Range of years	At birth	At age 1	At age 30	At age 65	At age 85
1884	1881–1890	47.20	54.34	63.64	76.06	88.86
1885		49.14	56.11	64.38	76.16	88.83
1895	1891–1900	51.08	57.88	65.11	76.25	88.79
1895		52.45	58.91	65.58	76.27	88.74
1900		53.83	59.93	66.05	76.29	88.70
1905	1901–1910	55.20	60.96	66.52	76.31	88.65
1910		56.52	61.86	67.16	76.54	88.64
1915		57.83	62.77	67.80	76.78	88.63
1920	1920–1922	59.15	63.67	68.44	77.01	88.62
1925		61.32	65.08	69.17	77.21	88.76
1933	1932–1934	63.48	66.49	69.90	77.40	88.90
1936		64.34	67.08	70.07	77.35	88.88
1942		65.21	67.66	70.23	77.30	88.86
1947	1946–1948	66.07	68.25	70.40	77.25	88.84
1950		66.61	68.56	70.65	77.29	88.93
1954	1953–1955	67.14	68.86	70.90	77.33	89.01
1961	1960–1962	67.92	69.46	71.12	77.47	89.08
1966	1965–1967	67.63	69.07	70.72	77.16	89.07
1971	1970–1972	67.81	69.25	70.94	77.21	89.13
1976	1975–1977	69.56	70.62	72.18	78.13	89.45
1981	1980–1982	71.23	72.05	73.51	78.80	89.67
1986	1985–1987	72.74	73.50	74.84	79.60	89.89
1991	1990–1992	74.32	74.92	76.07	80.41	90.23
1996	1995–1997	75.69	76.15	77.26	81.21	90.40
2001	2000–2002	77.40	77.80	78.80	82.40	90.60

(a) Data for single years were estimated by AIHW and are based on the range of years.

Source: AIHW population data holdings.

Table B4: Expected age at death (life expectancy), females, 1881–2001

Year^(a) or mid-point of range of years	Range of years	At birth	At age 1	At age 30	At age 65	At age 85
1884	1881–1890	50.84	57.44	66.13	77.27	88.90
1885		52.80	59.17	67.00	77.51	89.01
1895	1891–1900	54.76	60.89	67.86	77.75	89.12
1895		56.12	61.89	68.35	77.79	89.14
1900		57.48	62.89	68.84	77.84	89.17
1905	1901–1910	58.84	63.89	69.33	77.88	89.19
1910		60.33	64.94	70.05	78.12	89.15
1915		61.82	65.98	70.76	78.36	89.10
1920	1920–1922	63.31	67.03	71.48	78.60	89.06
1925		65.23	68.35	72.13	78.88	89.18
1933	1932–1934	67.14	69.67	72.77	79.15	89.30
1936		68.30	70.60	73.21	79.25	89.31
1942		69.47	71.06	73.43	79.30	89.31
1947	1946–1948	70.63	72.45	74.08	79.44	89.32
1950		71.69	73.34	74.76	79.73	89.42
1954	1953–1955	72.75	74.22	75.43	80.02	89.52
1961	1960–1962	74.18	75.49	76.49	80.68	89.79
1966	1965–1967	74.15	75.39	76.34	80.70	89.85
1971	1970–1972	74.49	75.74	76.67	80.90	90.03
1976	1975–1977	76.56	77.47	78.26	82.13	90.49
1981	1980–1982	78.27	78.98	79.67	83.00	90.74
1986	1985–1987	79.20	79.83	80.49	83.56	91.09
1991	1990–1992	80.39	80.91	81.48	84.26	91.40
1996	1995–1997	81.37	81.78	82.30	84.88	91.53
2001	2000–2002	82.60	83.00	83.40	85.80	91.80

(a) Data for single years were estimated by AIHW and are based on the range of years.

Source: AIHW population data holdings

Table B5: Age-standardised death rates by cause of death, males, 1907–2003

Year	Circulatory diseases	Cancer	Respiratory diseases	Injury and poisoning	Infectious diseases	Other diseases	All causes
1907	437.1	165.5	319.8	146.5	282.9	882.4	2,234.2
1910	430.9	165.9	225.2	146.7	213.9	923.8	2,106.4
1915	444.7	162.8	258.4	144.8	218.0	1,026.4	2,255.1
1920	497.0	179.6	240.2	127.4	188.9	949.3	2,182.4
1925	509.3	200.5	233.5	126.8	148.6	766.9	1,985.6
1930	513.7	196.0	191.7	128.4	126.4	601.3	1,757.5
1935	677.9	202.6	234.2	113.5	102.8	528.7	1,859.7
1940	746.9	208.0	178.8	118.0	96.0	488.7	1,836.4
1945	795.7	200.5	133.3	85.4	74.4	399.3	1,688.6
1950	892.8	209.3	141.5	111.0	55.1	308.0	1,717.7
1955	924.3	222.4	132.5	115.7	30.7	235.8	1,661.4
1960	929.6	226.5	136.9	108.2	20.5	198.5	1,620.2
1965	964.3	237.8	157.0	113.3	13.8	171.8	1,658.0
1970	977.3	263.2	181.7	113.7	11.7	171.7	1,719.3
1975	823.3	268.6	133.1	99.9	8.1	148.0	1,481.0
1980	700.3	278.0	125.3	86.5	5.8	142.8	1,338.7
1985	612.2	287.3	126.8	75.2	5.7	150.1	1,257.3
1990	490.8	277.7	102.0	70.6	7.0	146.9	1,095.0
1995	414.6	270.6	85.6	60.6	8.4	143.3	983.1
2000	318.5	246.6	80.5	61.0	11.0	135.7	853.3
2003	281.8	234.4	75.7	55.7	10.7	133.2	791.5

Table B6: Age-standardised death rates by cause of death, females, 1907–2003

Year	Circulatory diseases	Cancer	Respiratory diseases	Injury and poisoning	Infectious diseases	Other diseases	All causes
1907	379.1	154.3	263.1	55.1	229.9	762.9	1,844.4
1910	416.7	161.4	189.6	48.4	174.3	745.5	1,735.9
1915	374.9	160.5	195.2	47.3	164.0	827.0	1,768.9
1920	427.4	170.5	201.1	44.4	147.2	754.7	1,745.3
1925	445.8	176.8	185.2	44.4	109.3	614.7	1,576.2
1930	457.8	177.8	144.6	42.6	92.6	513.6	1,429.0
1935	551.3	185.5	182.8	46.8	65.9	440.1	1,472.4
1940	599.9	181.2	125.9	51.3	57.2	379.2	1,394.7
1945	625.5	178.6	98.6	45.0	44.9	320.8	1,313.4
1950	691.8	173.5	86.7	46.0	25.8	230.7	1,254.5
1955	677.6	166.8	67.2	51.1	13.8	172.0	1,148.5
1960	662.6	159.5	62.8	51.1	10.2	145.7	1,091.9
1965	660.3	155.7	61.2	55.8	6.8	129.2	1,069.0
1970	662.4	165.6	63.8	53.6	7.0	131.8	1,084.2
1975	550.5	158.0	43.0	44.3	5.1	103.1	904.0
1980	452.2	157.9	42.1	36.8	3.9	101.4	794.3
1985	410.4	167.2	48.5	30.9	4.4	111.9	773.3
1990	337.2	162.6	43.4	27.9	4.6	106.6	682.3
1995	282.4	160.3	42.4	24.4	5.2	108.1	622.8
2000	223.5	147.9	44.1	24.9	7.0	104.6	552.0
2003	197.3	145.4	45.4	22.2	6.8	105.9	523.0

Table B7: Age-standardised circulatory diseases death rates, males, 1907–2003

Year	Cerebrovascular disease	IHD ^(a)	Other circulatory plus IHD	Other circulatory ^(a)	All circulatory
1907	126.7	..	310.4	..	437.1
1910	97.5	..	333.4	..	430.9
1915	112.4	..	332.3	..	444.7
1920	120.3	..	376.7	..	497.0
1925	126.1	..	383.2	..	509.3
1930	113.3	..	400.4	..	513.7
1935	126.7	..	551.2	..	677.9
1940	131.5	136.6	..	478.8	746.9
1945	140.6	162.0	..	493.1	795.7
1950	171.9	287.0	..	433.9	892.8
1955	197.9	364.5	..	361.9	924.3
1960	193.9	434.8	..	300.9	929.6
1965	203.8	512.3	..	248.2	964.3
1970	213.3	574.7	..	189.3	977.3
1975	182.9	484.5	..	155.9	823.3
1980	148.2	409.4	..	142.7	700.3
1985	121.3	372.8	..	118.1	612.2
1990	93.7	300.8	..	96.3	490.8
1995	85.3	248.1	..	81.2	414.6
2000	68.5	184.6	..	65.4	318.5
2003	59.9	160.8	..	61.1	281.8

(a) Ischaemic heart disease was not recorded in the ICD until 1950. See page 66 for discussion of estimates for 1940–1950.

Table B8: Age-standardised circulatory diseases death rates, females, 1907–2003

Year	Cerebrovascular disease	IHD ^(a)	Other circulatory plus IHD	Other circulatory ^(a)	All circulatory
1907	129.9	..	249.2	..	379.1
1910	111.0	..	305.7	..	416.7
1915	114.8	..	260.1	..	374.9
1920	117.1	..	310.3	..	427.4
1925	133.3	..	312.5	..	445.8
1930	130.4	..	327.4	..	457.8
1935	147.5	..	403.8	..	551.3
1940	152.2	63.8	..	383.9	599.9
1945	159.7	74.3	..	391.5	625.5
1950	200.4	140.1	..	351.3	691.8
1955	209.8	172.6	..	295.2	677.6
1960	199.5	213.6	..	249.5	662.6
1965	204.3	251.6	..	204.4	660.3
1970	210.9	297.9	..	153.6	662.4
1975	180.8	251.5	..	118.2	550.5
1980	138.8	207.1	..	106.3	452.2
1985	118.0	198.7	..	93.7	410.4
1990	90.8	172.0	..	74.4	337.2
1995	78.5	140.5	..	63.4	282.4
2000	63.2	108.0	..	52.3	223.5
2003	57.0	92.5	..	47.8	197.3

(a) Ischaemic heart disease was not recorded in the ICD until 1950. See page 66 for discussion of estimates for 1940–1950.

Table B9: Age-standardised cancer death rates, males, 1922–2003

Year	Colorectal cancer	Stomach cancer	Prostate cancer	Lung cancer	Other cancers and lung cancer	Other cancers	All cancers
1922	27.6	54.4	8.4	(a)	98.6	..	189.0
1925	28.3	53.7	14.6	(a)	103.9	..	200.5
1930	30.8	53.8	14.5	(a)	96.9	..	196.0
1935	29.9	53.5	19.7	(a)	99.5	..	202.6
1940	34.7	50.4	22.9	(a)	100.0	..	208.0
1945	34.6	45.7	24.0	10.9	..	85.3	200.5
1950	34.8	44.8	29.0	19.0	..	81.7	209.3
1955	32.6	40.5	33.4	30.0	..	85.9	222.4
1960	32.0	32.2	32.8	39.5	..	90.0	226.5
1965	31.9	27.0	33.9	53.6	..	91.4	237.8
1970	32.5	26.1	36.0	65.8	..	102.8	263.2
1975	36.5	22.3	34.3	72.3	..	103.2	268.6
1980	34.9	19.2	33.2	79.4	..	111.3	278.0
1985	37.9	17.2	35.7	77.2	..	119.3	287.3
1990	35.2	13.2	39.9	68.5	..	120.9	277.7
1995	33.9	11.9	41.4	63.9	..	119.5	270.6
2000	30.5	9.5	35.9	54.7	..	116.0	246.6
2003	26.6	7.8	34.2	49.1	..	116.7	234.4

(a) Lung cancer not separately identified before 1945.

Table B10: Age-standardised cancer death rates, females, 1922–2003

Year	Colorectal cancer	Cancers of the cervix and uterus	Breast cancer	Stomach cancer	Lung cancer	Other cancers and lung cancer	Other cancers	All cancers
1922	27.2	23.9	24.7	33.8	(a)	70.3	..	179.9
1925	27.8	22.7	28.0	31.7	(a)	66.6	..	176.8
1930	32.1	19.9	26.4	34.4	(a)	65.0	..	177.8
1935	30.5	22.4	30.7	32.1	(a)	69.8	..	185.5
1940	31.4	20.8	31.3	30.2	(a)	67.5	..	181.2
1945	33.2	19.9	32.0	27.8	3.3	..	62.4	178.6
1950	33.5	16.6	30.5	25.2	4.1	..	63.6	173.5
1955	31.4	15.3	31.2	22.5	4.5	..	61.9	166.8
1960	30.0	14.6	28.8	18.5	5.2	..	62.4	159.5
1965	29.4	13.1	28.3	14.8	6.5	..	63.6	155.7
1970	30.9	12.5	29.7	13.7	9.5	..	69.3	165.6
1975	29.4	10.6	29.2	11.3	10.5	..	67.0	158.0
1980	28.6	8.5	28.8	8.5	14.4	..	69.1	157.9
1985	28.0	9.0	31.3	7.4	17.1	..	74.4	167.2
1990	24.3	6.8	31.0	6.0	19.3	..	75.2	162.6
1995	22.6	6.8	29.5	4.9	21.9	..	74.6	160.3
2000	20.5	5.2	24.7	3.9	22.2	..	71.4	147.9
2003	17.6	4.8	24.6	4.1	22.2	..	72.1	145.4

(a) Lung cancer not separately identified before 1945.

Table B11: Age-standardised respiratory diseases death rates, males, 1907–2003

Year	Pneumonia	Influenza ^(a)	COPD	Other and COPD	Other	All respiratory
1907	116.4	62.0	..	141.4	..	319.8
1910	91.3	18.5	..	115.4	..	225.2
1915	121.3	16.5	..	120.6	..	258.4
1920	113.0	15.1	..	112.1	..	240.2
1925	104.0	8.7	27.7	..	93.1	233.5
1930	95.8	8.6	21.0	..	66.3	191.7
1935	128.0	33.5	15.4	..	57.3	234.2
1940	110.7	8.6	13.8	..	45.7	178.8
1945	85.3	2.9	13.7	..	31.4	133.3
1950	69.5	8.3	24.0	..	39.7	141.5
1955	63.9	2.0	31.5	..	35.1	132.5
1960	68.5	1.8	42.9	..	23.7	136.9
1965	69.8	2.8	62.0	..	22.4	157.0
1970	53.7	12.8	91.4	..	23.8	181.7
1975	34.5	2.3	79.7	..	16.6	133.1
1980	25.8	1.5	81.0	..	17.0	125.3
1985	21.0	4.0	81.4	..	20.4	126.8
1990	16.7	0.4	66.0	..	18.9	102.0
1995	13.3	0.6	55.2	..	16.5	85.6
2000	18.9	0.4	43.5	..	17.7	80.5
2003	19.7	0.4	37.6	..	18.0	75.7

(a) The Spanish influenza pandemic, which caused the deaths of over 12,000 persons, was in 1918–1919.

Table B12: Age-standardised respiratory diseases death rates, females, 1907–2003

Year	Pneumonia	Influenza ^(a)	COPD	Other and COPD	Other	All respiratory
1907	85.3	59.6	..	118.2	..	263.1
1910	72.9	17.6	..	99.1	..	189.6
1915	88.9	20.0	..	86.3	..	195.2
1920	94.6	14.3	..	92.2	..	201.1
1925	92.5	11.7	14.6	..	66.4	185.2
1930	76.4	6.3	13.5	..	48.4	144.6
1935	98.6	29.6	10.1	..	44.5	182.8
1940	77.2	6.5	9.4	..	32.8	125.9
1945	64.4	1.9	8.1	..	24.2	98.6
1950	48.6	7.0	7.0	..	24.1	86.7
1955	41.5	2.4	5.3	..	18.0	67.2
1960	42.2	2.3	5.9	..	12.4	62.8
1965	39.3	1.5	7.0	..	13.4	61.2
1970	31.7	7.9	11.2	..	13.0	63.8
1975	20.2	1.5	12.3	..	9.0	43.0
1980	15.3	1.4	13.7	..	11.7	42.1
1985	13.8	3.7	18.1	..	12.9	48.5
1990	11.5	0.5	18.9	..	12.5	43.4
1995	8.7	0.5	20.3	..	12.9	42.4
2000	13.2	0.4	18.5	..	12.0	44.1
2003	14.8	0.3	18.5	..	11.8	45.4

(a) The Spanish influenza pandemic, which caused the deaths of over 12,000 persons, was in 1918–1919.

Table B13: Age-standardised injury and poisoning death rates, males, 1907–2003

Year	Suicide	Motor vehicle accident deaths	Other including motor vehicle	Other injury and poisoning	All injury and poisoning
1907	26.7	..	119.8	..	146.5
1910	26.7	..	120.0	..	146.7
1915	29.1	..	115.7	..	144.8
1920	26.4	..	101.0	..	127.4
1925	25.2	16.2	..	85.4	126.8
1930	29.8	25.8	..	72.8	128.4
1935	22.0	26.3	..	65.2	113.5
1940	18.6	34.3	..	65.1	118.0
1945	12.8	17.1	..	55.5	85.4
1950	16.7	39.0	..	55.3	111.0
1955	18.1	41.8	..	55.8	115.7
1960	18.0	43.0	..	47.2	108.2
1965	22.3	45.8	..	45.2	113.3
1970	20.6	48.9	..	44.2	113.7
1975	17.6	41.6	..	40.7	99.9
1980	18.1	35.2	..	33.2	86.5
1985	19.1	26.5	..	29.6	75.2
1990	21.0	21.2	..	28.4	70.6
1995	21.1	16.2	..	23.3	60.6
2000	19.8	14.3	..	26.9	61.0
2003	17.7	12.3	..	25.7	55.7

Table B14: Age-standardised injury and poisoning death rates, females, 1907–2003

Year	Suicide	Motor vehicle accident deaths	Other including motor vehicle	Other injury and poisoning	All injury and poisoning
1907	5.2	..	49.9	..	55.1
1910	4.9	..	43.5	..	48.4
1915	6.1	..	41.2	..	47.3
1920	5.4	..	39.0	..	44.4
1925	5.3	4.6	..	34.5	44.4
1930	5.3	7.5	..	29.8	42.6
1935	6.0	6.9	..	33.9	46.8
1940	5.5	9.3	..	36.5	51.3
1945	5.1	4.8	..	35.1	45.0
1950	5.1	7.5	..	33.4	46.0
1955	6.1	10.1	..	34.9	51.1
1960	7.1	14.2	..	29.8	51.1
1965	12.5	15.1	..	28.2	55.8
1970	8.8	17.5	..	27.3	53.6
1975	8.0	13.9	..	22.4	44.3
1980	6.1	12.8	..	17.9	36.8
1985	5.3	10.9	..	14.7	30.9
1990	5.1	8.8	..	14.0	27.9
1995	5.5	7.2	..	11.7	24.4
2000	5.2	5.5	..	14.2	24.9
2003	4.7	4.4	..	13.1	22.2

Table B15: Age-standardised infectious diseases death rates, males, 1907–2003

Year	Septicaemia	Tuberculosis	Diarrhoea	Other infectious	All infectious
1907	4.4	120.5	60.3	102.8	288.0
1910	3.2	103.9	60.3	56.5	223.9
1915	5.1	94.3	73.7	51.8	224.9
1920	3.4	90.6	58.8	45.8	198.6
1925	2.8	76.7	38.8	30.3	148.6
1930	1.9	66.8	29.0	28.7	126.4
1935	2.3	59.5	13.4	27.6	102.8
1940	1.3	51.2	14.0	29.5	96.0
1945	1.0	46.4	9.5	17.5	74.4
1950	0.9	34.8	5.3	14.1	55.1
1955	0.7	16.1	4.7	9.2	30.7
1960	0.6	10.2	4.7	5.0	20.5
1965	0.6	6.6	2.7	3.9	13.8
1970	0.5	4.1	3.5	3.6	11.7
1975	0.6	2.5	2.9	2.1	8.1
1980	1.4	1.9	1.2	1.3	5.8
1985	2.1	0.9	0.9	1.8	5.7
1990	3.4	0.9	0.8	1.9	7.0
1995	3.9	0.7	0.7	3.1	8.4
2000	6.3	0.4	0.6	3.7	11.0
2003	6.3	0.3	0.8	3.3	10.7

Table B16: Age-standardised infectious diseases death rates, females, 1907–2003

Year	Septicaemia	Tuberculosis	Diarrhoea	Other infectious	All infectious
1907	3.5	93.1	60.3	71.9	228.8
1910	3.3	80.4	60.3	34.3	178.3
1915	4.9	68.6	73.5	21.9	168.9
1920	2.7	57.3	58.8	32.4	151.2
1925	2.9	51.8	38.8	15.8	109.3
1930	1.7	45.1	33.2	12.6	92.6
1935	1.0	35.7	12.1	17.1	65.9
1940	0.8	29.4	11.0	16.0	57.2
1945	0.4	25.3	8.7	10.5	44.9
1950	0.3	13.3	5.1	7.1	25.8
1955	0.6	4.6	3.9	4.7	13.8
1960	0.6	2.9	3.0	3.7	10.2
1965	0.4	1.5	2.2	2.7	6.8
1970	0.5	0.9	3.1	2.5	7.0
1975	0.7	0.5	2.3	1.6	5.1
1980	1.4	0.4	1.0	1.1	3.9
1985	2.0	0.5	1.0	0.9	4.4
1990	2.4	0.4	1.0	0.8	4.6
1995	2.8	0.3	0.7	1.4	5.2
2000	4.2	0.2	0.7	1.9	7.0
2003	4.5	0.2	0.8	1.3	6.8

Table B17: Age-standardised 'other'^(a) diseases death rates, males, 1907–2003

Year	Senility	Mental and behavioural disorders^(b)	Diseases of the nervous system^(b)	Other	Total 'other'
1907	397.8	479.5	877.3
1910	408.7	505.1	913.8
1915	500.4	519.1	1,019.5
1920	452.3	487.4	939.7
1925	342.5	424.2	766.7
1930	230.4	370.9	601.3
1935	170.7	358.1	528.8
1940	132.5	356.3	488.8
1945	110.3	289.0	399.3
1950	66.8	241.2	308.0
1955	28.4	207.5	235.9
1960	19.3	179.2	198.5
1965	7.1	164.8	171.9
1970	7.0	10.0	13.4	141.3	171.7
1975	3.7	13.0	13.0	118.3	148.0
1980	1.4	11.5	13.9	116.0	142.8
1985	1.0	16.4	19.1	113.7	150.2
1990	0.7	17.7	19.3	109.2	146.9
1995	0.4	22.1	21.4	99.4	143.3
2000	0.3	17.6	23.8	94.0	135.7
2003	0.2	15.6	22.5	95.0	133.3

(a) 'Other' diseases include broad causes as described in Chapter 4 Broad causes of mortality-Other diseases page 38.

(b) Included in 'other' from 1968.

Table B18: Age-standardised 'other'^(a) diseases death rates, females, 1907–2003

Year	Senility	Mental and behavioural disorders^(b)	Diseases of the nervous system^(b)	Other	Total 'other'
1907	388.2	375.8	764.0
1910	349.8	391.7	741.5
1915	427.7	394.3	822.0
1920	378.4	372.4	750.8
1925	279.4	335.4	614.8
1930	211.8	301.9	513.7
1935	146.3	293.7	440.0
1940	110.0	269.2	379.2
1945	92.0	228.8	320.8
1950	62.4	168.4	230.8
1955	28.2	143.9	172.1
1960	18.9	126.8	145.7
1965	8.1	121.1	129.2
1970	5.7	7.1	9.5	109.5	131.8
1975	2.9	7.4	9.2	83.6	103.1
1980	1.9	7.5	9.5	82.5	101.4
1985	1.0	12.0	12.1	86.7	111.8
1990	0.9	13.5	14.3	77.9	106.6
1995	0.6	18.2	16.6	72.8	108.2
2000	0.6	14.6	19.5	70.0	104.7
2003	0.7	14.9	19.5	70.8	105.9

(a) 'Other' diseases include broad causes as described in Chapter 4 Broad causes of mortality-Other diseases page 38.

(b) Included in 'other' from 1968.

Table B19: Death rates, males aged 0–4, 1907–2003

Year	Injury and poisoning	Diarrhoea ^(a)	Infectious diseases	Respiratory diseases	SIDS	Cancer	Circulatory diseases	Congenital conditions	Conditions originating in the perinatal period	Other	All causes
1907	96.8	700.0	314.8	329.0	..	5.6	28.5	102.6	700.0	1,026.7	2,604.0
1910	80.9	700.0	383.6	197.5	..	6.8	22.0	102.6	700.0	938.9	2,432.3
1915	71.5	590.3	390.1	266.1	..	6.7	21.8	113.8	780.9	812.1	2,463.0
1920	65.9	617.1	435.1	244.9	..	8.6	14.8	106.5	792.7	806.3	2,474.8
1925	60.3	263.2	439.7	238.1	..	7.6	4.6	94.6	620.9	155.7	1,621.5
1930	70.3	217.8	382.6	197.6	..	12.0	9.8	106.5	561.0	126.0	1,465.8
1935	67.9	82.5	201.2	206.7	..	8.4	10.5	118.1	519.8	93.0	1,225.6
1940	63.1	90.8	196.7	188.8	..	13.7	6.2	122.3	518.8	105.3	1,214.9
1945	54.8	41.2	101.4	122.1	..	12.5	5.1	107.4	462.1	80.1	945.5
1950	54.3	34.9	70.1	68.1	..	11.4	3.1	87.5	369.7	65.9	730.1
1955	57.2	19.3	41.8	69.5	..	12.4	4.5	100.2	305.8	55.6	647.0
1960	45.5	16.3	28.4	59.1	..	10.9	4.6	97.0	276.6	44.5	566.6
1965	46.4	18.1	26.8	55.0	..	10.9	2.5	75.1	235.1	29.4	481.2
1970	47.9	17.4	28.8	62.0	4.1	8.9	3.6	94.5	252.8	29.4	532.0
1975	42.9	7.3	14.1	26.4	30.9	6.4	2.0	71.8	152.3	19.6	366.4
1980	33.5	2.9	5.5	10.3	41.7	6.4	3.1	67.9	107.6	15.6	291.6
1985	25.7	1.1	3.4	7.8	55.5	5.7	3.9	68.7	89.6	12.4	272.7
1990	22.6	0.6	4.0	5.4	45.7	5.9	1.7	48.5	82.5	13.1	229.4
1995	17.1	0.4	2.8	5.4	18.3	4.8	1.2	34.8	57.0	10.5	151.9
2000	15.9	0.2	2.6	2.3	12.4	2.7	1.5	29.7	54.6	12.6	134.3
2003	11.3	0.0	2.8	4.8	6.8	3.4	4.0	26.2	52.3	16.0	127.6

(a) Depending on the year, diarrhoea has been included as part of the broader groupings of either infectious disease or 'other' diseases. Its contribution to these broader groupings is included in the rates presented for them here. However, diarrhoea mortality rates are also shown separately in this table because it had such a large impact on deaths in the early twentieth century. Therefore, the all-causes rate shown here is the sum of all the columns except that for diarrhoea.

Table B20: Death rates, females aged 0–4, 1907–2003

Year	Injury and poisoning	Diarrhoea	Infectious diseases	Respiratory diseases	SIDS	Cancer	Circulatory diseases	Congenital conditions	Conditions originating in the perinatal period	Other	All causes
1907	79.6	578.5	493.5	267.3	..	5.4	26.9	66.6	595.6	678.8	2,213.7
1910	70.7	578.5	343.1	164.3	..	5.9	24.0	66.6	595.6	741.5	2,011.7
1915	54.8	499.6	349.9	212.9	..	4.4	16.1	86.2	645.3	647.3	2,016.9
1920	47.5	505.5	393.4	222.9	..	5.8	13.0	94.3	572.5	660.1	2,009.5
1925	54.1	201.7	382.2	207.4	..	6.6	8.5	84.6	481.6	113.8	1,338.8
1930	50.3	175.9	336.1	179.9	..	7.9	6.2	82.9	445.6	98.3	1,207.2
1935	45.8	62.4	192.9	166.4	..	9.5	7.2	92.3	410.4	78.5	1,003.0
1940	48.5	70.9	161.1	145.0	..	13.9	6.1	100.1	413.8	74.8	963.3
1945	36.3	38.7	88.5	99.1	..	12.4	3.5	101.2	351.4	53.5	745.9
1950	37.5	28.3	53.7	54.8	..	8.3	3.5	80.2	278.5	51.8	568.3
1955	40.9	19.7	37.6	55.5	..	14.4	3.7	90.1	227.1	37.4	506.7
1960	37.4	13.3	24.6	52.9	..	11.1	4.3	83.4	216.3	32.7	462.7
1965	30.7	13.7	21.3	43.7	..	10.0	1.9	71.7	179.1	23.1	381.5
1970	34.0	15.2	20.9	46.6	2.6	8.6	2.8	76.9	172.3	23.8	388.5
1975	28.1	6.5	11.5	16.9	22.8	6.7	3.0	61.0	107.7	14.8	272.5
1980	24.8	1.6	3.6	10.1	28.1	5.4	2.5	57.6	81.3	12.7	226.1
1985	18.8	1.7	3.4	6.3	35.5	5.1	2.6	58.3	75.0	10.9	215.9
1990	15.5	0.8	3.1	6.0	34.4	2.8	2.0	45.8	59.9	8.2	177.7
1995	11.7	0.3	1.1	6.8	13.3	3.6	1.6	33.0	45.4	7.1	123.6
2000	10.6	0.2	2.9	1.9	7.7	2.2	1.8	25.4	45.3	12.1	109.9
2003	12.7	0.2	2.1	3.1	4.7	1.8	2.3	22.2	42.8	12.5	104.2

(a) Depending on the year, diarrhoea has been included as part of the broader groupings of either infectious disease or 'other' diseases. Its contribution to these broader groupings is included in the rates presented for them here. However, diarrhoea mortality rates are also shown separately in this table because it had such a large impact on deaths in the early twentieth century. Therefore, the all-causes rate shown here is the sum of all the columns except that for diarrhoea.

Table B21: Death rates, males aged 5–14, 1907–2003

Year	Infectious diseases	Injury and poisoning	Respiratory diseases	Cancer	Circulatory diseases	Other	All causes
1907	67.6	45.5	19.2	3.8	16.1	35.1	187.3
1910	71.4	41.0	15.9	4.9	15.7	36.2	185.1
1915	78.4	38.1	21.2	3.0	18.1	36.9	195.7
1920	76.8	41.4	19.7	2.5	17.4	36.3	194.1
1925	32.1	40.7	16.0	4.5	20.8	41.0	155.1
1930	29.6	37.2	13.1	6.5	13.1	35.6	135.1
1935	34.3	41.2	18.2	5.2	14.5	32.2	145.6
1940	21.4	36.7	12.0	8.6	12.2	30.0	120.9
1945	18.5	38.0	8.0	8.3	11.2	21.3	105.3
1950	12.7	34.3	6.4	9.0	3.4	17.5	83.3
1955	5.2	26.1	4.9	8.0	2.8	12.1	59.1
1960	2.0	26.1	2.5	9.1	2.7	10.4	52.8
1965	1.9	24.2	3.3	8.1	1.4	7.8	46.7
1970	0.6	22.3	3.4	7.0	1.5	8.7	43.5
1975	0.8	21.4	1.5	5.6	0.8	6.9	37.0
1980	1.4	19.7	1.4	5.4	1.0	4.1	33.0
1985	0.6	15.1	1.2	5.4	1.2	5.6	29.1
1990	0.2	13.0	1.3	3.2	0.8	3.7	22.2
1995	0.5	8.5	0.5	4.2	0.8	3.7	18.2
2000	0.4	8.1	0.4	2.8	0.7	3.8	16.2
2003	< 0.1	5.0	0.6	3.0	0.3	3.4	12.4

Table B22: Death rates, females aged 5–14, 1907–2003

Year	Infectious diseases	Injury and poisoning	Respiratory diseases	Cancer	Circulatory diseases	Other	All causes
1907	73.9	21.7	23.0	2.1	20.7	30.3	171.7
1910	72.7	20.8	15.8	2.7	19.7	30.6	162.3
1915	86.0	15.0	22.3	1.0	21.8	36.5	182.6
1920	75.7	16.5	21.6	2.8	16.0	38.9	171.5
1925	29.1	17.8	13.0	3.1	20.9	29.1	113.0
1930	26.5	15.5	14.1	4.7	11.4	27.5	99.7
1935	24.5	14.8	17.0	3.9	12.3	29.9	102.4
1940	23.4	13.8	11.2	5.8	10.6	26.3	91.1
1945	15.1	13.2	6.8	6.3	11.4	19.7	72.5
1950	9.8	12.7	7.0	5.1	6.3	16.1	57.0
1955	4.3	10.9	4.0	6.0	3.5	13.6	42.3
1960	2.0	12.1	4.0	6.6	1.9	9.5	36.1
1965	1.4	10.8	3.3	6.4	1.1	8.4	31.4
1970	0.9	13.9	2.5	7.0	1.1	8.0	33.4
1975	0.6	10.2	1.9	5.2	1.1	5.4	24.4
1980	0.6	10.1	0.9	5.0	1.0	4.6	22.2
1985	0.4	7.1	1.0	3.8	0.8	4.7	17.8
1990	0.4	7.2	0.5	3.2	0.6	3.1	15.0
1995	0.4	6.0	1.0	2.6	1.0	5.3	16.3
2000	0.5	3.9	0.6	2.7	0.4	3.6	11.7
2003	< 0.1	4.2	0.7	2.0	0.6	2.5	10.1

Table B23: Death rates, males aged 15–24, 1907–2003

Year	Suicide	Motor vehicle accident deaths	Other injury and poisoning	Tuberculosis	Other infectious diseases	Respiratory diseases	Cancer	Circulatory diseases	Other	All causes
1907	6.2	n.a.	77.9	61.5	46.5	31.1	5.5	28.5	59.0	316.2
1910	6.7	n.a.	76.2	56.1	47.9	19.6	4.0	27.2	59.3	297.0
1915	7.9	n.a.	75.1	55.7	89.5	38.0	4.1	27.3	62.2	359.8
1920	7.9	n.a.	66.2	47.2	29.7	29.5	5.2	24.5	67.4	277.6
1925	9.9	19.2	70.3	30.0	12.4	20.9	6.2	19.6	58.4	246.9
1930	9.5	33.1	56.3	25.2	11.9	15.4	7.0	14.4	44.8	217.6
1935	7.5	39.0	38.5	19.0	10.2	23.1	8.2	17.0	44.2	206.7
1940	7.0	53.4	49.1	14.5	8.0	16.0	8.9	16.0	36.1	209.0
1945	5.3	16.7	28.9	11.1	4.2	7.7	5.3	14.1	23.9	117.2
1950	7.0	75.0	36.2	3.4	7.6	4.4	9.5	9.8	22.0	174.9
1955	9.8	73.9	33.1	0.3	3.7	3.1	10.8	8.8	19.5	163.0
1960	6.8	70.0	31.3	0.1	1.7	4.2	9.2	6.0	13.9	143.2
1965	10.6	76.1	25.1	0.0	0.5	5.3	11.2	5.4	11.2	145.4
1970	12.5	97.4	26.2	< 0.1	1.6	4.1	9.4	4.9	13.6	169.8
1975	13.6	94.2	23.4	< 0.1	1.3	3.0	8.9	4.5	11.9	160.9
1980	17.5	80.5	23.0	0.0	0.5	2.1	7.9	3.1	9.6	144.2
1985	24.0	60.6	21.9	<0.1	0.6	2.5	8.6	3.8	13.3	135.4
1990	27.0	42.7	20.7	0.0	1.0	2.0	6.3	4.6	11.4	115.7
1995	25.4	33.1	18.1	0.0	0.9	1.2	6.3	3.2	14.1	102.3
2000	19.7	28.3	17.0	0.0	0.9	0.6	4.8	2.0	16.8	90.1
2003	18.0	24.2	15.2	0.0	0.6	1.4	4.6	3.3	9.3	76.6

Table B24: Death rates, females aged 15–24, 1907–2003

Year	Suicide	Motor vehicle accident deaths	Other injury and poisoning	Tuberculosis	Other infectious diseases	Respiratory diseases	Cancer	Circulatory diseases	Other	All causes
1907	4.1	n.a.	16.5	98.1	37.4	22.8	2.7	29.1	86.5	297.2
1910	4.1	n.a.	14.0	85.5	33.1	14.9	4.4	33.6	88.4	278.0
1915	4.4	n.a.	12.0	73.7	40.8	22.2	4.0	26.2	86.9	270.2
1920	3.0	n.a.	11.9	64.9	23.0	17.4	5.0	21.1	87.9	234.2
1925	3.0	1.0	13.2	71.7	11.0	13.0	2.4	20.3	81.0	216.6
1930	5.3	3.9	11.3	57.0	9.1	15.1	5.1	15.1	66.1	188.0
1935	3.9	5.7	7.9	34.5	6.5	18.0	8.1	20.2	64.0	168.8
1940	3.1	8.7	6.2	27.0	6.3	9.3	6.2	14.7	56.1	137.6
1945	1.6	4.7	5.7	23.3	2.9	9.2	5.2	11.0	35.3	98.9
1950	2.2	9.8	6.4	8.8	5.1	5.3	8.4	8.9	28.6	83.5
1955	1.2	10.3	3.8	1.2	2.4	6.3	6.9	6.5	19.5	58.1
1960	2.0	17.6	3.6	0.6	2.0	3.0	5.3	5.9	14.6	54.6
1965	6.4	19.1	3.3	0.0	1.0	4.3	6.4	4.6	11.9	57.0
1970	4.7	24.5	4.5	0.0	0.9	3.0	6.0	2.9	10.7	57.2
1975	3.9	20.1	6.3	0.0	0.4	1.7	7.0	3.1	8.4	50.9
1980	4.4	19.7	7.3	0.0	0.7	1.7	5.6	2.4	7.0	48.8
1985	4.9	17.9	6.1	0.0	0.8	2.1	5.2	2.5	8.2	47.7
1990	4.4	14.6	6.0	<0.1	0.4	2.1	4.0	1.8	8.2	41.6
1995	6.4	11.3	5.3	0.0	0.3	0.9	3.8	1.4	8.9	38.3
2000	5.8	9.8	6.3	0.0	0.4	0.5	4.2	1.0	8.0	36.0
2003	3.7	7.5	5.3	0.0	0.7	0.2	3.8	2.1	6.6	29.9

Table B25: Death rates, males aged 25–44, 1907–2003

Year	Suicide	Motor vehicle accident deaths	Other injury and poisoning	Tuberculosis	Other infectious diseases	Respiratory diseases	Cancer	Circulatory diseases	Other diseases	All causes
1907	23.6	n.a.	90.2	137.2	58.1	60.9	22.8	68.2	116.9	577.9
1910	26.9	n.a.	91.1	122.8	61.5	50.1	21.2	66.5	114.0	554.1
1915	31.8	n.a.	92.4	116.4	76.0	57.5	23.5	67.8	128.3	593.7
1920	26.2	n.a.	77.2	114.7	37.7	50.7	21.1	60.0	113.8	501.4
1925	25.0	13.7	65.2	97.4	27.7	44.7	21.5	53.1	108.4	456.7
1930	30.9	24.9	51.4	79.3	16.4	35.0	23.9	49.0	78.8	389.6
1935	20.6	23.1	46.9	59.2	15.7	45.4	25.5	53.6	82.6	372.6
1940	16.8	32.4	42.3	41.1	13.1	27.0	30.5	51.6	69.3	324.1
1945	8.9	13.6	28.8	35.4	5.3	14.8	25.1	50.4	43.9	226.2
1950	14.7	36.1	35.1	18.4	4.7	10.8	25.0	56.1	43.9	244.8
1955	16.1	40.3	37.9	4.8	3.2	10.3	28.8	51.4	35.9	228.7
1960	20.4	39.9	32.0	2.6	2.5	10.1	28.8	57.4	28.9	222.6
1965	26.8	41.0	33.8	0.8	1.2	8.9	32.1	63.8	28.4	236.8
1970	23.0	41.9	32.5	0.4	1.2	10.8	32.3	55.5	25.1	222.7
1975	20.8	38.7	29.4	0.2	1.3	5.8	27.6	43.3	24.3	191.4
1980	23.0	33.4	26.9	0.1	0.8	4.2	24.9	35.3	19.6	168.2
1985	24.7	26.4	23.1	<0.1	0.8	4.6	22.8	27.8	23.2	153.4
1990	27.4	21.6	24.1	0.3	2.1	3.5	25.2	22.5	31.8	158.5
1995	30.6	17.0	21.6	0.0	2.9	2.5	22.6	20.3	36.7	154.2
2000	31.7	16.3	29.0	0.0	4.5	2.9	18.5	19.5	26.3	148.7
2003	26.9	14.4	24.3	0.0	3.1	2.6	19.5	18.2	20.9	129.9

Table B26: Death rates, females aged 25–44, 1907–2003

Year	Suicide	Motor vehicle accident deaths	Other injury and poisoning	Tuberculosis	Other infectious diseases	Respiratory diseases	Breast cancer	Other cancers	Circulatory diseases	Other diseases	All causes
1907	5.1	n.a.	16.5	133.7	38.3	49.7	6.1	29.8	65.9	209.7	554.8
1910	4.8	n.a.	14.5	125.1	37.3	34.3	6.9	31.9	63.7	194.1	512.6
1915	8.1	n.a.	13.9	105.7	34.1	37.6	6.8	31.5	65.9	180.9	484.5
1920	8.2	n.a.	13.8	86.9	31.3	42.4	6.4	29.3	56.3	183.7	458.3
1925	8.1	2.2	10.2	72.8	15.8	30.1	8.2	31.1	52.9	169.9	401.3
1930	6.2	4.5	7.0	66.0	12.4	23.8	8.7	37.4	45.5	138.8	350.3
1935	9.2	4.1	8.2	54.1	8.5	37.2	10.3	38.5	48.1	126.6	344.8
1940	6.5	4.5	6.7	43.0	8.4	18.6	7.5	30.3	41.2	98.0	264.7
1945	5.2	3.6	5.7	37.8	5.8	13.7	8.5	32.6	43.5	80.2	236.6
1950	5.5	4.6	5.9	17.5	3.4	10.1	8.9	32.5	41.1	52.9	182.4
1955	7.1	4.9	6.1	4.9	1.8	7.1	8.6	30.3	32.9	39.8	143.5
1960	7.6	7.3	6.0	2.5	1.9	7.1	9.3	33.2	33.2	33.9	142.0
1965	15.1	8.8	8.3	0.9	1.4	10.2	9.6	28.8	32.6	27.5	143.2
1970	9.7	11.9	6.8	0.3	1.5	9.5	10.2	27.2	32.4	27.4	136.9
1975	9.7	9.5	7.5	<0.1	1.4	5.2	8.4	21.7	23.9	18.2	105.5
1980	8.1	9.2	6.6	<0.1	0.6	3.1	7.9	21.5	13.6	13.0	83.6
1985	5.4	8.1	5.9	<0.1	0.6	3.6	9.6	19.9	12.2	11.7	77.0
1990	6.5	6.1	6.1	<0.1	0.5	2.8	9.3	20.3	8.0	11.5	71.1
1995	7.3	6.0	6.1	<0.1	0.8	2.0	7.5	17.6	8.1	13.4	68.8
2000	8.1	4.9	8.6	<0.1	1.0	2.2	7.4	15.3	8.2	12.7	68.8
2003	7.2	3.3	6.7	0.0	1.2	1.6	7.3	16.4	7.5	13.5	64.7

Table B27: Death rates, males aged 45–64, 1907–2003

Year	Infectious diseases	Injury and poisoning	Respiratory diseases	Cancer	Circulatory diseases	Other	All causes
1907	286.4	194.0	212.6	209.8	379.4	435.9	1,718.1
1910	270.5	206.2	148.6	217.4	368.0	437.3	1,648.0
1915	261.7	187.4	184.1	217.7	410.9	514.4	1,776.2
1920	225.6	157.9	163.2	246.5	438.0	466.4	1,697.6
1925	182.8	146.5	177.4	251.2	414.4	408.1	1,580.4
1930	157.5	162.4	136.7	221.6	400.9	312.2	1,391.3
1935	147.6	139.5	186.7	213.3	497.6	305.7	1,490.4
1940	148.4	136.6	134.4	211.8	574.9	283.3	1,489.4
1945	118.3	94.5	81.2	217.2	642.2	229.6	1,383.0
1950	86.2	110.1	93.7	229.6	742.5	193.6	1,455.7
1955	39.2	118.2	85.4	236.4	707.4	158.6	1,345.2
1960	22.5	110.7	74.5	246.9	721.6	131.6	1,307.8
1965	12.7	121.2	93.3	255.0	781.6	130.7	1,394.5
1970	9.2	118.0	109.5	290.4	758.9	129.8	1,415.8
1975	5.5	95.3	71.2	293.6	656.8	135.9	1,258.3
1980	4.6	83.5	56.7	314.4	532.1	117.6	1,108.9
1985	3.7	71.0	58.4	331.0	423.8	97.5	985.4
1990	5.1	63.0	43.1	288.4	307.1	91.0	797.7
1995	5.9	53.0	29.8	241.5	208.8	81.9	620.9
2000	6.9	48.7	24.2	210.8	145.1	73.9	509.6
2003	7.6	47.3	21.3	203.9	133.3	78.5	491.9

Table B28: Death rates, females aged 45–64, 1907–2003

Year	Infectious diseases	Injury and poisoning	Respiratory diseases	Cancer	Circulatory diseases	Other	All causes
1907	153.2	39.7	132.2	243.8	312.2	359.4	1,240.5
1910	141.1	35.6	92.5	228.7	314.9	346.4	1,159.2
1915	126.3	38.3	103.9	239.2	308.5	355.6	1,171.8
1920	113.6	33.1	96.5	245.4	326.5	308.3	1,123.4
1925	85.8	29.9	85.2	251.9	304.3	302.5	1,059.6
1930	72.2	33.9	65.6	243.2	309.7	257.3	981.9
1935	54.3	28.2	96.6	242.5	337.9	245.8	1,005.3
1940	46.9	30.4	64.3	249.6	354.2	219.1	964.5
1945	37.1	27.5	46.9	244.6	383.7	190.7	930.5
1950	24.3	27.8	37.8	230.6	425.8	144.1	890.4
1955	12.8	35.2	28.2	218.5	381.0	115.6	791.3
1960	8.0	43.6	25.9	205.1	347.6	98.2	728.4
1965	6.9	55.2	30.3	202.0	337.7	100.3	732.4
1970	6.3	48.7	39.0	228.2	326.2	101.4	749.8
1975	4.0	41.3	26.0	223.5	271.5	85.3	651.6
1980	3.0	32.0	27.0	221.8	201.3	67.1	552.2
1985	3.5	28.2	33.6	241.9	161.3	61.5	530.0
1990	2.7	23.2	26.0	223.1	110.7	50.0	435.7
1995	2.3	20.7	22.1	195.5	73.9	45.0	359.5
2000	3.4	17.8	17.7	170.9	55.1	44.8	309.7
2003	3.2	16.5	17.5	168.2	48.6	41.4	295.4

Table B29: Cancer death rates, males aged 45–64, 1922–2003

Year	Colorectal cancer	Prostate cancer	Stomach cancer	Lung cancer	Other cancers and lung cancer	Other cancers	All cancers
1922	31.4	6.2	90.0	(a)	127.1	..	254.7
1925	30.9	8.5	80.9	(a)	130.9	..	251.2
1930	30.4	11.6	65.8	(a)	113.8	..	221.6
1935	33.9	7.9	62.4	(a)	109.1	..	213.3
1940	34.0	10.7	55.3	(a)	111.8	..	211.8
1945	35.7	10.2	52.6	23.9	..	94.8	217.2
1950	33.5	9.2	45.4	41.6	..	99.9	229.6
1955	29.6	8.8	41.6	55.6	..	100.8	236.4
1960	29.4	10.1	28.6	69.3	..	109.5	246.9
1965	28.9	7.5	23.7	81.4	..	113.5	255.0
1970	32.4	9.1	25.2	95.9	..	127.8	290.4
1975	37.9	9.1	20.0	102.1	..	124.5	293.6
1980	40.7	10.5	19.4	107.9	..	135.9	314.4
1985	47.3	10.4	17.0	108.2	..	148.1	331.0
1990	38.4	10.4	11.9	83.3	..	144.4	288.4
1995	35.6	11.6	9.6	63.8	..	120.9	241.5
2000	31.2	8.3	7.7	50.6	..	113.0	210.8
2003	27.0	9.1	6.7	45.7	..	115.4	203.9

(a) Lung cancer not separately identified before 1945.

Table B30: Cancer death rates, females aged 45–64, 1922–2003

Year	Colorectal cancer	Cancers of the cervix and uterus	Breast cancer	Stomach cancer	Lung cancer	Other cancers and lung cancer	Other cancers	All cancers
1922	37.4	51.6	48.9	39.6	(a)	86.1	..	263.6
1925	30.3	42.3	46.1	41.3	(a)	91.9	..	251.9
1930	32.4	36.6	48.9	35.7	(a)	89.6	..	243.2
1935	32.0	38.9	52.4	27.1	(a)	92.1	..	242.5
1940	34.6	41.0	54.2	29.4	(a)	90.4	..	249.6
1945	34.7	39.5	55.9	25.5	5.8	..	83.2	244.6
1950	32.9	31.5	50.9	20.5	7.0	..	87.8	230.6
1955	33.6	25.6	54.0	17.0	6.6	..	81.7	218.5
1960	28.9	23.7	47.1	10.3	7.9	..	87.2	205.1
1965	28.0	22.5	47.9	10.2	10.4	..	83.0	202.0
1970	32.5	19.8	52.2	10.8	18.6	..	94.3	228.2
1975	33.7	16.9	51.6	10.7	19.3	..	91.3	223.5
1980	33.3	12.9	51.8	7.0	26.8	..	90.0	221.8
1985	34.9	12.6	59.1	6.2	30.4	..	98.7	241.9
1990	29.7	10.0	56.6	6.5	30.0	..	90.3	223.1
1995	23.6	10.4	52.2	4.0	28.3	..	77.0	195.5
2000	21.1	6.6	43.3	3.4	27.2	..	69.3	170.9
2003	17.0	6.8	43.2	3.8	27.0	..	70.4	168.2

(a) Lung cancer not separately identified before 1945.

Table B31: Circulatory diseases death rates, males aged 45–64, 1907–2003

Year	Cerebrovascular disease	IHD^(a)	Other circulatory plus IHD	Other circulatory	All circulatory
1907	100.4	..	279.0	..	379.4
1910	81.5	..	286.5	..	368.0
1915	92.3	..	318.6	..	410.9
1920	100.3	..	337.7	..	438.0
1925	92.9	..	321.5	..	414.4
1930	86.1	..	314.8	..	400.9
1935	94.8	..	402.8	..	497.6
1940	102.5	183.5	..	288.9	574.9
1945	120.1	231.3	..	290.8	642.2
1950	143.5	357.8	..	241.2	742.5
1955	131.4	414.3	..	161.7	707.4
1960	116.9	476.4	..	128.3	721.6
1965	112.7	555.9	..	113.0	781.6
1970	105.3	568.1	..	85.5	758.9
1975	96.5	491.2	..	69.1	656.8
1980	74.1	388.1	..	69.9	532.1
1985	51.7	316.3	..	55.8	423.8
1990	37.6	221.3	..	48.2	307.1
1995	25.8	147.9	..	35.1	208.8
2000	17.4	102.7	..	25.0	145.1
2003	16.7	92.1	..	24.5	133.3

(a) Ischaemic heart disease was not recorded in the ICD until 1950. See page 66 for discussion of estimates for 1940–1950.

Table B32: Circulatory diseases death rates, females aged 45–64, 1907–2003

Year	Cerebrovascular disease	IHD ^(a)	Other circulatory plus IHD	Other circulatory	All circulatory
1907	88.6	..	223.6	..	312.2
1910	81.0	..	233.9	..	314.9
1915	82.3	..	226.2	..	308.5
1920	102.7	..	223.8	..	326.5
1925	98.6	..	205.7	..	304.3
1930	104.3	..	205.4	..	309.7
1935	111.7	..	226.2	..	337.9
1940	115.5	56.5	..	182.2	354.2
1945	139.3	62.2	..	182.2	383.7
1950	162.1	117.1	..	146.6	425.8
1955	140.2	130.0	..	110.8	381.0
1960	111.7	148.3	..	87.6	347.6
1965	103.6	161.5	..	72.6	337.7
1970	95.3	175.6	..	55.3	326.2
1975	77.4	151.2	..	42.9	271.5
1980	54.0	114.0	..	33.3	201.3
1985	39.5	93.6	..	28.2	161.3
1990	24.6	63.4	..	22.7	110.7
1995	16.5	40.4	..	17	73.9
2000	13.7	27.3	..	14.1	55.1
2003	11.9	23.5	..	13.2	48.6

(a) Ischaemic heart disease was not recorded in the ICD until 1950. See page 66 for discussion of estimates for 1940–1950.

Table B33: Death rates, males aged 65–84, 1907–2003

Year	Senility	Infectious diseases	Injury and poisoning	Respiratory diseases	Cancer	Circulatory diseases	Other conditions	All causes
1907	1,396.9	460.2	281.8	1,150.0	783.3	1,976.8	1,691.9	7,740.9
1910	1,442.6	390.0	279.8	877.8	758.2	2,021.9	1,594.3	7,364.6
1915	1,755.8	345.2	274.8	910.7	730.7	1,954.7	1,556.1	7,528.0
1920	1,444.9	288.3	230.6	794.5	801.3	2,221.5	1,462.7	7,243.8
1925	871.6	203.7	231.2	704.6	892.6	2,136.2	1,304.6	6,344.5
1930	586.2	202.4	207.2	566.6	901.2	2,187.1	1,183.3	5,834.0
1935	441.0	188.3	204.8	768.7	995.9	2,981.7	1,246.8	6,827.2
1940	373.9	194.8	214.0	588.3	1,011.7	3,356.0	1,249.6	6,988.3
1945	297.8	180.7	183.4	496.9	963.4	3,646.4	1,042.3	6,810.9
1950	157.5	166.8	217.8	561.7	985.9	4,033.8	828.6	6,952.1
1955	53.5	109.7	216.1	507.6	1,004.2	4,087.4	737.6	6,716.1
1960	35.5	75.2	195.0	566.4	1,033.9	4,186.2	619.9	6,712.1
1965	11.1	54.4	201.5	671.2	1,146.1	4,388.3	585.6	7,058.2
1970	..	44.6	188.6	842.7	1,251.1	4,497.5	580.5	7,420.4
1975	..	30.9	151.2	583.0	1,252.1	3,575.3	472.8	6,071.3
1980	..	19.9	113.7	565.9	1308.5	3,075.4	483.1	5,567.4
1985	..	23.3	102.7	577.4	1367.6	2,766.4	512.3	5,351.2
1990	..	25.3	104.0	476.8	1,356.9	2,243.4	497.7	4,705.4
1995	..	30.0	80.6	416.4	1,363.7	1,877.5	505.6	4,274.3
2000	..	40.4	82.0	378.1	1,293.4	1,429.9	517.7	3,741.9
2003	..	42.4	78.1	361.1	1,238.3	1,236.6	530.4	3,486.9

Table B34: Death rates, females aged 65–84, 1907–2003

Year	Senility	Infectious diseases	Injury and poisoning	Respiratory diseases	Cancer	Circulatory diseases	Other conditions	All causes
1907	1,227.3	309.7	134.9	1,088.5	625.9	1,781.2	1,113.0	6,280.5
1910	1,171.2	327.6	130.8	735.2	669.2	2,036.0	1,013.3	6,083.3
1915	1,438.5	288.1	111.8	781.3	648.0	1,767.6	1,001.5	6,036.8
1920	1,238.0	240.0	96.0	746.2	694.6	1,951.6	926.4	5,892.8
1925	732.1	124.5	103.4	623.1	701.8	1,898.7	985.9	5,169.5
1930	497.5	114.4	87.3	447.8	698.2	1,975.7	906.0	4,726.9
1935	356.4	81.5	108.5	584.6	760.1	2,480.2	947.8	5,319.1
1940	300.2	93.4	147.8	415.5	761.2	2,679.4	877.2	5,274.7
1945	234.8	78.3	128.6	344.7	756.4	2,914.3	777.0	5,234.1
1950	147.1	49.5	116.9	296.2	717.3	3,112.2	559.3	4,998.5
1955	62.3	34.3	130.6	216.5	691.9	3,022.3	479.9	4,637.8
1960	32.1	27.0	121.9	198.8	642.5	2,946.2	434.1	4,402.6
1965	17.9	16.5	129.3	201.8	653.9	3,070.7	432.9	4,523.0
1970	..	20.2	128.1	242.8	696.2	3,206.1	472.2	4,777.3
1975	..	16.5	101.1	161.2	663.0	2,548.6	361.4	3,858.4
1980	..	15.1	75.9	165.2	675.4	2,020.0	369.0	3,324.2
1985	..	18.6	65.7	199.3	729.0	1,866.9	418.0	3,299.5
1990	..	18.2	63.2	196.1	724.7	1,544.0	399.5	2,946.7
1995	..	22.2	48.0	209.0	761.9	1,273.2	423.7	2,739.0
2000	..	32.1	50.2	208.7	718.5	977.7	416.2	2,403.7
2003	..	31.0	48.4	217.1	715.4	851.8	427.1	2,290.8

Table B35: Cancer death rates, males aged 65–84, 1922–2003

Year	Colorectal cancer	Prostate cancer	Stomach cancer	Lung cancer	Other cancers and lung cancer	Other cancers	All cancers
1922	132.2	48.5	252.0	(a)	445.2	. .	877.9
1925	147.3	73.3	251.5	(a)	420.5	. .	892.6
1930	144.3	80.9	273.8	(a)	402.2	. .	901.2
1935	138.9	116.1	282.0	(a)	458.9	. .	995.9
1940	171.1	138.2	273.9	(a)	428.5	. .	1,011.7
1945	176.0	140.2	238.8	39.7	. .	368.7	963.4
1950	168.8	155.4	242.9	81.7	. .	337.1	985.9
1955	159.6	173.5	192.9	138.1	. .	340.1	1,004.2
1960	148.8	165.0	165.3	185.1	. .	369.7	1,033.9
1965	160.5	172.1	136.3	279.3	. .	397.9	1,146.1
1970	158.9	181.4	124.8	347.9	. .	438.1	1,251.1
1975	163.8	163.6	109.1	378.7	. .	436.9	1,252.1
1980	158.6	164.4	90.6	412.9	. .	482.0	1,308.5
1985	177.6	174.8	82.6	402.0	. .	530.6	1,367.6
1990	178.5	203.9	65.8	369.4	. .	539.3	1,356.9
1995	167.6	209.7	60.2	361.0	. .	565.2	1,363.7
2000	158.2	190.4	51.3	316.6	. .	576.9	1,293.4
2003	140.0	181.0	41.3	289.7	. .	586.3	1,238.3

(a) Lung cancer not separately identified before 1945.

Table B36: Cancer death rates, females aged 65–84, 1922–2003

Year	Colorectal cancer	Cancers of the cervix and uterus	Breast cancer	Stomach cancer	Lung cancer	Other cancers and lung cancer	Other cancers	All cancers
1922	119.4	71.0	85.1	168.6	(a)	282.1	. .	726.2
1925	134.3	73.8	95.6	151.9	(a)	246.2	. .	701.8
1930	147.0	65.8	86.2	174.1	(a)	225.1	. .	698.2
1935	146.3	77.9	108.0	165.2	(a)	262.7	. .	760.1
1940	150.5	77.5	110.7	164.7	(a)	257.8	. .	761.2
1945	163.3	74.5	112.6	150.4	14.6	. .	241.0	756.4
1950	154.4	60.9	112.8	130.7	17.6	. .	240.9	717.3
1955	145.2	59.6	109.5	114.2	22.3	. .	241.1	691.9
1960	136.2	54.1	99.0	94.3	23.4	. .	235.5	642.5
1965	139.8	49.1	101.2	76.8	28.5	. .	258.5	653.9
1970	144.9	49.5	105.7	69.9	39.7	. .	286.5	696.2
1975	134.2	43.9	99.9	56.4	45.9	. .	282.7	663.0
1980	130.6	34.0	100.7	40.8	65.3	. .	304.0	675.4
1985	128.1	37.7	106.0	36.3	83.1	. .	337.8	729.0
1990	113.4	24.7	108.9	27.2	98.4	. .	352.1	724.7
1995	111.4	28.9	107.9	25.6	118.5	. .	369.6	761.9
2000	102.6	23.5	88.3	18.9	123.1	. .	362.1	718.5
2003	91.2	20.3	89.9	20.7	123.4	. .	369.9	715.4

(a) Lung cancer not separately identified before 1945.

Table B37: Circulatory diseases death rates, males aged 65–84, 1907–2003

Year	Cerebrovascular disease	IHD ^(a)	Other circulatory plus IHD	Other circulatory	All circulatory
1907	574.4	..	1,402.4	..	1,976.8
1910	444.1	..	1,577.8	..	2,021.9
1915	531.8	..	1,422.9	..	1,954.7
1920	546.1	..	1,675.4	..	2,221.5
1925	537.7	..	1,598.5	..	2,136.2
1930	513.7	..	1,673.4	..	2,187.1
1935	595.5	..	2,386.2	..	2,981.7
1940	606.6	617.0	..	2,132.4	3,356.0
1945	671.8	756.9	..	2,217.7	3,646.4
1950	817.7	1,370.4	..	1,845.7	4,033.8
1955	887.8	1,737.3	..	1,462.3	4,087.4
1960	906.1	2,066.9	..	1,213.2	4,186.2
1965	945.6	2,437.5	..	1,005.2	4,388.3
1970	1,011.9	2,731.7	..	753.9	4,497.5
1975	790.5	2,203.1	..	581.7	3,575.3
1980	644.1	1,906.6	..	524.7	3,075.4
1985	531.0	1,776.1	..	459.3	2,766.4
1990	405.1	1,438.5	..	399.8	2,243.4
1995	367.9	1,169.5	..	340.1	1,877.5
2000	297.8	861.3	..	270.8	1,429.9
2003	259.6	720.4	..	256.6	1,236.6

(a) Ischaemic heart disease was not recorded in the ICD until 1950. See page 66 for discussion of estimates for 1940–1950.

Table B38: Circulatory diseases death rates, females aged 65–84, 1907–2003

Year	Cerebrovascular disease	IHD ^(a)	Other circulatory plus IHD	Other circulatory	All circulatory
1907	619.4	..	1,161.8	..	1,781.2
1910	567.9	..	1,468.1	..	2,036.0
1915	579.3	..	1,188.3	..	1,767.6
1920	557.9	..	1,393.7	..	1,951.6
1925	589.3	..	1,309.4	..	1,898.7
1930	586.0	..	1,389.7	..	1,975.7
1935	693.1	..	1,787.1	..	2,480.2
1940	714.5	321.4	..	1,643.5	2,679.4
1945	778.8	400.8	..	1,734.7	2,914.3
1950	937.0	734.0	..	1,441.2	3,112.2
1955	969.0	889.3	..	1,164.0	3,022.3
1960	911.9	1,089.6	..	944.7	2,946.2
1965	946.0	1,337.8	..	786.9	3,070.7
1970	1,023.8	1,560.5	..	621.8	3,206.1
1975	833.5	1,256.9	..	458.2	2,548.6
1980	595.5	1,030.5	..	394.0	2,020.0
1985	507.3	1,008.2	..	351.4	1,866.9
1990	391.8	861.4	..	290.8	1,544.0
1995	342.2	677.9	..	253.1	1,273.2
2000	266.9	497.9	..	212.9	977.7
2003	239.1	415.9	..	196.8	851.8

(a) Ischaemic heart disease was not recorded in the ICD until 1950. See page 66 for discussion of estimates for 1940–1950.

Table B39: Death rates, males aged 85 or over, 1907–2003

Year	Senility	Infectious diseases	Injury and poisoning	Mental health and nervous system ^(a)	Respiratory diseases	Cancer	Circulatory diseases	Other conditions	All causes
1907	13,531.8	746.1	576.5	..	5,053.2	1,051.3	4,714.1	3,561.2	29,234.2
1910	14,159.2	602.5	421.8	..	2,952.3	1,024.3	4,338.1	3,223.5	26,721.7
1915	16,447.9	633.6	684.3	..	3,522.7	1,140.5	5,170.1	3,142.4	30,741.5
1920	15,674.5	918.2	852.6	..	3,825.7	1,355.4	5,968.1	2,754.6	31,349.1
1925	13,650.0	375.0	825.0	..	4,375.0	1,800.0	7,450.0	4,375.0	32,850.0
1930	9,020.4	265.3	673.5	..	3,755.1	1,734.7	7,530.6	4,142.8	27,122.4
1935	6,793.7	190.5	460.3	..	3,698.4	1,460.3	10,365.1	2,984.1	25,952.4
1940	5,536.2	304.3	623.2	..	3,434.8	1,681.2	12,188.4	3,956.5	27,724.6
1945	4,814.4	278.4	670.1	..	2,670.1	1,793.8	12,762.9	3,195.9	26,185.6
1950	3,038.8	147.3	674.4	..	2,821.7	1,868.2	13,992.2	2,728.7	25,271.3
1955	1,426.6	160.8	727.3	..	2,720.3	2,118.9	15,524.5	2,181.7	24,860.1
1960	973.9	169.9	719.0	..	2,797.4	2,274.5	15,169.9	2,156.8	24,261.4
1965	390.8	103.4	626.4	..	3,103.4	2,155.2	16,011.5	2,109.3	24,500.0
1970	..	95.2	646.6	225.6	3,042.6	2,416.0	15,990.0	1,684.2	24,446.1
1975	..	87.8	626.9	242.4	2,595.5	2,649.8	14,344.2	1,471.2	22,210.1
1980	..	95.3	505.7	388.5	2,503.0	2,689.9	12,313.6	1,685.8	20,273.4
1985	..	95.7	465.9	734.4	2,771.0	3,073.4	11,781.4	1,999.6	20,980.0
1990	..	105.9	452.3	890.2	2,203.7	3,050.6	9,709.9	1,938.9	18,385.2
1995	..	149.7	392.8	1,067.5	1,865.5	3,179.6	9,168.7	1,810.8	17,659.3
2000	..	201.2	438.6	910.8	2,074.4	3,000.6	7,852.4	1,897.5	16,394.9
2003	..	204.5	433.5	1,032.5	2,012.8	2,893.0	7,453.2	1,890.7	15,936.9

(a) Work has not been done to enable valid comparisons across ICD versions before 1970.

Table B40: Death rates, females aged 85 or over, 1907–2003

Year	Senility	Infectious diseases	Injury and poisoning	Mental health and nervous system ^(a)	Respiratory diseases	Cancer	Circulatory diseases	Other conditions	All causes
1907	14,545.1	742.4	776.2	..	3,509.7	742.4	3,914.7	1,687.4	25,917.9
1910	12,464.2	554.6	554.6	..	3,240.1	904.9	3,999.1	1,517.9	23,235.4
1915	15,271.4	828.0	621.0	..	2,483.9	920.0	3,541.9	1,908.8	25,575.0
1920	13,771.7	772.4	678.2	..	3,202.7	1,243.4	5,406.9	2,091.2	27,166.5
1925	11,519.2	500.0	615.4	..	3,653.8	1,230.8	7,019.2	2,384.7	26,923.1
1930	9,161.3	274.2	629.0	..	3,096.8	1,387.1	7,500.0	2,387.1	24,435.5
1935	6,325.6	232.6	802.3	..	3,779.1	1,430.2	8,941.9	2,209.2	23,720.9
1940	4,802.1	114.6	864.6	..	2,833.3	1,333.3	11,718.8	2,260.4	23,927.1
1945	4,274.6	169.0	866.2	..	2,288.7	1,323.9	11,415.5	1,887.5	22,225.4
1950	3,000.0	94.2	931.9	..	2,240.8	1,560.2	13,717.3	1,345.7	22,890.1
1955	1,373.4	85.8	1,008.6	..	1,806.9	1,433.5	14,094.4	1,360.5	21,163.1
1960	1,033.7	97.4	872.7	..	1,794.0	1,513.1	14,674.2	1,168.5	21,153.6
1965	411.4	62.9	760.0	..	1,631.4	1,402.9	14,397.1	1,288.6	19,954.3
1970	..	74.0	726.0	198.8	1,419.7	1,433.6	14,527.5	1,172.3	19,859.4
1975	..	67.9	525.6	230.6	1,088.7	1,399.7	12,983.6	1,036.8	17,488.4
1980	..	57.4	424.3	340.3	1,065.8	1,410.9	11,573.6	1,336.9	16,312.8
1985	..	68.7	359.2	633.9	1,198.0	1,493.1	11,123.6	1,655.2	16,590.3
1990	..	99.4	317.2	854.1	1,002.8	1,480.0	9,559.8	1,586.0	14,958.0
1995	..	124.3	269.5	1,165.8	916.4	1,485.9	8,729.9	1,530.4	14,259.4
2000	..	155.9	332.3	1,138.4	1,196.1	1,538.1	7,512.9	1,596.9	13,510.6
2003	..	162.4	325.2	1,212.3	1,292.2	1,534.9	6,954.6	1,725.3	13,259.3

(a) Work has not been done to enable valid comparisons across ICD versions before 1970.

Table B41: Cancer death rates, males aged 85 or over, 1922–2003

Year	Colorectal cancer	Stomach cancer	Prostate cancer	Lung cancer ^(a)	Other cancers and lung cancer	Other cancers	All cancers
1922	170.2	170.2	85.1	(a)	702.2	..	1,127.7
1925	150.0	325.0	175.0	(a)	1,150.0	..	1,800.0
1930	285.7	428.6	102.0	(a)	918.4	..	1,734.7
1935	285.7	285.7	206.3	(a)	682.6	..	1,460.3
1940	362.3	289.9	202.9	(a)	826.1	..	1,681.2
1945	309.3	329.9	299.0	61.9	..	793.7	1,793.8
1950	379.8	294.6	457.4	46.5	..	689.9	1,868.2
1955	321.7	433.6	531.5	62.9	..	769.2	2,118.9
1960	405.2	333.3	601.3	137.3	..	797.4	2,274.5
1965	327.6	321.8	683.9	178.2	..	643.7	2,155.2
1970	310.8	325.8	706.8	230.6	..	842.0	2,416.0
1975	463.9	280.0	685.4	255.0	..	965.5	2,649.8
1980	377.5	241.9	623.0	432.4	..	1,048.1	2,722.9
1985	391.9	222.2	749.8	469.0	..	1,265.2	3,098.1
1990	348.8	163.6	851.7	466.7	..	1,219.8	3,050.6
1995	380.5	158.5	903.7	477.4	..	1,259.5	3,179.6
2000	321.2	112.2	816.6	450.2	..	1,300.4	3,000.6
2003	299.0	91.1	813.6	389.0	..	1,300.3	2,893.0

(a) Lung cancer not separately identified before 1945.

Table B42: Cancer death rates, females aged 85 or over, 1922–2003

Year	Colorectal cancer	Cancers of the cervix and uterus	Breast cancer	Stomach cancer	Lung cancer ^(a)	Other cancers and lung cancer	Other cancers	All cancers
1922	196.4	53.6	250.0	267.9	(a)	517.8	..	1,285.7
1925	192.3	76.9	173.1	192.3	(a)	596.2	..	1,230.8
1930	274.2	64.5	145.2	290.3	(a)	612.9	..	1,387.1
1935	279.1	69.8	186.0	325.6	(a)	569.7	..	1,430.2
1940	291.7	52.1	239.6	229.2	(a)	520.7	..	1,333.3
1945	288.7	56.3	225.4	239.4	14.1	..	500.0	1,323.9
1950	403.1	68.1	225.1	324.6	5.2	..	534.1	1,560.2
1955	334.8	77.3	244.6	296.1	8.6	..	472.1	1,433.5
1960	393.3	78.7	262.2	280.9	11.2	..	486.8	1,513.1
1965	351.4	80.0	205.7	211.4	45.7	..	508.7	1,402.9
1970	374.6	99.4	187.3	205.8	27.7	..	538.8	1,433.6
1975	352.2	67.9	221.7	144.8	41.1	..	572.0	1,399.7
1980	334.7	65.8	214.3	126.0	44.8	..	616.3	1,401.9
1985	349.1	79.9	221.8	114.9	63.1	..	664.3	1,493.1
1990	303.0	53.0	211.2	89.0	83.3	..	740.5	1,480.0
1995	279.2	46.9	212.9	64.0	107.9	..	775.0	1,485.9
2000	278.0	40.0	191.3	58.2	135.9	..	834.7	1,538.1
2003	231.6	38.7	194.9	56.0	140.5	..	873.2	1,534.9

(a) Lung cancer not separately identified before 1945.

Table B43: Circulatory diseases death rates, males aged 85 or over, 1907–2003

Year	Cerebrovascular disease	IHD ^(a)	Other circulatory plus IHD	Other circulatory	All circulatory
1907	1,560.1	..	3,154.0	..	4,714.1
1910	1,235.2	..	3,102.9	..	4,338.1
1915	1,292.5	..	3,877.6	..	5,170.1
1920	1,530.3	..	4,437.8	..	5,968.1
1925	2,025.0	..	5,425.0	..	7,450.0
1930	1,387.8	..	6,142.8	..	7,530.6
1935	1,634.9	..	8,730.2	..	10,365.1
1940	2,043.5	1,202.9	..	8,942.0	12,188.4
1945	1,969.1	1,185.6	..	9,608.2	12,762.9
1950	2,294.6	2,550.4	..	9,147.2	13,992.2
1955	3,419.6	3,664.3	..	8,440.6	15,524.5
1960	3,228.8	4,673.2	..	7,267.9	15,169.9
1965	3,925.3	5,873.6	..	6,212.6	16,011.5
1970	3,969.9	7,213.0	..	4,807.1	15,990.0
1975	3,740.7	6,424.0	..	4,179.5	14,344.2
1980	3,045.4	5,519.1	..	3,749.1	12,313.6
1985	2,823.5	5,804.3	..	3,153.6	11,781.4
1990	2,290.3	5,049.8	..	2,369.8	9,709.9
1995	2,291.7	4,801.9	..	2,075.1	9,168.7
2000	2,012.4	4,012.0	..	1,828.0	7,852.4
2003	1,831.6	3,914.4	..	1,707.2	7,453.2

(a) Ischaemic heart disease was not recorded in the ICD until 1950. See page 66 for discussion of estimates for 1940–1950.

Table B44: Circulatory diseases death rates, females aged 85 or over, 1907–2003

Year	Cerebrovascular disease	IHD ^(a)	Other circulatory plus IHD	Other circulatory	All circulatory
1907	1,754.9	..	2,159.8	..	3,914.7
1910	1,109.2	..	2,889.9	..	3,999.1
1915	1,310.9	..	2,231.0	..	3,541.9
1920	1,243.4	..	4,163.5	..	5,406.9
1925	2,000.0	..	5,019.2	..	7,019.2
1930	1,806.5	..	5,693.5	..	7,500.0
1935	2,011.6	..	6,930.3	..	8,941.9
1940	2,354.2	781.3	..	8,583.3	11,718.8
1945	2,028.2	816.9	..	8,570.4	11,415.5
1950	3,104.7	1,733.0	..	8,879.6	13,717.3
1955	3,665.2	2,459.2	..	7,970.0	14,094.4
1960	3,988.8	3,314.6	..	7,370.8	14,674.2
1965	4,388.6	3,840.0	..	6,168.5	14,397.1
1970	4,633.6	5,329.6	..	4,564.3	14,527.5
1975	4,388.6	4,889.2	..	3,705.8	12,983.6
1980	3,796.7	4,204.2	..	3,572.7	11,573.6
1985	3,483.8	4,389.1	..	3,250.7	11,123.6
1990	2,828.4	4,221.2	..	2,510.2	9,559.8
1995	2,595.8	3,961.1	..	2,173.0	8,729.9
2000	2,239.2	3,453.0	..	1,820.7	7,512.9
2003	2,109.0	3,171.2	..	1,674.4	6,954.6

(a) Ischaemic heart disease was not recorded in the ICD until 1950. See page 66 for discussion of estimates for 1940–1950.

Table B45: Septicaemia death rates, males and females aged 65–84 and 85 or over, 1907–2003

Year	Males 65–84	Males 85 or over	Females 65–84	Females 85 or over
1907	14.1	0.0	1.3	0.0
1910	5.2	30.1	7.1	0.0
1915	12.2	0.0	13.3	0.0
1920	11.9	0.0	1.8	0.0
1925	7.2	25.0	4.9	0.0
1930	2.6	0.0	1.7	16.1
1935	2.7	31.7	2.7	11.6
1940	0.8	0.0	1.5	0.0
1945	0.4	20.6	0.0	7.0
1950	1.7	0.0	1.5	0.0
1955	2.1	0.0	1.0	8.6
1960	1.4	0.0	0.9	7.5
1965	1.3	11.5	0.8	5.7
1970	1.9	0.0	1.4	6.9
1975	2.1	0.0	2.0	7.2
1980	6.5	18.3	6.2	28.0
1985	9.1	52.5	9.6	41.7
1990	15.0	67.4	10.1	68.2
1995	15.8	107.5	12.2	90.8
2000	27.2	171.6	21.3	118.2
2003	28.1	170.0	21.5	128.8

Table B46: Tuberculosis death rates, males and females, 1910–2003

Year	Males					Females				
	15–24	25–44	45–64	65–84	85 or over	15–24	25–44	45–64	65–84	85 or over
1910	56.1	122.8	176.1	112.3	90.4	85.5	125.1	88.7	62.4	0.0
1915	55.7	116.4	155.0	112.6	50.7	73.7	105.7	71.2	60.5	46.0
1920	47.2	114.7	152.1	83.1	43.7	64.9	86.9	66.0	39.8	0.0
1925	30.0	97.4	125.6	108.1	25.0	71.7	72.8	54.9	44.3	19.2
1930	25.2	79.3	114.3	102.0	20.4	57.0	66.0	45.9	42.0	48.4
1935	19.0	59.2	105.3	112.0	47.6	34.5	54.1	38.7	33.8	34.9
1940	14.5	41.1	103.3	99.9	29.0	27.0	43.0	29.1	38.2	31.3
1945	11.1	35.4	89.5	115.9	30.9	23.3	37.8	25.6	32.0	21.1
1950	3.4	18.4	66.6	112.0	62.0	8.8	17.5	16.6	19.9	20.9
1955	0.3	4.8	25.7	69.8	42.0	1.2	4.9	5.7	11.7	8.6
1960	0.1	2.6	14.6	47.1	32.7	0.6	2.5	4.0	8.3	7.5
1965	0.0	0.8	7.0	32.0	57.5	0.0	0.9	2.6	3.8	8.6
1970	< 0.1	0.4	4.3	20.4	35.1	0.0	0.3	1.9	3.2	2.3
1975	< 0.1	0.2	1.9	11.1	41.8	0.0	< 0.1	1.4	1.9	0.0
1980	0.0	0.1	1.7	8.3	29.3	0.0	< 0.1	0.6	2.3	2.8
1985	< 0.1	< 0.1	1.3	5.1	0.0	0.0	< 0.1	1.0	1.9	0.0
1990	0.0	0.3	1.0	5.0	0.0	< 0.1	< 0.1	0.7	2.0	0.0
1995	0.0	0.0	0.3	3.2	15.9	0.0	< 0.1	0.2	1.6	5.2
2000	0.0	0.0	0.1	2.8	3.9	0.0	< 0.1	0.2	1.2	1.7
2003	0.0	0.0	0.2	1.4	6.7	0.0	0.0	< 0.1	1.8	1.0

Table B47: Motor vehicle accident death rates, males and females, 1925–2003

Year	0–4	5–14	15–24	25–44	45–64	65–84	85 or over
Males							
1925	3.1	10.5	19.2	13.7	14.4	28.8	75.0
1930	8.8	10.8	33.1	24.9	26.7	37.0	40.8
1935	8.4	14.0	39.0	23.1	27.4	42.5	15.9
1940	11.3	12.7	53.4	32.4	33.5	55.4	29.0
1945	8.5	10.5	16.7	13.6	18.8	30.0	51.5
1950	9.0	12.2	75.0	36.1	33.4	59.3	100.8
1955	12.4	9.3	73.9	40.3	40.3	59.6	69.9
1960	9.3	10.9	70.0	39.9	40.4	72.5	104.6
1965	13.2	10.9	76.1	41.0	46.0	74.2	97.7
1970	16.1	12.0	97.4	41.9	46.4	69.6	85.2
1975	13.9	10.8	94.2	38.7	34.1	48.1	96.1
1980	10.2	11.7	80.5	33.4	26.3	44.1	69.6
1985	9.0	9.0	60.6	26.4	19.1	31.9	43.2
1990	6.0	7.0	42.7	21.6	15.4	27.5	50.5
1995	3.6	4.4	33.1	17.0	11.7	19.8	54.6
2000	2.9	4.1	28.3	16.3	10.8	15.7	34.8
2003	4.0	2.9	24.2	14.4	9.5	13.3	22.2
Females							
1925	2.2	4.8	1.0	2.2	4.5	10.5	38.5
1930	5.9	6.2	3.9	4.5	9.1	16.6	16.1
1935	6.4	4.1	5.7	4.1	7.4	18.0	0.0
1940	7.8	5.5	8.7	4.5	9.9	22.8	41.7
1945	3.8	2.6	4.7	3.6	5.2	10.5	14.1
1950	6.7	5.2	9.8	4.6	6.9	15.5	10.5
1955	9.3	5.7	10.3	4.9	10.1	25.3	42.9
1960	11.3	6.5	17.6	7.3	17.0	31.5	33.7
1965	9.3	5.6	19.1	8.8	17.3	36.1	28.6
1970	9.1	8.5	24.5	11.9	19.1	33.9	41.6
1975	9.9	6.5	20.1	9.5	14.7	27.7	21.5
1980	8.7	6.7	19.7	9.2	12.5	23.2	28.0
1985	6.5	4.8	17.9	8.1	11.1	18.6	22.5
1990	4.2	4.5	14.6	6.1	8.8	16.6	12.3
1995	3.5	3.2	11.3	6.0	6.6	12.3	14.1
2000	3.2	2.0	9.8	4.9	4.4	9.1	9.7
2003	3.2	1.9	7.5	3.3	3.7	7.8	11.7

Table B48: Suicide death rates, males, 1907–2003

Year	15–24	25–44	45–64	65–84	85 or over
1907	6.2	23.6	47.1	54.4	67.8
1910	6.7	26.9	50.8	36.4	90.4
1915	7.9	31.8	47.8	59.1	50.7
1920	7.9	26.2	41.1	52.6	109.3
1925	9.9	25.0	37.6	54.4	125.0
1930	9.5	30.9	56.3	50.2	40.8
1935	7.5	20.6	41.8	35.6	47.6
1940	7.0	16.8	36.6	30.0	43.5
1945	5.3	8.9	21.9	33.5	20.6
1950	7.0	14.7	25.0	41.4	46.5
1955	9.8	16.1	27.8	45.9	28.0
1960	6.8	20.4	28.0	36.4	32.7
1965	10.6	26.8	35.5	40.5	5.7
1970	12.5	23.0	32.7	33.4	35.1
1975	13.6	20.8	23.6	29.9	33.4
1980	17.5	23.0	23.2	24.5	36.6
1985	24.0	24.7	21.6	24.2	49.4
1990	27.0	27.4	22.9	27.0	50.5
1995	25.4	30.6	23.6	21.7	40.5
2000	19.7	31.7	19.9	20.3	46.4
2003	18.0	26.9	19.1	20.0	31.1

Table B49: Suicide death rates, females, 1907–2003

Year	15–24	25–44	45–64	65–84	85 or over^(a)
1907	4.1	5.1	8.2	10.3	0.0
1910	4.1	4.8	10.5	4.7	0.0
1915	4.4	8.1	12.5	2.1	0.0
1920	3.0	8.2	8.5	4.5	0.0
1925	3.0	8.1	7.5	4.9	0.0
1930	5.3	6.2	9.9	4.4	0.0
1935	3.9	9.2	8.5	5.9	0.0
1940	3.1	6.5	9.1	8.9	0.0
1945	1.6	5.2	11.1	5.8	7.0
1950	2.2	5.5	10.1	7.9	0.0
1955	1.2	7.1	12.7	7.2	4.3
1960	2.0	7.6	14.4	10.6	3.7
1965	6.4	15.1	22.4	16.4	5.7
1970	4.7	9.7	15.8	13.1	4.6
1975	3.9	9.7	14.1	9.4	1.8
1980	4.4	8.1	8.5	8.0	5.6
1985	4.9	5.4	8.5	8.6	6.8
1990	4.4	6.5	6.5	8.2	7.6
1995	6.4	7.3	7.5	6.0	3.0
2000	5.8	8.1	5.4	5.6	3.4
2003	3.7	7.2	6.4	4.2	2.5

(a) There were no deaths from suicide recorded for females aged 85 or over until 1945.

Table B50: Respiratory and SIDS deaths, numbers of persons aged under 1 year, 1968–2003

Year	Respiratory deaths	SIDS deaths	Respiratory + SIDS deaths
1968	483	26	509
1969	549	19	568
1970	515	39	554
1971	572	53	625
1972	480	88	568
1973	344	241	585
1974	278	284	562
1975	205	323	528
1976	165	300	465
1977	114	338	452
1978	107	414	521
1979	85	381	466
1980	77	387	464
1981	54	424	478
1982	56	451	507
1983	64	450	514
1984	53	497	550
1985	62	522	584
1986	43	525	568
1987	39	515	554
1988	47	468	515
1989	47	470	517
1990	49	492	541
1991	51	357	408
1992	41	294	335
1993	32	252	284
1994	25	209	234
1995	52	199	251
1996	34	206	240
1997	21	148	169
1998	28	135	163
1999	24	155	179
2000	21	129	150
2001	23	101	124
2002	21	119	140
2003	32	73	105

Table B51: Infant mortality rates per 1,000 live births, 1912–2003

Year	Males	Females
1912	80.06	63.04
1915	74.25	60.47
1920	76.66	61.15
1925	58.80	47.73
1930	51.90	42.31
1935	44.59	34.76
1940	42.82	33.83
1945	32.56	26.02
1950	27.20	21.60
1955	24.66	19.23
1960	22.39	17.80
1965	20.59	16.24
1970	21.92	16.08
1975	16.29	12.13
1980	11.93	9.44
1985	11.02	8.74
1990	9.05	7.23
1995	6.14	5.15
2000	5.66	4.62
2003	5.2	4.3

Sources: CBCS Bulletins 1907–1963; ABS Deaths 3302.0, 2003.

Table B52: HIV/AIDS death rates, males, 1988–2003

Year	25–44	45–64	65–84
1988	3.7	1.8	0.6
1989	7.0	3.4	0.7
1990	9.0	5.9	0.0
1991	9.7	6.2	0.1
1992	11.1	7.4	1.1
1993	13.9	8.4	1.0
1994	11.4	8.4	0.7
1995	10.9	7.6	1.3
1996	12.1	7.0	0.9
1997	4.7	3.5	0.5
1998	3.0	2.2	0.9
1999	2.3	1.8	0.2
2000	2.0	1.6	0.3
2001	2.0	1.7	0.8
2002	1.6	2.1	0.3
2003	1.7	1.6	1.2

Table B53: Age-standardised death rates for miscellaneous conditions, 1907–2003

Year	Meningococcal		Viral hepatitis		Diabetes	
	Males	Females	Males	Females	Males	Females
1907	15.7	21.0
1910	13.3	23.1
1915	12.4	5.7	17.1	21.2
1920	1.8	0.6	17.7	22.5
1925	1.1	0.8	15.5	25.9
1930	0.8	0.4	14.7	23.6
1935	0.6	0.5	18.3	33.8
1940	0.6	0.3	22.9	34.4
1945	1.0	0.7	18.4	32.7
1950	0.9	0.7	0.7	0.6	15.6	24.4
1955	0.6	0.6	1.0	1.1	14.3	22.1
1960	0.3	0.3	0.6	0.7	16.0	19.3
1965	0.1	0.2	0.4	0.5	17.6	20.8
1970	< 0.1	< 0.1	0.5	0.5	23.0	23.4
1975	0.1	< 0.1	0.2	0.2	20.9	17.5
1980	0.1	< 0.1	0.2	0.2	16.3	15.1
1985	< 0.1	< 0.1	0.3	0.2	17.7	14.9
1990	0.2	0.2	0.4	0.2	18.6	13.6
1995	0.1	< 0.1	1.1	0.4	19.9	14.5
2000	0.2	0.1	1.1	0.6	20.3	12.9
2003	0.1	0.1	1.2	0.5	20.9	12.9