

## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on [13 11 14](tel:131114), or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)

## Suicide & self-harm monitoring data

### Over 3,000 deaths by suicide occur each year in Australia

In 2022, there were 3,249 deaths by suicide - an average of about 9 deaths per day - with a rate of 12.3 per 100,000 population.

### Suicide and self-inflicted injuries was the second leading cause of premature death from injury or disease

In 2018, males lost an average of 42.2 years and females lost an average of 41.6 years to 'suicide and self-inflicted injuries'.

### Males are 3 times more likely to take their own life than females

In 2022, there were 2,455 male deaths at a rate of 18.8 per 100,000; there were 794 female deaths at a rate of 5.9 per 100,000. In 2022, the number of deaths by suicide was higher for males than females in all reported age groups.

### Females are more likely to attempt suicide or be hospitalised for intentional self-harm than males

Rates of ambulance attendances for suicide attempt and self-injury were higher for females than males. In 2021-22 females made up two-thirds (67%) of intentional self-harm hospitalisations.

### Suicide is the leading cause of death for young people

Almost one third of all deaths among Australians aged 15-24 years were due to suicide in 2022.

### The rate of suicide among young First Nations people is three times that of young non-Indigenous Australians

In 2018-2022, the age-specific rate of suicide deaths was 3.1 times higher in First Nations people aged 0-24.

### The highest proportion of deaths by suicide occur during mid-life

More than half of all deaths by suicide in 2022 (55%) occurred in people aged 30-59 (1,774 deaths).

### Suicide rates are highest among middle aged and older males

Since 2008, the highest suicide rates have generally been among males aged 40-49 and over 85.

### Results of a birth cohort analysis show trends in suicides have changed over time

Suicide rates for the most recently born female cohorts are higher than those for earlier female cohorts at the same age while suicide rates for the most recently-born male cohorts are similar to, or lower than, earlier male cohorts at the same age.

### Using linked data, the estimated suicide risk is higher among those with fewer years of education

Among males aged 25-54 with secondary school or no education, the cumulative suicide risk is 2.6 times higher than among males with a university degree. This gradient between highest and lowest levels of educational attainment for females was consistent with that seen for males - with a smaller ratio (1.6 times).

## Suicide and Self-harm monitoring data

*Suicide and Self-harm Monitoring* brings together key statistical data on suicide and self-harm from multiple national sources that will be updated regularly as new data become available. Here, you can examine the data through interactive visualisations and read information on the demographics, trends, methods and risk factors of suicide and self-harm in Australia.

This website represents only one part of a comprehensive program of work on suicide and self-harm in Australia by the AIHW (for more information visit [About the Suicide and Self-harm Monitoring System](#)).

## Why is it important to collect data about suicide and self-harm?

Monitoring of suicide and intentional self-harm—how many people harm themselves, when, where and how—can provide a better understanding of the nature of suicide and self-harm in Australia and help determine who may be at increased risk. Reporting of this data can raise community awareness of suicide and self-harm, further research, improve responses and support services for those that need them, and inform the design and targeting of suicide prevention activities.

## Considerations when using these data

There are several considerations to keep in mind when examining suicide and self-harm data and information in Australia.

### Deaths by suicide

The assembling and national reporting of deaths by suicide has up to an 18-month time lag. Deaths by suicide may be presented by year of occurrence of death or year of registration. Although reporting of deaths by suicide by year of death can provide more reliable information on trends in occurrence than reporting by year of registration, the latest data available may underestimate the number of deaths, especially those in the later months of the year, due to a lag in registration. For this reason, and unless otherwise specified, year of registration of death has been used to allow the latest year of data to be compared with previous years. In both cases, the latest years of data are coded with preliminary causes of death information and may underestimate causes of death that are usually certified by a coroner, including deaths by suicide. For more information on how deaths are registered, coded and updated, visit [Technical notes](#).

### Suspected deaths by suicide

State and Territory suicide registers can provide more timely data on suspected deaths by suicide. Recent surveillance data from suicide registers are preliminary and may change over time, typically upon completion of the coronial investigation. Suicide registers are operational in all Australian jurisdictions, except for Western Australia. The AIHW contributed to the development of suicide registers in South Australia (established in 2022), the Australian Capital Territory (established in 2021) and the Northern Territory (established in 2023).

Suicide registers in New South Wales, Victoria and Queensland publish monthly data reports on suspected or confirmed suicide deaths in their respective jurisdictions. The AIHW receives data on suspected or confirmed suicide deaths from several other jurisdictions' registers on a fortnightly or monthly basis. The data can be used to inform governments' decision making and responses to emerging issues that may influence suicide risk in the community, such as cost of living concerns. Due to the highly sensitive nature of coronial investigations, the AIHW will not publicly release jurisdictional data unless they have been published by the relevant data custodians (visit [Data from suicide registers](#)).

### Hospital admissions

Hospital admissions data are collated as an annual release with a 12-month lag.

### Ambulance attendances

Ambulance data are currently available for some states and territories for selected months from 2018 to 2021 (visit [Ambulance attendances: suicidal and self-harm behaviours](#)), with monthly data from January 2021. In addition, monthly ambulance data for Victoria from January to March 2021 are also reported (visit [COVID-19](#)). Further information on the collection of data and sources is available in the [Technical notes](#).

### Issues with small numbers and the need for caution

Deaths by suicide are statistically rare events. Small numbers can raise privacy and confidentiality issues but also statistical concerns. For this report, values based on small numbers of deaths, hospitalisations for intentional self-harm or ambulance attendances have been suppressed in order to maintain data confidentiality, and/or avoid publishing statistics of low reliability. Visit [Technical notes](#) for further information.

The statistics on deaths by suicide reported here fluctuate from one period to the next—mostly due to small counts (and in the case of females, very small counts)—especially in many smaller subgroups (for example, individual age groups or small geographic areas). Estimates of rates are also subject to random variability. Statistics based on small numbers of deaths by suicide should be interpreted with caution and all rates and their comparison with rates in other populations should be reported in context. For further insight into the methodological challenges and statistical issues of monitoring suicide and self-harm, visit [Suicide Mortality in Australia: Estimating and Projecting Monthly Variation and Trends From 2007 to 2018 and Beyond](#).

## How to use the interactive data visualisations

- Due to large data sets, visualisations may take time to load.
- Visualisations are compatible with Chrome, Microsoft Edge and Firefox.
- Each panel may contain more than 1 visualisation. You can interact with the visualisations to see the specific data you are interested in by either selecting from the filter(s) at the bottom of the chart, or in the case of maps, from the pop up box by clicking on an area of interest.
- Hover over each data point to see the underlying data and, if available, further details.

- The [Data downloads](#) page provides the source data as Excel (.xlsx) files. The relevant source supplementary table is cited at the bottom of each visualisation.
  - Each visualisation may be downloaded and exported or shared.
  - A print friendly PDF of all pages of text and the default visualisations related to suicide and self-harm may also be downloaded—click on the ‘Download all data pages’ button. Visit [Technical notes](#) for information about data sources, data quality and methodology.
- 



## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on [13 11 14](tel:131114), or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)

## Suicide & self-harm monitoring: Deaths by suicide in Australia

If at any point you feel worried about harming yourself while viewing the information on this website - or if you think someone else may be in danger - please stop reading and [seek help](#).

### Important points to remember about deaths by suicide:

**Each statistic represents a person - with a family and community grieving for their loss**

Although it is a relatively rare cause of death - in 2022, 1.7% of all deaths registered were by suicide - it can have devastating and long-lasting effects on those left behind.

### The reasons people take their own life are complex

Suicide can affect anyone - regardless of their personal characteristics and family background - but some populations are at greater risk. There is also no single reason why a person chooses to end their life - the reasons are often complex. For information on risk factors see [Behaviours and risk factors](#).

### Deaths by suicide are preventable

Monitoring the number, trends and rates of suicide in Australia is key to understanding who is at risk and for the planning and targeting of suicide prevention activities.

It is our endeavour that by bringing together various data sources we can strengthen the evidence base to build a more coherent picture of suicide and self-harm in Australia in order to improve the effectiveness of suicide prevention.



## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on [13 11 14](tel:131114), or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)



## Deaths by suicide over time

Numbers and rates of deaths by suicide change over time as social, economic and environmental factors influence suicide risk. The data visualisations below provide an overview of the characteristics of people who have died by suicide in Australia since 1907, looking at trends and variations by sex and age - how many there were, how old they were when they died, and the methods used over time. This analysis may provide useful information on potentially preventable factors, such as restricting access to means of suicide and reducing the risks posed by social or economic factors. Over time, the accuracy and quality of the data collected have been influenced by a number of factors including changes in legislation, technology and a reduction in social stigma.

Suicide deaths by sex, Australia, 1907 to 2022.

The line graph shows age-standardised rates of suicide for males, females and persons from 1907 to 2022. Users can also choose to view the number of deaths by suicide and male to female rate ratios from 1907 to 2022 and median age at death by sex from 1964 to 2022. The data can be viewed for any period between the years for which data are available.

## Suicide deaths by sex, Australia, 1907 to 2022



Source: ABS Causes of Death, Australia 2023  
Supplementary Table: NMD S1  
Latest data: 2022 (annual release)

See notes ►

Numbers of deaths by suicide increased steadily over the first half of the 20th Century (from 461 in 1907 to 760 in 1950), with peaks and troughs in numbers of suicides corresponding with significant world events (see below). However, since the 1950s numbers of deaths by suicide increased more steeply over time - in part driven by population growth. Peaks in numbers of deaths by suicide occurred during the 1960s and late 1990s. Since the mid-2000s numbers of deaths by suicide in Australia have increased, reaching over 3,000 Australians dying by suicide by 2015.

### Have suicide rates changed over time?

Between 1907 and 2022, age-standardised suicide rates in Australia ranged from 8.4 deaths per 100,000 population per year (in 1943 and 1944) to 18.4 in 1963.

- Suicide rates peaked in 1913 (18.0 deaths per 100,000 population), 1915 (18.2), 1930 (17.8), 1963 (18.4) and 1967 (17.7). These peaks tended to coincide with major social and economic events or changes, see [Impact of social and economic events](#)
- Suicide rates tended to increase from 1907 to 1915 (from 16.9 to 18.2 deaths per 100,000 population). Rates then fluctuated throughout the late 1910s and early 1920s (from 13.1 deaths per 100,000 population in 1918 to 16.2 in 1920, returning to 12.8 in 1922), before increasing to a peak of 17.8 in 1930.
- Rates then declined throughout the 1930s and early 1940s, reaching a low of 8.4 deaths per 100,000 population in 1943 and 1944 (however, suicide rates for the war years may have been underestimated, see [Impact of social and economic events](#)).
- Rates tended to increase throughout the 1950s, peaking at 18.4 deaths per 100,000 population in 1963. Rates remained high throughout the 1960s while the 1970s and early 1980s saw a decline in rates (from 15.4 deaths per 100,000 population in 1971 to 11.6 in 1984).
- Rates began to rise in 1985 and fluctuated from 14.3 in 1987 to 11.9 in 1993 with a recent peak of 14.8 in 1997. This was followed by sustained declines over the early 2000s, with a low of 10.2 per 100,000 population in 2006.
- After 2006, suicide rates began to rise, partly due to improvements in data quality and capture (see below). In 2022, the rate was 12.3 deaths per 100,000 population - down from a post-2006 high of 13.2 in 2017 and 2019. It is important to note that deaths registered in 2021 and 2022 are preliminary and as such, are subject to revision (see below).

It is important to note that deaths by suicide were underestimated in the collection of routine deaths data, particularly in the years before 2006 (AIHW: Harrison et al 2009; De Leo, 2010; AIHW: Harrison & Henley 2015). Since then, the Australian Bureau of Statistics (ABS) has introduced a revisions process to improve data quality by enabling the revision of cause of death for open coroner's cases over time. Deaths registered in 2021 and 2022 are preliminary and data for 2020 are revised and therefore, data for these years are subject to further revision by the Australian Bureau of Statistics. Data from 2006 to 2019 are final (for further information see [Technical notes](#)).

### What's changed in the last decade?

Please note: small numbers can result in large yearly variation in suicide rates. Caution is advised when making year to year comparisons.

- Over the last decade, the age-standardised suicide rate for males increased from 16.2 deaths per 100,000 population in 2011 to 18.8 in 2022. Female rates also increased from 5.1 deaths per 100,000 population in 2011 to 5.9 in 2022.

For detailed analysis of recent trends in suicide in Australia, see [Suicide Mortality in Australia: Estimating and Projecting Monthly Variation and Trends From 2007 to 2018 and Beyond](#).

### Impact of social and economic events

While the reasons for an individual's suicide death are personal and often complex, overall peaks and troughs in rates and numbers of deaths by suicide historically coincide - more or less - with social and economic events.

Falls in the male suicide rate coincided with both World Wars 1 and 2. These falls are at least partly a statistical artefact due to the fact that deaths from all causes (including deaths by suicide) of Australian service personnel while overseas were not included in Australian death registration data, while population estimates were not adjusted to allow for the absence of these personnel (AIHW 2005; AIHW: Harrison & Henley 2014).

The highest annual age-standardised rate for males in the last century occurred in 1930 (29.8 deaths per 100,000 population), during the Great Depression - a period of high unemployment, particularly among males. The rise in both male and female suicide rates in the 1960s has been attributed, in part, to the unrestricted availability of barbiturate sedatives (Oliver & Hetzel 1972; Whitlock 1975). Subsequent falls in these rates in the late 1960s and early 1970s have in turn been attributed to the introduction of restrictions to the availability of these drugs in July 1967 (AIHW: Harrison & Henley 2014). While high rates of suicide in the late 1980s and early 1990s also coincided with a period of economic uncertainty in Australia, the social and economic disruption related to the [COVID-19](#) pandemic has not seen an increase in the number of suspected deaths by suicide referred to coroners courts.

### Males have consistently higher rates of suicide than females

Since 1907, the male age-standardised suicide rate has been consistently higher and more variable than the female rate. Variations in the overall suicide rate in Australia have been largely driven by changes in the male suicide rate.

The peak in overall suicide rates in 1930 was driven by an increase in male suicide rates, peaking at 29.8 deaths per 100,000 in 1930 - the highest rate ever recorded. Similarly, the increase in overall suicide rates in the 1990s was also mainly driven by an increase in male rates. The peak in the 1960s reflects a rise in suicide rates for both males and females.

The male suicide rate ranged from a high of 5.6 times that of females in 1930 to lows of less than twice the female rate in the 1960s and early 1970s - mainly due to the marked rise in female suicide rates at this time. Since then, the male suicide rate has fluctuated around 3 to 4 times that of the female rate.

Although males are more likely to die by suicide, females are hospitalised for intentional self-harm (with and without suicidal intent) almost twice as frequently as males (see [Intentional self-harm hospitalisations](#)). Furthermore, ambulance attendance data reporting on attendances for suicide attempts between 2018 and 2022 suggest females are more likely to attempt suicide than males (see [Ambulance attendances, suicidal and self-harm behaviours](#)).

### Patterns of suicide by age have changed over time

Age-specific suicide rates for males are higher than those for females across all reported age groups for all years. Use the year slider to see how patterns of suicide in males and females have changed in Australia over time. Hover over the graph to display the tooltip to see the trend in deaths by suicide by sex over time for each age group. The age distribution of deaths by suicide is similar for males and females. The highest proportion of deaths by suicide occur during mid-life. More than half of all deaths by suicide (54.6%) in 2022 occurred in people aged 30-59 (1,774 deaths), compared with 20.5% for those aged 15-29, and 24.5% for those aged 60 and over.


In 2022, the highest suicide rates for males occurred in those aged 85 and over and those aged 45-49 (32.7 and 32.6 deaths per 100,000 population, respectively). For males aged 85 and over, the number of deaths by suicide was the lowest (71 deaths), whereas males aged 45-49 had the highest number of suicide deaths recorded (262). High rates of suicide were also recorded in males aged 50-54 and 80-84 (27.4 and 22.2, respectively). Males aged 40-54 accounted for one quarter (29.1%) of deaths by suicide among males. The highest suicide rate for females was in those aged 85 and over (10.6). Females aged 40-54 accounted for 25.8% of deaths by suicide among females.

Suicide deaths by age and sex, Australia, 2022.

The bar chart shows the age-specific rates of suicide for males and females by age groups (five year age bands from 15-19, 20-24, etc to 80-84 and 85 and over). Users can choose to view numbers of deaths by suicide for males and females in these age groups. Data can also be viewed by year from 1907.

## Suicide deaths by age and sex, Australia, 2022

Age-specific rate (per 100,000)



Note: Age-specific rates may refer to a small number of deaths.

Source: ABS Causes of Death, Australia 2023

Supplementary Table: NMD S2

Latest data: 2022 (annual release)

[See notes ►](#)

For approximately the first half of the period 1907 to 2022, age-specific suicide rates in males generally increased with age; however, by the start of the 1990s this pattern had changed substantially with suicide rates highest in younger males aged 20-39 and males aged 80 and over. Since 2008, the highest suicide rates have been observed in middle-aged males (aged 40-54) and older males aged 85 and over; however, it should be noted that rates of death by suicide in males aged 85 and over have historically been based on relatively small numbers compared to other age groups and as such, the rates can be quite volatile over time and should be interpreted with caution.

Throughout 1907 to 2022, the lowest suicide rates in males were observed in those aged 15-19.

- From 1907 to 1970, suicide rates in males aged 15-19 were less than 10 deaths per 100,000 population. Rates then increased throughout the 1970s and 1980s peaking at 21.0 in 1988, while remaining the lowest of the reported age groups.
- In 2022, the suicide rate for males aged 15-19 was 9.7 deaths per 100,000 population.

Males aged 20-24 had the second-lowest age-specific suicide rates of all males for most of the 20th Century; however, this changed from the late 1960s.

- From 1907 to 1966, suicide rates for males aged 20-24 were around 11 deaths per 100,000 population with peaks of 16.8 in 1914, 17.0 in 1958, and 19.1 in 1963 and a low of 1.9 in 1944.
- From the late 1960s to the late 1990s, suicide rates in this age group increased steadily, reaching a high of 43.1 deaths per 100,000 population in 1997.
- Rates then fell steadily to 16.3 deaths per 100,000 population in 2009, before rising again to 25.4 in 2020. In 2022, the rate for death by suicide for males aged 20-24 years was 20.5.
- A similar pattern was observed for those aged 25-29.

The pattern of age-specific suicide rates for middle-aged males (aged 40-59) was different to that of younger age groups, with the highest rates being observed in the first part of the 20th Century and then falling to lower levels.

- The highest age-specific suicide rate for middle-aged males was 64.9 deaths per 100,000 population in 1913 for males aged 50-54. Peaks of more than 56 were also seen in 1930 (56.6). Age-specific rates then fell to a low of 14.5 in 1944. Similar patterns were seen for 40-44, 45-49 and 55-59 age groups with the second highest age-specific rate of 63.9 deaths per 100,000 population for males aged 55-59 in 1931 and the lowest age-specific rate of 10.5 for males aged 40-44 in 1944.
- Rates tended to increase throughout the 1950s and 1960s peaking again at 42.0 deaths per 100,000 population in 1962 for males aged 55-59, before falling to 19.1 in 1983 for males aged 45-49. The greatest decline during this time period was seen for males aged 55-59 falling from 41.6 deaths per 100,000 population in 1968 to 18.4 in 1977.
- Since then, rates for these age groups have fluctuated to a high of 34.4 deaths per 100,000 population in 1987 for males aged 55-59 and a recent high of 33.8 in 2017 for males aged 45-49.
- A similar pattern was seen in males aged 60 and older.



It should be noted that the number of deaths by suicide recorded for older males historically has been low, particularly for males aged 75 and older. This causes fluctuation in the age-specific rates. Therefore, caution should be used when interpreting trends for these age groups over time.

- The age-specific suicide rate for males aged 60 and older was about 40 deaths per 100,000 population from 1907 to 1967.
- From 1968, suicide rates for males aged 60 and older generally fell. For example, suicide rates for males aged 65-69 fell to an all-time low of 12.6 per 100,000 population in 2005. In 2022 the rate of suicide for males aged 65-69 was 19.9.

Age-specific suicide rates for females showed comparatively little variation over time - except for a peak in multiple age groups during the 1960s.

- For the first half of the 20th Century, age-specific rates in females aged 40-59 was about 9 deaths per 100,000 population, with peaks of 21.5 in 1915 and 21.2 in 1953, in the 55-59 age group. The highest rate recorded for females was 29.2 deaths per 100,000 population in 1963 for the 50-54 age group and remained around 20 until peaking a second time in 1967 at 27.1 for the 65-69 age group. Rates then fell to a low of 4.1 deaths per 100,000 population in 2004 and 2005 for females aged 55-59. Age-specific suicide rates have increased in this age group to 9.7 deaths per 100,000 population in 2019 and fell to 6.1 in 2020 and rose again to 7.1 in 2022.
- Similar patterns were seen for females aged 20-39 and 60 and older, albeit with lower suicide rates.
- A different pattern has been observed in females aged 15-19. Suicide rates fluctuated from around 2 to 6 deaths per 100,000 population from 1907 to the late 1930s. The fluctuations in rates have been mainly due to small numbers of deaths by suicide in this age group. Rates then declined to around 1 to 2 deaths per 100,000 population during the 1940s and 1950s. Rates then increased in the 1960s to the late 1990s, fluctuating between 2 and 6 deaths per 100,000 population. Since then, suicide rates have increased to between 3 and 8 deaths per 100,000 population with the highest rate recorded in this age group in 2012 (8.3 deaths per 100,000 population). In 2022, the rate of death by suicide for females aged 15-19 years was 6.6 per 100,000 population.

### How have methods of suicide changed over time?

Understanding the methods used for suicide can play an important role in suicide prevention. These data are provided to inform discussion around restriction of access to means as a policy intervention for the prevention of suicide.

Please consider your need to read the following information. If this material raises concerns for you or if you need immediate assistance, please contact a [crisis support service](#), available free of charge, 24 hours a day, 7 days a week.

Please consider the [Mindframe guidelines](https://mindframe.org.au/suicide/communicating-about-suicide/mindframe-guidelines) (https://mindframe.org.au/suicide/communicating-about-suicide/mindframe-guidelines) if reporting on these statistics.

The pattern of methods used for suicide has changed greatly, sometimes rapidly, over the last century as new methods have become available or as restrictions to the availability of some methods have been introduced. The methods of suicide used by males and females differed over the period 1907 to 2022; however, as males account for the majority of deaths by suicide the methods used by males have a greater influence on the overall pattern than the methods used by females.

The classification system used to code causes of deaths data, ICD-10, uses the term 'mechanism' to refer to the external cause of death. Throughout *Suicide & self-harm monitoring* 'mechanism' has been used in data visualisations, while the term 'method' has been used in the accompanying text.

Suicide deaths by sex and mechanism, Australia, 1907 to 2022.

The line graph shows age-standardised suicide rates by mechanism for poisons, gas, firearms, hanging and other mechanisms from 1907 to 2022. Users can also choose to view age-standardised rates and numbers of deaths by suicide, by sex and mechanism (including all mechanisms) from 1907 to 2022 and median age at death by sex and mechanism from 1964 to 2022. The data can be viewed for any period between the years for which data are available.

---

**Content warning:**

The data in this visualisation might be distressing to some readers as it contains data on the modality of suicide deaths and attempts. Please consider your need to read the following information. If this material raises concerns for you or if you need immediate assistance, please contact a [crisis support service](#), available free of charge, 24 hours a day, 7 days a week.

Please consider the [Mindframe guidelines](#) if reporting on these statistics.

[Proceed to visualisation](#)

Hanging (ICD-10 X70) has become the most common method of suicide in Australia and use of this method increased substantially over the last 25 years. Age-standardised rates of suicide by hanging remain much higher for males than females, but have increased for both sexes.

- Rates of suicide by hanging were relatively steady from 1930 to the late 1980s. Over this period, rates of suicide by hanging ranged between 2.0 and 5.7 per 100,000 population for males and between 0.5 and 1.2 for females. Prior to 1930, rates of suicide by hanging were more volatile.
- From the late 1980s, rates of suicide by hanging increased as other methods of suicide (firearms and poisoning by gas) declined.
- Hanging became the most common method of suicide for males in 1989 and for females in 1997. Age-standardised suicide rates by hanging in males have more than doubled since then - from 5.7 per 100,000 population in 1989 to 12.6 in 2019, then falling to 11.8 in 2022. In 2022, hanging accounted for almost two-thirds (62.0%) of male deaths by suicide.
- Similarly, the rate of suicide by hanging in females increased from 1.9 deaths per 100,000 population in 1997 to 3.3 in 2022. In 2022, hanging caused around half (52.8%) of all deaths by suicide in females, having increased steadily from 30.3% of deaths by suicide in 1997.

Use of firearms (ICD-10 X72-X75) was the most common method of suicide for males from 1907 to the late-1980s.

- In males over this period, the rate of suicide by use of firearms peaked at 10.2 deaths per 100,000 population in 1914 and fell below 5 deaths per 100,000 population between 1941 and 1946.
- Rates of suicide by use of firearms declined steeply for both males and females from 1987 and continued to decline from 1996, coinciding with the introduction of gun control restrictions and reforms.
- In contrast, female rates of suicide by this method were low (maximum of 0.6 deaths per 100,000 population) between 1907 and 2022.

In the 1920s, poisoning by gas (ICD-10 X67), largely due to carbon monoxide poisoning, became a new method of suicide in Australia with the introduction of the domestic gas supply and the motor vehicle to Australia.

- Rates of poisoning by gas peaked in 1963 in females (2.1 deaths per 100,000 population) and were also high for males (4.8). Rates then declined throughout the 1970s - this has been attributed to the replacement of toxic 'town gas' by less toxic gases in most of Australia at this time (AIHW: Harrison & Henley 2014).
- Rates of poisoning by gas subsequently increased again in the 1980s and 1990s, peaking for males (5.8 deaths per 100,000 population) and for a second time in females at a much lower level (1.2 deaths per 100,000 population) in 1997 as a result of the increasing use of motor vehicle exhaust gas (AIHW: Harrison & Henley 2014).
- A decline in poisoning by gas after 1997 was likely due to the introduction of emission controls that greatly reduced the amount of carbon monoxide permitted in the exhaust gas of new motor vehicles (AIHW: Harrison & Henley 2014).

Exposure to poisonous substances excluding gas (ICD-10 X60-X66, X68-X69) was the most common method of suicide for females from 1907 until 1997.

- For most of the first half of the 20th Century, rates of poisoning by substances (excluding gas) were approximately 2 deaths per 100,000 population in females; however, during the 1960s rates increased to 4 times that - peaking at 8.4 in 1967 - before returning to previous levels in the 1980s.

- A similar peak in suicide rates by this method was seen in males, with rates more than doubling in the 1960s to a peak of 8.2 deaths per 100,000 population in 1963 before falling again in the 1970s and 1980s.
- These peaks in suicide rates due to poisonous substances (excluding gas) during the 1960s have been attributed mainly to the unrestricted availability of barbiturate sedatives (AIHW: Harrison & Henley 2014). These trends were not associated with compensatory falls in the use of other methods of suicide during this time. In July 1967, in response to concerns over misuse of these drugs, the supply of barbiturates was limited and deaths by suicide from poisoning (excluding gas) in both males and females declined soon after (AIHW: Harrison & Henley 2014).
- In 2022, poisoning by substances (excluding gas) was the second most common means of suicide among females with a rate of 1.5 deaths per 100,000 population - accounting for almost a third of female deaths by suicide each year for the last decade.

Age-standardised rates for suicides by other methods (ICD-10 X71, X76-X84, Y87.0) are only available from 1964.

- Rates for these methods were relatively stable over the period from 1964 to 2022 for both males and females.
- It is not possible to report on these different methods individually, as the numbers are too small to report for privacy or data reliability reasons.

## References

---

AIHW (Australian Institute of Health and Welfare) 2005. Mortality over the twentieth century in Australia: trends and patterns in major causes of death. Mortality surveillance series no. 4. Cat. no. PHE 73. Canberra: AIHW.

AIHW: Harrison JE, Pointer S and Elnour AA 2009. A review of suicide statistics in Australia. Cat. no. INJCAT 121. Canberra: AIHW.

AIHW: Harrison JE & Henley G 2014. Suicide and hospitalised self-harm in Australia: trends and analysis. Injury research and statistics series no. 93. Cat. no. INJCAT 169. Canberra: AIHW.

AIHW: Harrison JE & Henley G 2015. Injury deaths data, Australia: technical report on issues associated with reporting for reference years 1999-2010. Injury research and statistics series no. 94. Cat. no. INJCAT 170. Canberra: AIHW.

Oliver R & Hetzel R 1973. An analysis of recent trends in suicide rates in Australia (<https://academic.oup.com/ije/article-abstract/2/1/91/648690?redirectedFrom=fulltext>). International Journal of Epidemiology 2(1):91-101.

Whitlock F 1975. Suicide in Brisbane, 1956 to 1973: the drug-death epidemic (<https://pubmed.ncbi.nlm.nih.gov/239307/>). Medical Journal of Australia 1(24):737-43.

---



## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on [13 11 14](tel:131114), or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)

## Birth cohort analysis of deaths by suicide

Analysing deaths by suicide according to the period in which people were born can provide additional insights to that obtained by examining suicide rates by period of death (see [Deaths by suicide over time](#)).

A 'birth cohort' is a group of people born within the same defined period. People in a birth cohort age together over time and experience the same events and changes in technology or cultural norms at the same age.

This birth cohort analysis relates deaths by suicide to period of birth (birth cohort) and age at death. It examines how suicide rates change within birth cohorts as they age and how they vary between birth cohorts when compared at the same age.

### Data sources and methods

This analysis is based on data from the AIHW National Mortality Database, which holds records for deaths in Australia from 1964.

Suicide rates by age at death (5-year age groups; ages 10-14 years and older) were calculated for each birth cohort. Birth cohorts can be defined in terms of any range of birth dates for which data are available; the cohorts presented here are those born in each 5-year period from 1954-58 through to 2004-08. The earliest birth cohort, those born in 1954-58, can be followed for over 60 years. For more information on data sources and methods, see [Suicide in Australia: Trends and analysis 1964 to 2018](#).


### How do suicide rates change among birth cohorts?

The interactive data visualisation shows how suicide rates have changed as people in each birth cohort have aged—with each line representing a birth cohort. By comparing the earlier birth cohorts with those born more recently, see how the age groups most at risk change.

Trends in suicide rates for birth cohorts from 1954-1958 to 2004-2008, by age at death, sex and mechanism, Australia.

The line graph shows age-specific rates of suicide for 5-year birth cohorts from 1954-1958 to 2004-2008 by age at death from 15-19 to 60-64 for males by all mechanisms. Users can also choose to view suicide rates by sex, mechanism and age at death. The highest suicide rate was in males born 1969-1973 who died aged 25-29, followed by males in this cohort who died aged 20-24.

## Trends in suicide rates for birth cohorts from 1954–58 to 2004–08, by age at death, sex and mechanism, Australia



Source: AIHW National Mortality Database  
Supplementary table: S2.6, S3.3, S4.3, S5.3, S6.3  
Latest data: 2018

[See notes ►](#)

In the earlier male birth cohorts (born 1954-58 to 1974-78) peaks in suicide rates for each subsequent birth cohort tended to be higher and occur at successively younger ages of death—with peaks tending to coincide with deaths occurring in the 1990s (period of death). For more information, see [Suicide deaths over time](#). Suicide rates in these cohorts then tended to decline as they aged.

For example, peak suicide rates in males born in:

- 1954-58 occurred at age 40-44 (29.4 deaths per 100,000 cohort members)
- 1959-63 occurred at age 35-39 (31.0)
- 1964-68 occurred at age 30-34 (34.0)
- 1969-73 occurred at age 25-29 (36.9)
- 1974-78 occurred at age 20-24 (33.4).

For the majority of the male cohorts born in the later years, from 1974-78 onwards, suicide rates were still rising at the end of the available data; the oldest people in these cohorts were aged 42-46 years in 2018.

Suicide rates in female cohorts were much lower than those of male cohorts and for the earlier born cohorts tended to increase as they aged.

- For example, the highest suicide rates in female cohorts were in those born in 1964-68 and 1969-73—the same cohorts that had the highest rates in males. However, peaks in suicide rates for these female cohorts tended to occur at older ages (9.7 and 9.5 at age 50-54 and 45-49, respectively) than in male cohorts (which peaked in early adulthood and then declined).


### How do suicide rates vary between birth cohorts when compared at the same age?

The interactive data visualisation shows how suicide rates have changed for people of the same age, but born at different times—each line representing the same age group. By following the suicide rate of a specific age group, see how suicide rates have changed for people born between 1954 and 2008.

Trends in suicide rates at age of death across birth cohorts from 1954-1958 to 2004-2008, by sex and mechanism, Australia.

The line graph shows age-specific suicide rates for ages of death from 10-14 to 60-64, by 5-year birth cohorts from 1954-1958 to 2004-2008 by all mechanisms for females. Users can also choose to view suicide rates by sex, mechanism and for selected age-ranges at death. The rates of suicide among young females aged 15-19 at death showed the greatest change between the earliest and latest born cohorts for which data are available, almost doubling from 3.6 per 100,000 population in the 1954-1958 cohort to 6.4 in the 1999-2003 cohort, with some fluctuation in between these cohorts.

## Trends in suicide rates at age of death across birth cohorts from 1954–58 to 2004–08, by sex and mechanism, Australia



Source: AIHW National Mortality Database  
Supplementary table: S2.6, S3.3, S4.3, S5.3, S6.3  
Latest data: 2018

See notes ►

In females, the suicide rate at age 15-19 for those born most recently (1999-2003) was 1.8 times higher than the earliest cohort born in 1954-58. This pattern was not observed in males of the same age.

- For females born in 1999-2003, the suicide rate reached 6.4 deaths per 100,000 cohort members at age 15-19—considerably higher than females born in 1954-58 (3.6 deaths per 100,000 cohort members).

Suicide rates at age 45-49 have increased with each successive birth cohort in both males (from 24.1 in those born in 1954-58 to 29.5 deaths per 100,000 cohort members in those born in 1969-73) and females (from 6.7 to 9.5 deaths per 100,000 cohort members in the same cohorts).

Suicide rates across male cohorts compared at the same age show no clear pattern. Rates at younger ages of death (15-19 and 20-24) tended to be higher for those born prior to 1979-83 than in those born in more recent cohorts (1984-1988 onwards).

- For males born in 1984-88 the suicide rate at age 20-24 was almost half that of the cohort born in 1969-73 (18.0 deaths per 100,000 cohort members compared with 35.1).
- Rates of suicide at age 15-19 for males born in 1974-78 were 1.8 times higher than those with the lowest rate born in 1954-58 (18.0 deaths per 100,000 cohort members and 10.0, respectively). Rates at age 15-19 were 12.3 deaths per 100,000 cohort members in the most recent male birth cohort for which data are available (1999-2003).

## Trends in methods of suicide by birth cohort and age at death

Understanding the methods used for suicide can play an important role in suicide prevention. These data are provided to inform discussion around restriction of access to means as a policy intervention for the prevention of suicide.

Please consider your need to read the following information. If this material raises concerns for you or if you need immediate assistance, please contact a [crisis support service](#), available free of charge, 24 hours a day, 7 days a week.

Please consider the [Mindframe guidelines](https://mindframe.org.au/suicide/communicating-about-suicide/mindframe-guidelines) (https://mindframe.org.au/suicide/communicating-about-suicide/mindframe-guidelines) if reporting on these statistics.

The classification system used to code causes of deaths data, ICD-10, uses the term 'mechanism' to refer to the external cause of death. Throughout *Suicide & self-harm monitoring* 'mechanism' has been used in data visualisations, while the term 'method' has been used in the accompanying text.

The interactive data visualisations show which methods underlie changes in suicide rates as people in each birth cohort have aged (top visualisation)—and underlie changes in suicide rates for people of the same age, but born at different times (second visualisation).

Rates of suicide by hanging (ICD-10 X70):

- tended to increase for both male and female birth cohorts as the cohort aged (top visualisation).
- tended to increase in most age groups with each successive birth cohort in females; the pattern in males was less consistent (second visualisation). For example:
  - rates of suicide by hanging at ages 15-19 increased for each successive female birth cohort from a low of 0.1 per 100,000 cohort members in those born in the earliest cohort (1954-58) to a high of 4.9 in those born in the most recent cohort (1999-2003).
  - in the 2 most recently born female cohorts for which there are data available at ages 15-19 (born in 1994-98 and 1999-2003), rates of suicide by hanging were as high or higher than, rates at almost any other age in all other female cohorts.
  - for males, rates of suicide by hanging at ages 15-19 do not show the same pattern as females; rates in male cohorts increased up until those born in 1979-1983 and have since remained at about the same level (9.5 deaths per 100,000 cohort members for the latest birth cohort, born 1999-2003).

Rates of suicide by use of firearms (ICD-10 X72-X75) for both males and females peaked at younger ages (15-19 or 20-24) in all birth cohorts and then declined as cohorts aged (top visualisation). Suicide rates by this method tended to be lower for each successive birth cohort at all ages for which there are data available.

- Each more recently born male cohort (born 1969-73 to 1989-93) had successively lower suicide rates by use of firearms at age 20-24 (7.3, 3.6, 1.5, 1.0 and 0.8 deaths per 100,000 cohort members).
- A similar pattern was seen for female cohorts; however, rates were low.

Rates of suicide due to exposure to poisons excluding gas (ICD-10 X60-X66, X68-X69) in female cohorts were similar to that of male cohorts throughout the period 1964 to 2018 (0-3.6 deaths per 100,000 cohort members compared with 0-4.3, respectively)—unlike that of other suicide methods (top visualisation). Rates of suicide by this method were still rising for most male and female cohorts at the end of the available data.

## Reference

Australian Institute of Health and Welfare. Henley G & Harrison JE 2020. Suicide in Australia: Trends and analysis-1964 to 2018. Injury research and statistics series no. 132. Cat. no. INJCAT 212. Canberra: AIHW.





## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on [13 11 14](tel:131114), or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)

## Deaths by suicide, by states and territories

Patterns of deaths by suicide between states and territories can reveal insights that may be masked by results for the whole of Australia. Variations in the rates of deaths by suicide across states and territories may help to highlight different risk factors and assist in better targeting of suicide prevention activities. For example, differences in the ratio of urban to regional and remote areas may explain some of the differences across states and territories given that the rates of suicide tend to be higher in regional and remote areas, see [Suicide by remoteness areas](#).

Information based on the deceased's usual state or territory of residence is available for deaths registered after 1979. Deaths by suicide may be presented by either year of death or by year of registration. Reporting by year of death can provide more reliable information on trends in occurrence than reporting by year of registration; however, the latest data available underestimates the occurrence of recent deaths due to a lag in registration, for more information, see [Technical notes](#). Here, data based on both year of registration of death and year of occurrence of death are presented.


Suicide deaths by states and territories, Australia, 1979 to 2022.

The line graph shows age-standardised suicide rates for each state and territory and the Australian total from 1979 to 2022. Users can choose to view age-standardised suicide rates, numbers of deaths by suicide, year-on-year change in age-standardised suicide rate and year-on-year change in numbers of deaths by suicide. Data can be viewed either by year of registration or year of death.



## Suicide deaths by states and territories, Australia, 1979 to 2022

Death by year of registration  
Age-standardised rate (per 100,000)



Select state/territory:  
(hold Ctrl for multi-select and Esc to show all)

Aus NSW Vic Qld WA SA Tas ACT NT

Select year range: 1979 to 2022

Select measure:  
Age-standardised rate (per 100,000)

Select counting period:  
Death by year of registration

\*Age-standardised rates are not shown where there were fewer than 20 deaths in an area due to unstable rates.

Source: AIHW National Mortality Database and ABS Causes of Death, Australia 2022

Supplementary Table: NMD S4, NMD S5

Latest data: 2023 (annual release)

See notes ►

### How do suicide rates vary across states and territories?

From 1979 to 2022, age-standardised rates of death by suicide (according to year of registration):

- tended to be lower for New South Wales, Victoria and the Australian Capital Territory compared to the overall Australian suicide rate
- tended to be highest in the Northern Territory. However, the Northern Territory also had one of the lowest number of deaths by suicide, when comparing all jurisdictions.

In 2022:

- the age-standardised rate of death by suicide ranged from 10.8 per 100,000 population in New South Wales to 20.5 per 100,000 in the Northern Territory.

Age-standardised suicide rates allow for comparisons between states and territories by adjusting for differences in age structures and population size. Rates fluctuate over time—particularly in the smaller jurisdictions—due to the small number of deaths by suicide that are registered each year. Differences in coronial processes, data processing or coding practices should also be taken into consideration when comparing data across jurisdictions and over time. Caution is advised when comparing state and territory data.

In 1979, the largest number of deaths by suicide (according to year of registration) was in:

- New South Wales (539 deaths), followed by Victoria (462), Queensland (296), South Australia (178) and Western Australia (116).

By 2022, the largest number of deaths by suicide was in:

- New South Wales (911), followed by Queensland (773), Victoria (754), Western Australia (377) and South Australia (242).

However, it should be noted that New South Wales and Victoria have the largest populations in Australia and the populations of both Queensland and Western Australia increased considerably (ABS, 2023).

### What is the effect of reporting suicide deaths by year of occurrence?

The data for age-standardised rates and number of suicide deaths are broadly similar when analysed by year of death or by year of registration. Data for the most recent years, regardless of whether by year of death or year of registration, are impacted by incomplete coronial processes. The most recent years of data by year of death, are additionally impacted by a lag between the occurrence of a death and the registration of the death. This contributes to differences seen between data reporting suicide by year of death and data reporting by year of registration.

## References

ABS (Australian Bureau of Statistics) (March 2023), *National, state and territory population*

<https://www.abs.gov.au/statistics/people/population/national-state-and-territory-population/latest-release>, ABS Website, accessed 4 October 2023.

---

© Australian Institute of Health and Welfare 2024



## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on [13 11 14](tel:131114), or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)

## Australian prevalence estimates of suicidal behaviours

If at any point you feel worried about harming yourself while viewing the information on this website - or if you think someone else may be in danger - please stop reading and [seek help](#).

Suicidal thoughts and behaviours include suicidal ideation (thoughts about taking one's own life), making suicide plans and attempting suicide.

People who experience suicidal ideation and make suicide plans are at increased risk of suicide attempts and those who experience all forms of suicidal thoughts and behaviours are at greater risk of dying by suicide (Demesmaeker et al. 2022, Mendez-Bustos et al. 2013).

Nationally representative population surveys can collect data on the prevalence of suicidal thoughts and behaviours in Australia, the incidence of which may be underestimated in administrative datasets such as the National Hospital Morbidity Database and the National Ambulance Surveillance System (see [Intentional self-harm hospitalisations](#) and [Ambulance attendances](#)). National survey data play an important role in identifying population groups at increased risk of suicidal behaviours and informing suicide prevention activities and efforts to reduce stigma and increase help-seeking behaviour.

The National Survey of Mental Health and Wellbeing program of surveys began in the late 1990s. In 2020 the Australian Bureau of Statistics (ABS) National Study of Mental Health and Wellbeing was introduced as a component of the wider Intergenerational Health and Mental Health Study. The first of these, the 2020-22 National Study of Mental Health and Wellbeing measured the 12-month and lifetime prevalence of mental illnesses in Australia for the first time since the 2007 National Survey of Mental Health and Wellbeing. In addition to data on suicidal thoughts and behaviours, which were collected in 2007, the 2020-22 study also collected data on: health service use due to thoughts or plans to take one's own life; confiding in another person about thoughts or plans to take one's own life, being close to someone who took or attempted to take their own life; whether services were used afterwards; and self-harm behaviours (intentional self-harm without suicidal intent). The 2020-22 study was also the first ABS collection to use the Standard for Sex, Gender, Variations of Sex Characteristics and Sexual Orientation Variables (ABS 2020, 2020-22b). However, to date, only data disaggregated by sex assigned at birth (male/female) have been reported for suicide and self-harm variables.

Summary statistics for the 2020-22 National Study of Mental Health and Wellbeing were released by the ABS on 5 October 2023 (ABS 2020-22a). In this publication, the results for suicide and self-harm are disaggregated by age-group and sex assigned at birth (male/female). The results for suicidal thoughts and behaviours are not directly comparable with those from the 2007 National Survey of Mental Health and Wellbeing as different questions were used.

Results from the 2020-22 National Study of Mental Health and Wellbeing (ABS 2020-22a) indicate that:

- One in 6 (16.7% or around 3.3 million) Australians aged 16-85 had experienced serious thoughts about taking their own life at some point in their lives.
- Around 1.5 million or 7.4% of Australians aged 16-85 years had made a suicide plan and around 970,000 or 4.9% had attempted suicide during their lifetime.
- Females were more likely to be suicidal than males, with a higher prevalence of suicidal thoughts and behaviours in their lifetime (18.3% compared with 15.0%). These findings are in contrast to the data on deaths by suicide, which show that males are more likely than females to die by suicide; visit [Deaths by suicide over time](#).
- Young people and adults aged 16-34 years reported the highest prevalence of suicidal thoughts and behaviours in the 12 months before the administration of the study (4.9% of people aged 16-34 years).

- In their lifetimes, 36.2% of Australians aged 16-85 years were close to someone who took or attempted to take their own life, while 4.7% were close to someone who took or attempted to take their own life in the 12 months prior to the study.
- The prevalence of lifetime and past 12 months self-harm (without suicidal intent) was highest in the youngest age group (16-24, 20.4% and 6.0%, respectively) and decreased with increasing age (2.2% and 0.4%, respectively for those aged 55-85).
- Self-harm prevalence was higher for females than for males. 27.9% of females aged 16-24 had self-harmed in their lifetimes, 8.7% in the past 12 months, compared with 13.6% and 3.3% of males in this age group, respectively.

For full results visit [National Study of Mental Health and Wellbeing \(https://www.abs.gov.au/statistics/health/mental-health/national-study-mental-health-and-wellbeing/2020-2022\)](https://www.abs.gov.au/statistics/health/mental-health/national-study-mental-health-and-wellbeing/2020-2022). Please note that some of the 12-month prevalence estimates have high relative standard error (RSE) and margin of error (MOE), and thus wide confidence intervals and greater uncertainty. These results should be interpreted with caution. Refer to the ABS technical notes on interpretation of results with high RSEs & MOEs (visit [National Study of Mental Health and Wellbeing methodology \(https://www.abs.gov.au/methodologies/national-study-mental-health-and-wellbeing-methodology/2020-2022\)](https://www.abs.gov.au/methodologies/national-study-mental-health-and-wellbeing-methodology/2020-2022)). See [glossary](#) for a definition of confidence intervals.

## References

ABS (Australian Bureau of Statistics) (2020) *Standard for Sex, Gender, Variations of Sex Characteristics and Sexual Orientation Variables* (<https://www.abs.gov.au/statistics/standards/standard-sex-gender-variations-sex-characteristics-and-sexual-orientation-variables/2020>), ABS website, accessed 4 July 2023.

ABS (Australian Bureau of Statistics) (2020-2022a) *National Study of Mental Health and Wellbeing*, ABS website (<https://www.abs.gov.au/statistics/health/mental-health/national-study-mental-health-and-wellbeing/2020-2022>), accessed 17 October 2023.

ABS (Australian Bureau of Statistics) (2020-22b) *National Study of Mental Health and Wellbeing methodology* (<https://www.abs.gov.au/methodologies/national-study-mental-health-and-wellbeing-methodology/2020-2022>), ABS website, accessed 2 June 2023.

Demesmaeker A, Chazard E, Hoang A, Vaiva G & Amad A (2022) 'Suicide mortality after a nonfatal suicide attempt: A systematic review and meta-analysis', *The Australian and New Zealand Journal of Psychiatry*, 56(6): 603-616, <https://doi.org/10.1177/00048674211043455>.

Mendez-Bustos P, de Leon-Martinez V, Miret M, Baca-Garcia E, Lopez-Castroman J (2013) 'Suicide reattempters: a systematic review', *Harvard Review of Psychiatry*, 21(6): 281-295, doi: 10.1097/HRP.0000000000000001.



## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on [13 11 14](tel:131114), or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)

## Patterns of health service use in the last year of life among those who died by suicide

A significant proportion of people who die by suicide have contact with the health system in their last year of life. These contacts points provide a potential touch point for suicide prevention activities. As Clapperton et al. (2021) argue, many prevention activities focus on people who access hospitals. However, Clapperton et al. (2021) also show significant proportions of people who die by suicide (particularly men) do not attend hospitals in their last year of life. It is important to focus not just on people who use services but also on people who do not access services.

Through this project we have used the National Integrated Health Services Information Analysis Asset (NIHSI AA) version 0.5 to look at patterns of health service use in the last year of life for people who have died by suicide. The main value add of this project, compared to earlier studies, comes from fact that the NIHSI AA includes both Medicare Benefits Schedule (MBS) and Pharmaceutical Benefits Scheme (PBS) data. This is an important distinction as we show that people who die by suicide are considerably more likely to access MBS and PBS services in their last year of life than hospital services.

While this project provides the most comprehensive analysis of health service use in the last year of life for people who die by suicide it does have some important limitations. First the NIHSI AA does not include all health services. For example, it does not include community or residential mental health services. Second the emergency department (ED) data in the NIHSI AA does not identify intentional self-harm well as intentional self-harm is not identified well in ED data in most States and Territories (AIHW 2022a).

An additional limitation of this analysis relates to the fact that not all mental services under the MBS are billed as specific mental health items – some are billed under general GP items (AIHW 2022b).

The project has been established to measure and compare the patterns of health service use of Australians in their last year of life.

Specifically, in this analysis we aim to:

- measure the patterns of health service use in the last year of life
- identify key factors related to variation in the patterns of health service use in the last year of life such as age and sex.

Using linked datasets including the National Deaths Index, National Hospital Morbidity Database, National Non-admitted Patient Emergency Department Care Database, MBS and PBS, the NIHSI AA presents descriptive statistics to answer the research questions. While the MBS and PBS databases include national data, hospital data pertains to only New South Wales, Victoria (excluding Albury-Wodonga), South Australia and Tasmania public hospitals within the NIHSI AA v0.5. Admitted patient information also contains information from private hospitals in Victoria. To ensure accurate comparisons with hospitals data, only deaths registered in these jurisdictions are included in the analysis.

The analysis population was those who had died between 1 July 2010 and 31 December 2017 in the linked National Deaths Index. Only people whose age at death was between 15 to 64 years were included in the analysis. This was due to people in this age range making up the majority of those who die from suicide and to allow for better comparisons with deaths from other causes, which mostly occur in people older than 65 (AIHW 2022c). In the analysis population, people aged 15-64 years represented 82% of suicides (10,013 suicide deaths).

For further information on the dataset and methods used, visit [Technical notes - Data sources](#).

## People who died by suicide accessed fewer health services in their last year of life than those who died from other causes

Overall, 49% of 15-64 year olds who died by suicide **did not have** any contact with the hospital (emergency department (ED) presentation or hospital separation), compared to 24% who died by other causes. This is similar to results from Clapperton et al. (2021), who found that 50% of people who died by suicide in Victoria did not have any ED presentations or hospital separations in their last year of life, using data from the Victorian Suicide Register and including all age groups.


In addition to looking at overall access to hospitals it is also worth exploring access to individual services and how access to these services vary by age and sex.

- Females (59%) who died by suicide were considerably more likely than males to attend hospital for any reason in their last year of life (48%). There is also variation by age and sex with females aged 34-44 being the most likely to attend hospital (61%) and males aged 55-64 being the least likely to attend (45%).
- Females (53%) were more likely than males to attend an emergency department for any reason in their last year of life than males (44%). The highest rate was for females aged 25-34 (56%) while the lowest rate is for males aged 55-64 (39%).
- 9% of females and 6% of males who died by suicide had a mental health related ED presentation in their last year of life.
- While it is not possible to identify intentional self-harm ED presentations in the NIHSI AA it is possible to identify intentional self-harm ED presentations in Victoria. Clapperton et al. (2021) found that the vast majority of both males (92%) and females (84%) who died by suicide did not attend hospital for intentional self-harm in their last year of life as either an admitted patient or in an emergency department.
- A higher proportion of females (47%) who died by suicide were an admitted patient in hospital in their last year of life than males (32%). The highest proportion was for females aged 45-54 (49%) while the lowest proportion was for men aged 15-24 (26%).
- Only 13% of women and 6% of men who died by suicide were an admitted patient in hospital for intentional self-harm in their last year of life.
- Females (30%) who died by suicide were more likely than males (19%) to have had a mental health hospitalisation (this excludes ED presentations) in their last year of life.
- Both females (90% v 59%) and males (79% v 48%) who died by suicide were considerably more likely to have used MBS services than hospital services in their last year of life.
- Females who died by suicide (57%) were considerably more likely to have used MBS mental health services than males (37%) in their last year of life.
- Females (71%) who died by suicide were also more likely than males (50%) to have had a PBS mental health prescription in their last year of life.

The interactive data visualisation shows the proportion of health services used in the last year of life for people who died by suicide and for people who died by other causes. The service type can also be selected. It is displayed by age group from 15-64 and sex for deaths between 1 July 2010 and 31 December 2017.

### Proportion of health services used by sex, age, service type cause of death and whether had a service , 2010-11 to 2017

Proportion of people (Any hospitalisation (including ED))



1. Includes deaths registered to New South Wales, Victoria (excluding Albury-Wodonga), South Australia and Tasmania  
 2. *Did not have a service* refers to people who did not had the selected service within 12 months of death. *Did have a service* refers to people who had the selected service within 12 months of death.  
 3. *Did not have a service* and *Did have a service* proportions add up to 100%. The denominator is the total number of people who died by the selected cause of death.  
 Source: NIHSI  
 Supplementary Table: NIHSI S2


[See notes ►](#)

Of those who died from suicide from 1 July 2010 and 31 December 2017:

- 11% did not access any of the health services analysed in their last year of life
- Over 1 in 10 males (13%) who died by suicide did not access a service in their last year of life. Males were less likely to have accessed any services than females in every age group.
- Overall, younger age groups and males had higher proportions of not accessing a service in their last year of life compared to older age groups and females.
- Males aged 15-24 who died by suicide had the highest proportion of not accessing services in their last year of life (19%), while females aged 55-64 years who died of other causes had the lowest proportion (2.6%).
- Around 10% of females aged 15-24 who died by suicide did not use any of the health services analysed in their last year of life.

The data visualisation shows the amount of health services used in the last year of life for people who died by suicide and for people who died by other causes. It is displayed by age groups between 15-64 and sex for deaths registered between 1 July 2010 and 31 December 2017. For males and females of all age groups a higher percentage of services was used when the cause of death was not suicide.

Proportion of health service usage in last year of life by sex, age, cause of death, 2010-11 to 2017



Notes: Includes deaths registered to New South Wales, Victoria (excluding Albury-Wodonga), South Australia and Tasmania  
 Source: NIHSI  
 Supplementary Table: NIHSI S1

[See notes ►](#)


### People who died by suicide had more ED presentations in their last year of life than those who died by other causes

Of those who did access a health service in their last year of life, MBS and PBS services represented the highest proportion of services among those who died by either suicide or other causes. For those who died by suicide, the next most prevalent health service after MBS and PBS was ED presentations (3.8%) then hospital separations (2.4%).

- Out of the four selected service groups, the proportion of ED presentations and hospital separations decreased in those who died by suicide with increasing age, for both males and females.
- MBS services also decreased with increasing age among those who died by suicide. For instance, the proportion of MBS services out of the four selected service groups was 64% in females aged 15-24 and 47% in females aged 55-64.

The interactive data visualisation shows the type of health services used in the last year of life. The user can display the data by female, male or persons. Data is categorised by age groups from 15-64, causes of death by suicide and other causes of death and service type used between 1 July 2010 and 31 December 2017.

**Proportion of health services used in last year of life by sex, age, service type and cause of death, 2010-11 to 2017**



Notes: Includes deaths registered to New South Wales, Victoria (excluding Albury-Wodonga), South Australia and Tasmania  
 Source: NIHSI  
 Supplementary Table: NIHSI S3

[See notes ►](#)

**Young people who died by suicide had higher health service use than those who died by other causes**

Suicide is the leading cause of death among people aged 15-44, while chronic diseases feature more prominently among people aged 45 and over. In general people who die by suicide are younger than people who die by other causes (AIHW 2022c). This is reflected in these data for health service use. For example, among those who died by suicide and had any hospital contact in last year of life (including ED), 59% of those hospital contacts were in the 15-44 age group, compared to 17% of the same age who died of other causes. Most people who died by other causes and had a hospital contact were aged 45-64 (83%). Any mental health, suicidal ideation or intentional self-harm hospitalisations or ED mental health presentations also follow this pattern.


- Of those who presented to ED and died from suicide, one quarter (25%) were people aged 35-44, whereas only 11% of those who presented to ED and died from other causes were in this age group.
- Out of those who died by suicide, the average number of services per person for any mental health MBS services was higher than for those who died by other causes, across all age groups.

The interactive data visualisation shows health services used in last year of life for those that died by suicide or other causes, by age group for deaths between 1 July 2010 and 31 December 2017. The user can display the data by measure (proportion of people, proportion of services, average number of services per person), sex (males, females, persons) and service type used.



## Health services used in last year of life, by cause of death, age and service type, 2010–11 to 2017

Proportion of people (Any hospitalisation (including ED))



### Notes:

- Includes deaths registered to New South Wales, Victoria (excluding Albury-Wodonga), South Australia and Tasmania
- Age group proportions add up to 100%. The denominator is the total number of people/services for the selected service type and cause of death.

Source: NIHSI

Supplementary Table: NIHSI S3

[See notes ►](#)

## A higher proportion of health services occurred in the month prior to death

Among those who died by suicide, the highest proportion of hospital contacts (ED or hospital separation) occurred 1 month prior to death (18%), out of a 12 month period. This might indicate increased risk following the use of some services. However, the average number of services is similar across all months meaning that those who did receive a service one month prior to death did, on average, not attend more often than in previous months. Note that in this analysis, people who died in hospital or during their ED presentation were excluded to capture their service use prior to death, except for those who had an intentional self-harm diagnosis during their episode of care (see [Technical notes - Data sources](#) for an explanation on analytical method used).


Per person, of those who died by suicide:

- The average number of any MBS mental health or mental health treatment plan service is higher than those who died from other causes between 12 months and 1 month prior to death.
- The average number of any MBS service decreases sharply between 3 months prior to death and 1 month prior to death, while remaining somewhat steady in those who died from other causes.

The interactive data visualisation shows service use in the 12 months leading up to death. Users can display data by measure (proportion of people, proportion of services, average number of services per person), sex (males, females, persons) and service type. Most health services occur in the month prior to death.

### Month of service prior to death, by cause of death and service type, 2010-11 to 2017

Proportion of people (Any hospitalisation (including ED))



Select measure:  Proportion of people  Proportion of services  Average number of services per person

Select sex:  Males  Females  Persons

Select service type:  Any hospitalisation (including ED)

Suicide  Other causes

Note: Includes deaths registered to New South Wales, Victoria (excluding Albury-Wodonga), South Australia and Tasmania  
 Source: NIHSI  
 Supplementary Table: NIHSI S4

[See notes ►](#)


Among those who died by suicide:

- The cumulative total proportion of ED mental health presentations in the lead up to death is lower than those who died from other causes between 12 to 2 months prior to death, indicating lower service usage until 1 month prior to death.
- The cumulative total proportion of any MBS service is lower than those who died by other causes between 12 to 1 month/s prior to death.

The interactive data visualisation shows the cumulative service use by sex and cause of death in the 12 months leading up to death. Users can display data by sex and service type.

### Cumulative proportion of people using health services, by sex, cause of death and months prior to death, 2010-11 to 2017

Cumulative proportion



Select sex:  Males  Females  Persons

Select service type:  Any hospitalisation (including ED)

Suicide  Other causes

Notes: Includes deaths registered to New South Wales, Victoria (excluding Albury-Wodonga), South Australia and Tasmania  
 Source: NIHSI  
 Supplementary Table: NIHSI S4

[See notes ►](#)

The AIHW will undertake further analysis on these data including multivariate modelling. While these data are informative more insights can be gained by looking at how the use of these various health services compares to the population in general. For example, while the majority of people who die by suicide did not have a mental health hospitalisation in their last year of life they are considerably more likely to have done so than the population in general.

## References

---

Australian Institute of Health and Welfare (AIHW) (2022a) [Suicide and self-harm monitoring: Intentional self-harm hospitalisations](#), AIHW, Australian Government, accessed 14 October 2022.

Australian Institute of Health and Welfare (2022b) [Mental health services in Australia](#), AIHW, Australian Government, accessed 09 September 2022.

AIHW (2022c) [Deaths in Australia](#), AIHW, Australian Government, accessed 13 September 2022.

AIHW (2021) [Suicide & self-harm monitoring: Social factors and deaths by suicide](#), AIHW, Australian Government, , accessed 01 February 2022.

Clapperton A, Dwyer J, Millar C, Tolhurst P and Berecki-Gisolf J (2021) [Sociodemographic characteristics associated with hospital contact in the year prior to suicide: A data linkage cohort study in Victoria, Australia](#) (<https://doi.org/10.1371/journal.pone.0252682>), *PLoS ONE*, 16(6): e0252682, doi:10.1371/journal.pone.0252682.

---

© Australian Institute of Health and Welfare 2024



## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on [13 11 14](tel:131114), or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)

## Suicide registers

In Australia, the coroners court in each state and territory is responsible for investigating suspected deaths by suicide. Most Australian jurisdictions have established suicide registers to record the information provided to coroners at the time a suspected suicide death is referred for investigation. These surveillance systems provide close to real-time data and are valuable for informing responses, research, and policy in suicide prevention locally, and across national and international levels.

Prior to the COVID-19 pandemic, suicide registers existed in Queensland (established in 1990), Victoria (2012) and Tasmania (2017). New South Wales established a suicide register in October 2020. Through the Suicide and Self-harm Monitoring Project, the AIHW has worked with governments in the Australian Capital Territory, South Australia, and the Northern Territory to establish suicide registers in these jurisdictions. These registers became operational in 2021, 2022 and 2023, respectively. Visit [Data development activities](#) to read more.

Several jurisdictions have published reports on their suicide register data, including:

- New South Wales (monthly reports up to January 2024)
- Victoria (monthly reports up to March 2024)
- Queensland (monthly reports up to February 2024)

These reports are discussed further in the next section. It is important to note that suicide is not influenced or caused by one factor - but results from a complex interaction between multiple risk factors (Leske et al. 2022).

Data on suspected deaths by suicide are based on initial police reports and other information available at the time of referral to the coroner. As such, they are not directly comparable with cause of death data released by the Australian Bureau of Statistics, which are based on final coronial determinations. However, the differences are generally small. For example, in the case of the Victorian Suicide Register (VSR):

‘VSR analyses have shown that over time, there is consistently less than 5% difference between the number of suicides initially identified as suicide, and the number of deaths ultimately confirmed as suicides’ (CCOV 2022).

The state and territory suicide registers also differ from each other in their processes and counting rules for identifying suspected suicide deaths. Therefore, data from one register cannot be directly compared with those from another.

For more information on suicide register data custodians with published data, visit [Data sources](#).

## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on [13 11 14](tel:131114), or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)

## Data from suicide registers

### New South Wales

The New South Wales Suicide Monitoring System (NSW SuMS) was established in October 2020. The NSW SuMS is a collaboration between NSW Health, the Department of Communities and Justice (DCJ), the State Coroner and NSW Police. NSW Health publishes monthly reports on suspected and confirmed deaths by suicide occurring in New South Wales. Data on suspected deaths by suicide are an estimate, and numbers for the same period may differ slightly between reports as the coroners' determinations into the deaths are finalised. Caution is advised against drawing any conclusions about suicide trends in NSW based on short-term changes.


The latest NSW SuMS report for January 2024 shows that (NSW Health 2024):

- A total of 940 suspected or confirmed suicide deaths was recorded for the full year in 2023. This was lower than the number of suicide deaths in 2022 (960) and higher than in 2021 (907), and 2020 (896).
- 83 suspected deaths by suicide were recorded between 1 and 31 January 2024. This compares to 89 suspected or confirmed suicide deaths for the same period in 2023, 80 in 2022, 101 in 2021 and 79 in 2020.

The number (frequency) of suspected or confirmed suicide deaths per month varies considerably from month to month, as can be seen in the visualisation below when the "Frequency" view is selected. The "Cumulative Frequency" view shows the year-to-date numbers of suicide deaths. From this view it appears that the increase in suicide numbers in 2022 began from around April 2022.

### Frequency of suspected and confirmed deaths by suicide in New South Wales, by month, January 2019 to January 2024

The interactive data visualisation shows the number of suspected and confirmed deaths by suicide in New South Wales, by month, beginning from January 2019 up to February 2023. Viewing can be changed between frequency and cumulative frequency. An average trendline has been included.



Notes: Data are subject to revision as coronial investigations progress and are finalised. Data for the same period may therefore vary between reports.  
 Source: NSW Health (2024)

The NSW SuMS also reports on suicide deaths by gender, age group and residential location (NSW Health 2024):

- In 2023 there were 733 suspected or confirmed suicide deaths reported for males in NSW compared with 745 in 2022, 670 in 2021 and 673 in 2020.
- For females, there were 207 suspected or confirmed suicide deaths in 2023, compared with 214 in 2022, 237 in 2021, and 223 in 2020.
- Nearly four-fifths (78%) of suspected or confirmed suicide deaths in NSW in 2023 were among males.
- For males, the age groups of 45 to 54 years (147), 55 to 64 years (131), 35 to 44 years (120), and 25 to 34 years (114) recorded the largest number of suspected or confirmed suicide deaths, accounting for 20%, 18%, 16% and 16% of all male suicide deaths, respectively.
- Similarly, for females, the age groups of 35 to 44 years (46), 25 to 34 years (35), 45 to 54 years (34) and 55 to 64 years (31) experienced the greatest number of suspected or confirmed deaths by suicide, accounting for 22%, 17%, 16% and 15% of all female suicide deaths, respectively.
- Among males there was an overall decrease in the number of suspected or confirmed suicides of 12 in 2023 compared with 2022. The age group of males with the largest decrease (23) was 35 to 44 (143 deaths in 2022 compared to 120 in 2023).
- Among females there was an overall decrease in the number of suspected or confirmed suicides of 7 in 2023 compared with 2022. Among individual age groups of females there were no decreases greater than 5. In the female 35 to 44 years age group, the number of suspected or confirmed suicides increased by 9 (37 in 2022 to 46 in 2023).
- In each year from 2019 to 2023, around half of suicide deaths in NSW occurred among residents of Greater Sydney, with the remainder comprising residents of the Rest of NSW and a small number of interstate/overseas residents.

#### Frequency of suspected and confirmed deaths by suicide in New South Wales, by age group and sex, and location, 2019 to 2023

The interactive data visualisation shows the number of suspected and confirmed deaths by suicide in New South Wales, starting from 2019 up to 2022. The population group is divided by sex (males, females) and age groups, ranging from people under the age of 18 years to 85 years and over. Viewing by location of usual residence can also be selected.

Year-to-date data to 31 January 2024 from the SuMS show:

- Four-fifths (80%) of suspected deaths by suicide were among males. This compares with the full-year data for previous years, with males accounting for 78% of suspected or confirmed suicide deaths in 2023, 78% in 2022, 74% in 2021, and 75% in 2020.

#### Frequency of suspected and confirmed deaths by suicide in New South Wales, by age group and sex, and location, 1 to 31 January 2019 to 2024

The interactive data visualisation shows the number of suspected deaths by suicide in New South Wales. The population group is divided by age group, ranging from people under the age of 18 years to over 85, and sex (males, females). Viewing by location of usual residence can also be selected.

The Coroners Court of Victoria (CCOV) established the Victorian Suicide Register (VSR) in 2012 and publishes monthly data reports on suspected and confirmed deaths by suicide. VSR data are regularly reviewed, where deaths may be added or removed from the register as coronial investigations progress and are finalised. VSR data may therefore change over time.

The latest *Monthly Suicide Data Report* shows (CCOV 2024c):

- There was a total of 801 suspected or confirmed suicide deaths in 2023. This was higher than the number of suspected or confirmed suicide deaths in 2022 (766), 2021 (681), and 2020 (674).

The CCOV stated that there was an increase in the number of suicides in 2022 and 2023 compared to the period from 2019 to 2021, which was relatively stable (CCOV 2024b).

Year-to-date data to March 2024 from *Monthly Suicide Data Report* shows (CCOV 2024c):


- Between 1 January and 31 March 2024, 222 suspected deaths by suicide were reported in Victoria. This is higher than in the same period in 2023 (189), 2022 (172), 2021 (177), and 2020 (186).

As illustrated below, the monthly frequency data show considerable variation which, according to the CCOV, usually results from random factors rather than underlying systemic issues or emerging clusters (CCOV 2022). The data therefore should be interpreted cautiously, with great care taken in drawing conclusions about any apparent short-term increase or decrease that is observed.

The “Cumulative Frequency” view in the visualisation below shows that the increase in 2022 began in the second half of the year and ‘is in contrast to the preceding four years, in which Victoria had seen a plateau in suicide numbers’ (CCOV 2023a). The number of suspected or confirmed suicides between January and July 2022 was consistent with previous years, with a monthly average of 58 deaths. However, between August and December 2022, the average monthly frequency increased to 71 deaths, which ‘might signal an emerging trend’ (CCOV 2023a). This increase in suspected or confirmed suicide deaths appears to have continued in 2023 and the first quarter of 2024, with monthly averages of 67 and 74 deaths respectively, both higher than previous years.

#### Frequency of suspected and confirmed deaths by suicide in Victoria, by month, January 2016 to March 2024

The interactive data visualisation shows the number of suspected deaths by suicide in Victoria, by month, starting from January 2016 to January 2023. Viewing can be changed between frequency and cumulative frequency. An average trendline has been included.



Notes: Data are subject to revision as coronial investigations progress and are finalised. Data for the same period may therefore vary between reports. Data for 2016, 2017, 2018, and 2019 are correct as of January 2021, January 2022, February 2023, and February 2024 respectively, as the Coroners Court of Victoria (CCOV) has not published more recent data for these years. Any revisions to 2020 data and onwards will continue to be updated as published by the CCOV.  
Source: CCOV (2021, 2022, 2023c, 2023d, 2024c)

The CCOV also reports on suicide deaths by age group and sex, and incident location (CCOV 2024a, 2024b). For 2023:

- Males accounted for 72% of suspected or confirmed suicide deaths. However, the increase in deaths in 2023 was more pronounced for females than males. In 2023, the total number of suspected or confirmed suicide deaths among females increased by 24 from 2022 (12% increase from 2022). For males, the increase from 2022 to 2023 was 16 (2.9% increase from 2022).
- Among males, the age groups of 35 to 44 years (120), 45 to 54 years (108) and 25 to 34 years (105) recorded the largest number of suicide deaths, accounting for 21%, 19% and 18% of all male suicide deaths, respectively.

- Similarly, for females, the age groups of 35 to 44 years (48), 25 to 34 years (45), and 45 to 54 years (41) experienced the greatest number of deaths, accounting for 21%, 20% and 18% of all female suicide deaths, respectively.
- The highest increase was seen in people aged 35 to 44 years, with 168 suspected or confirmed suicide deaths compared to 129 in 2022.
- The largest percentage increase was seen in those under 18 years, with an 86% increase in 2023 from 2022 (26 deaths in 2023 compared with 14 in 2022). In April 2023, the CCOV investigated the increase in suicide deaths among under 18s and released a statement noting that the deaths ‘occurred in diverse circumstances across communities in both Metropolitan Melbourne and Regional Victoria, with no clear links established to date between any of the deaths’ (CCOV 2023b).
- Consistent with previous years, the proportion of suicide deaths remained higher in Metropolitan Melbourne (65%) compared to Regional Victoria (35%).

For a detailed breakdowns of full-year frequencies of suspected and confirmed suicide deaths in Victoria by sex and age group, and incident location can be viewed on the visualisation below.

### Frequency of suspected and confirmed deaths by suicide in Victoria, by age group and sex, and location, 2018 to 2023

The interactive data visualisation shows the number of suspected and confirmed deaths by suicide in Victoria, starting from 2018. The population group is divided by sex (males, females) and age groups, ranging from people under the age of 18 to over 65. Viewing by incident location can also be selected.

Year-to-date data to March 2024 for age group and sex, and incident location show (CCOV 2024c):

- Males recorded 166 suspected deaths by suicide, which is higher than the same period in 2023 (137), 2022 (116), 2021 (137), and 2020 (153).
- Females recorded 56 suspected deaths by suicide, compared with 52 for the same period in 2023, 56 in 2022, 40 in 2021, and 33 in 2020.
- The increased number of suspected suicide deaths observed in the first quarter of 2024 mainly occurred in Metropolitan Melbourne. From 1 January to 31 March 2024, the number of suspected suicide deaths in Metropolitan Melbourne (147) was higher than reported for the same period in previous years (117 in 2023, 108 in 2022, 109 in 2021, and 128 in 2020).

Detailed breakdowns of the year-to-date frequency of suspected and confirmed suicide deaths in Victoria by age group, sex, and incident location can be viewed on the visualisation below.

### Year-to-date frequency of suspected and confirmed deaths by suicide in Victoria, by age group and sex, and location, 1 January to 31 March 2019 to 2024

The interactive data visualisation shows the number of suspected deaths by suicide in Victoria, from month end January 2019 to month end April 2023. The population group is divided by sex (males, females) and age group, ranging from people under the age of 18 years to 65 years and over. Viewing by location can also be selected.

The CCOV has also published data on suicides of Aboriginal and Torres Strait Islander (First Nations) people. The number of suspected deaths by suicide in 2023 for First Nations people in Victoria was 28, compared to 18 in 2022, 34 in 2021, 21 in 2020, 19 in 2019, and 14 in 2018. Of those 28 people who were suspected to have died by suicide in 2023, 22 were male. This compares to 13 in 2022, 25 in 2021, 14 in 2020, 12 in 2019 and 10 in 2018. There were 6 female suspected deaths by suicide in 2023, compared to 5 in 2022, 9 in 2021, 7 in both 2020 and 2019, and 4 in 2018 (CCOV 2024d).

Between 2018 to 2023, First Nations people made up an average of 3.1% of people who were suspected or confirmed to have died by suicide in Victoria. In Victoria, the average annual crude rate of death by suicide (suspected or confirmed) was more than twice as high among First Nations people compared with non-indigenous people (28.4 and 10.8 per 100,000 population, respectively) (CCOV 2024d).

Suicide deaths among First Nations people tended to occur at a younger age compared to non-Indigenous people, with 54% of all suicides among First Nations people occurring in those under 35 years of age, compared to 32% for non-Indigenous people (CCOV 2024d).

## Queensland

In Queensland there are two systems that are used to monitor suicide deaths, the Queensland Suicide Register (QSR), which includes suicide data since 1990 and is used to monitor longer-term trends, and the interim Queensland Suicide Register (iQSR), which was established in 2011 to provide real-time information on suicide deaths. The QSR contains information on suicide deaths in Queensland for which the coroners' investigations have been finalised, whereas the iQSR records interim data on deaths suspected to be from suicide, shortly after the death occurs. Data on suspected suicide deaths are based on initial police reports and other information that is available to police at the time they refer the death to the coroner.

Management of the QSR and iQSR was transferred from the Australian Institute for Suicide Research and Prevention (AISRAP) at Griffith University to the Queensland Mental Health Commission (QMHC) in September 2023. Previously, AISRAP published annual reports on suicide in Queensland from the QSR and iQSR. In January 2024, the QMHC commenced publishing monthly data reports from the iQSR. To date, monthly reports have been published for October, November and December 2023, with data dating back to January 2016.

The latest iQSR Monthly Suicide Data Report for February 2024 shows (QMHC 2024b):



- There were 782 suspected suicide deaths in 2023. This was less than the number of suspected suicide deaths recorded in 2022 (795), 2021(816) and 2020 (790).
- 138 suspected deaths by suicide were recorded between 1 January and 29 February 2024. This compares to 137 for the same period in 2023, 135 in 2022, 151 in 2021 and 151 in 2020.

In the visualisation below:

- The number (frequency) of suspected suicide deaths per month varies considerably from month to month, when the “Frequency” view is selected. The number of suspected male suicide deaths appears to be higher in the summer months compared to the winter months. When females are selected this pattern is less evident (See male and female).
- The “Cumulative Frequency” view shows the year-to-date numbers of suspected suicide deaths for each month. For persons, males and females, suspected suicide deaths in 2016 are lower than from 2017 to 2023.

#### Frequency of suspected deaths by suicide in Queensland, by persons, male and female, January 2016 to February 2024

The interactive data visualisation shows the number of suspected deaths by suicide in Queensland, by month. Viewing can be changed between frequency and cumulative frequency. An average trendline has been included.

The iQSR reports on suspected suicide deaths by age group and residential location (QMHC 2024b):

- In 2023, three-quarters of suspected suicide deaths in Queensland were among males. However, while the number of suspected suicide deaths among males decreased from 611 in 2022 to 583 in 2023, the number of suspect suicide deaths among females increased from 184 in 2022 to 199 in 2023.
- In 2023 the highest number of suspected suicide deaths by age group was for people aged 45 to 54 years (149), 25 to 34 years (146), and 35 to 44 years (139) accounting for 19%, 19% and 18% of all suspected suicide deaths respectively. People aged 17 years and under had the fewest deaths (21, 2.7%).
- In 2023 there were 13 fewer suspected suicide deaths than in 2022. The largest decrease in suspected suicide deaths between 2022 and 2023 was in people aged 45 to 54 years with 22 fewer deaths. The largest increase between 2022 and 2023 was for those aged 25 to 34 years with 10 more deaths. For all other age groups, the differences in suspected suicide deaths between 2022 and 2023 was 6 or less.
- In each year from 2020 to 2023 over half of the suspected suicide deaths were among people who resided in major cities, around 40% from inner or outer regional areas, and less than 5% from remote or very remote areas.

The visualisation below contains three different views. The first shows the frequency of suspected deaths by suicide in Queensland by age group and sex from 2020 to 2023. The second shows the data presented by age group (persons only) for years 2016 to 2023, while the third display the data by residential location for years 2016 to 2023.

#### Frequency of suspected deaths by suicide in Queensland by age group, sex, and location 2016-2023

The interactive data visualisation shows the number of suspected deaths by suicide in Queensland, starting from 2016. The population group is divided by age groups, ranging from people under the age of 18 years to 65 or 75 years and over, and sex (males and females). Viewing by incident residential location can also be selected.

The year-to-date iQSR reports (to February 2024) on suspected suicide deaths by age group, sex and residential location show (QMHC 2024b):


- For males there was an increase in the number of suspected deaths by suicide (100 in 2023 to 113 in 2024) while females showed a decrease (37 in 2023 to 25 in 2024).
- The largest year-to-date to February increase was for males aged 55 to 64 years (7 in 2023 to 20 in 2024), although the number in 2024 was similar to 2020, 2021 and 2022 (18, 16, and 16 deaths, respectively).
- By residential location there was a decrease in the number of suspected deaths by suicide in Major Cities (70 in 2023 to 64 in 2024) with an increase in Outer Regional (21 in 2023 to 28 in 2024).


The visualisation below shows the frequency of suspected deaths by suicide in Queensland by age group, sex and residential location for the period of 1 January to the end of February 2020 to 2024.


#### Frequency of suspected deaths by suicide in Queensland by age group, sex, and location, 1 January to end February 2020-2024


The interactive data visualisation shows the number of suspected deaths by suicide in Queensland, starting from 2016. The population group is divided by age groups, ranging from people under the age of 18 years to 65 or 75 years and over, and sex (males and females). Viewing by incident residential location can also be selected.


## References

Coroners Court of Victoria (CCOV) (2021)  [Monthly Suicide Data Report, December 2020 update](https://www.coronerscourt.vic.gov.au/sites/default/files/2021-01/Coroners%20Court%20Monthly%20Suicide%20Data%20Report%20-%20December%202020.pdf)  
<https://www.coronerscourt.vic.gov.au/sites/default/files/2021-01/Coroners%20Court%20Monthly%20Suicide%20Data%20Report%20-%20December%202020.pdf>, 18 January 2021.

CCOV (2022)  [Monthly Suicide Data Report, December 2021 update](https://www.coronerscourt.vic.gov.au/sites/default/files/2021-01/Coroners%20Court%20Monthly%20Suicide%20Data%20Report%20-%20December%202020.pdf) (https://www.coronerscourt.vic.gov.au/sites/default/files/2021-01/Coroners%20Court%20Monthly%20Suicide%20Data%20Report%20-%20December%202020.pdf), 20 January 2022.

CCOV (2023a)  [Coroners Court report shows concerning increase in suicides in 2022](https://www.coronerscourt.vic.gov.au/coroners-court-report-shows-concerning-increase-suicides-2022), (https://www.coronerscourt.vic.gov.au/coroners-court-report-shows-concerning-increase-suicides-2022) 6 February 2023.

CCOV (2023b)  [Increase in youth suicide observed in first three months of 2023](https://www.coronerscourt.vic.gov.au/increase-youth-suicide-observed-first-three-months-2023#:~:text=Troubling%20figures%20released%20today%20by,youth%20suicides%20in%20the%20state.) (https://www.coronerscourt.vic.gov.au/increase-youth-suicide-observed-first-three-months-2023#:~:text=Troubling%20figures%20released%20today%20by,youth%20suicides%20in%20the%20state.), 26 April 2023.


CCOV (2023c)  [Monthly Suicide Data Report, December 2022 update](https://www.coronerscourt.vic.gov.au/sites/default/files/2023-02/Coroners%20Court%20Suicide%20Data%20Report%20December%202022.pdf) (https://www.coronerscourt.vic.gov.au/sites/default/files/2023-02/Coroners%20Court%20Suicide%20Data%20Report%20December%202022.pdf), 6 February 2023.

CCOV (2024a) [Coroners Court 2023 Annual Suicide Data Report](https://www.coronerscourt.vic.gov.au/sites/default/files/2024-02/Coroners%20Court%202023%20Annual%20Suicide%20Data%20Report%20-%20December%202023.pdf) (https://www.coronerscourt.vic.gov.au/sites/default/files/2024-02/Coroners%20Court%202023%20Annual%20Suicide%20Data%20Report%20-%20December%202023.pdf), 16 February 2024.

CCOV (2024b) [Media Release](https://www.coronerscourt.vic.gov.au/sites/default/files/2024-02/New%20report%20shows%20suicides%20continued%20to%20rise%20in%20Victoria%20in%202023.pdf) (https://www.coronerscourt.vic.gov.au/sites/default/files/2024-02/New%20report%20shows%20suicides%20continued%20to%20rise%20in%20Victoria%20in%202023.pdf), 16 February 2024.

CCOV (2024c) [Monthly Suicide Data Report, March 2024 update](https://www.coronerscourt.vic.gov.au/sites/default/files/2024-04/CCOV%20Monthly%20Suicide%20Data%20Report%20-%20March%202024%20Update.pdf) (https://www.coronerscourt.vic.gov.au/sites/default/files/2024-04/CCOV%20Monthly%20Suicide%20Data%20Report%20-%20March%202024%20Update.pdf), 11 April 2024.

CCOV (2024d), [Suicides of Aboriginal and Torres Strait Islander people](https://www.coronerscourt.vic.gov.au/sites/default/files/2024-03/Suicides%20of%20Aboriginal%20and%20Torres%20Strait%20Islander%20people%20in%20Victoria%2C%202018%E2%80%932023.pdf) (https://www.coronerscourt.vic.gov.au/sites/default/files/2024-03/Suicides%20of%20Aboriginal%20and%20Torres%20Strait%20Islander%20people%20in%20Victoria%2C%202018%E2%80%932023.pdf), 7 March 2024.

Leske S, Adam G, Catakovic A, Weir B, & Kôlves K (2022)  [Suicide in Queensland: Annual Report 2022](https://www.griffith.edu.au/_data/assets/pdf_file/0033/1639473/AISRAP-Annual-Report-2022.pdf) (https://www.griffith.edu.au/\_data/assets/pdf\_file/0033/1639473/AISRAP-Annual-Report-2022.pdf), Australian Institute for Suicide Research and Prevention, School of Applied Psychology, Griffith University, September 2022.

NSW Health (2024) [NSW Suicide Monitoring System. Report 41. Data to January 2024](https://www.health.nsw.gov.au/mentalhealth/resources/Publications/sums-report-jan-2024.pdf) (https://www.health.nsw.gov.au/mentalhealth/resources/Publications/sums-report-jan-2024.pdf), 28 March 2024.

Queensland Mental Health Commission (QMHC) (2024) [Queensland Suicide data - Monthly Report - December 2023](https://6232990.fs1.hubspotusercontent-na1.net/hubfs/6232990/Monthly%20reporting%20template%20as%20at%20December%202023.pdf) (https://6232990.fs1.hubspotusercontent-na1.net/hubfs/6232990/Monthly%20reporting%20template%20as%20at%20December%202023.pdf), 13 February 2024.

## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on [13 11 14](tel:131114), or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)

## Ambulance attendances: suicidal and self-harm behaviours

The complete extent of non-fatal suicidal and self-harming behaviours in the community is unknown in Australia. This is because, in part, not all people seek medical treatment or support for suicide attempts and self-harm injuries. Further, there are also limitations to and gaps in our current data that capture information about these behaviours. For example, data on hospitalisations due to intentional self-harm under-report the true incidence of these behaviours in the community, as only those with serious physical or mental ill-health are admitted to hospital for further treatment. Additionally, it is not possible to identify those presenting to Emergency Departments with intentional self-harm and suicide ideation in national wide data assets, although some states and territories have developed methodologies to do so within their individual datasets.

Clinical data from ambulance attendances have the potential to help broaden understandings of suicidal and self-harm behaviours in Australia, and to identify opportunities for improved intervention or postvention.

The National Ambulance Surveillance System (NASS) is a world-first public health monitoring system providing timely and comprehensive data on ambulance attendances in Australia. The NASS is a partnership between Turning Point, Monash University and state or territory ambulance services across Australia. The NASS collates and codes monthly ambulance attendances data for participating states and territories for self-harm behaviours (suicidal ideation, suicide attempt, death by suicide, and intentional self-injury).

---

## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on [13 11 14](tel:131114), or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)

## Ambulance attendances: Suicidal ideation, and suicidal and self-harm behaviours

AIHW began receiving monthly ambulance attendance data for New South Wales (NSW), Victoria (Vic), Queensland (Qld), Tasmania (Tas) and the Australian Capital Territory (ACT) from January 2021. Monthly data have been received until December 2022 for Tas, Qld, and Vic, and until September 2022 for NSW and ACT. Prior to this, data included 1-month per quarter snapshots from Vic, Tas, ACT and NSW from March 2018 to December 2020, and Qld from March 2020 to December 2020.

See [Data development activities](#) to learn more about the ongoing developments relating to ambulance attendance data funded through this project.

Self-harm related ambulance attendances are included if self-harm occurred in the preceding 24 hours or during the ambulance attendance, with 4 categories of self-harm related ambulance attendances defined and coded as:

- self-injury (non-fatal intentional injury without suicidal intent)
- suicidal ideation (thinking about taking one's own life without acting on the thoughts)
- suicide attempt (non-fatal intentional injury with suicidal intent, regardless of likelihood of lethality)
- suicide (fatal intentional injury with suicidal intent).

Suicide, suicide attempt and suicidal ideation are coded as mutually exclusive in these data; however, self-injury could be simultaneously coded with any other self-harm case category.

The number of attendances related to suicide is under-represented as ambulances do not attend all attempts, injuries, or deaths. Furthermore, when they do attend there may be insufficient information to determine suicidal intent at the scene. Rates of death by suicide have not been calculated because of small numbers, which may affect the reliability of the estimates.

Note that minor industrial action occurred in NSW on 7th April 2022, with a minimal impact on ambulance services and demand. This should be considered in any interpretation of data presented for April 2022.

For more information, see [Data sources - National Ambulance Surveillance System \(NASS\)](#).

## State and territory variations

It is important to interpret ambulance attendance rates carefully when comparing across states and territories, as there are several factors that can influence differences. For instance, the number of ambulance services available may vary by state and territory. Some states offer free ambulance services to their residents, while others are covered by private health insurance or out-of-pocket costs or are only free to vulnerable populations (Queensland Government 2020; Tasmanian Government Department of Health 2021; NSW Ambulance n.d.; ACT Emergency Services Agency n.d. & Victorian Government Department of Health 2022). Furthermore, paramedics record information that is relevant to patient care, rather than for research purposes. Data collected by paramedics may not be consistent between state and territories (Lubman et al. 2020). Although, Turning Point (the data custodian) obtains a degree of consistency between states and territories through their coding practices. Therefore, factors such as these are unmeasurable in these data and may influence some variation in the rate of attendances across states and territories.

## 2021

Throughout 2021, across NSW, Vic, Qld, Tas and the ACT, ambulances attended around 90,100 incidents, or an average around 9,000 incidents each month, that involved suicidal thoughts and behaviours (suicidal ideation or suicide attempt).

During 2021, the average monthly rates of ambulance attendances (per 100,000 population) for either suicidal ideation or suicide attempt were:

- 34 in NSW (the lowest monthly rate was 29 and the highest was 40)
- 31 in Vic (the lowest monthly rate was 26 and the highest 36)
- 50 in Qld (the lowest monthly rate was 43 and the highest 55)
- 28 in Tas (the lowest monthly rate was 24 and the highest 34)
- 38 in the ACT (the lowest monthly rate was 31 and the highest 47)

## 2022

Ambulance attendance data for Qld, Tas and Vic are available up till December 2022 with data for NSW and ACT available up till September 2022.

These data show that the average monthly rates of ambulance attendance per 100,000 population for suicidal ideation or suicide attempts combined were:

- 27 in NSW (the lowest monthly rate was 23 and the highest 32)
- 25 in Victoria (the lowest monthly rate was 22 and the highest 28)
- 47 in Qld (the lowest monthly rate was 38 and the highest 54)
- 36 in Tas (the lowest monthly rate was 31 and the highest 40)
- 29 in ACT (the lowest monthly rate was 22 and the highest 37)

Generally, for the available states and territories across both 2021 and 2022, average monthly rates of ambulance attendances per 100,000 population is highest for suicidal ideation, followed by suicide attempts, and lowest for self-injury. However, for Tas and the ACT, there are individual months where the rate of suicide attempts is higher than the rate of suicidal ideation.

The interactive data visualisation shows the average monthly rate of ambulance attendances (averaged across 12 months of a calendar year) for suicidal ideation and self-harm behaviours (self-injury and suicide attempt). These categories are subdivided by state and territory. Selection for years where monthly ambulance attendance data are also available (from 2021 onwards).

### Average monthly rate of ambulance attendances for suicidal ideation and self-harm behaviours by state or territory, 2021

Average monthly crude rate (per 100,000)



Select year:

- 2021  
 2022

Select attendance type:

Suicidal Ideation

Notes:

1. Averages are calculated by averaging the sum of monthly rates by calendar year.
2. Data included for NSW, Vic, Qld, Tas and ACT. However, NSW June 2021 data are not available and are not included in totals for 2021.

Supplementary Table: AMB S7

Source: Monash University, Turning Point

See notes ►

## Gender variations

There are distinct differences between males and females when examining deaths by suicide and intentional self-harm hospitalisations; higher rates of deaths by suicide are seen in males compared with female (see [Deaths by suicide over time \(https://www.aihw.gov.au/suicide-self-harm-monitoring/data/deaths-by-suicide-in-australia/suicide-deaths-over-time\)](https://www.aihw.gov.au/suicide-self-harm-monitoring/data/deaths-by-suicide-in-australia/suicide-deaths-over-time)) while females have higher rates of hospitalisations for intentional self-harm


(see [Intentional self-harm hospitalisations](https://www.aihw.gov.au/suicide-self-harm-monitoring/data/intentional-self-harm-hospitalisations) (https://www.aihw.gov.au/suicide-self-harm-monitoring/data/intentional-self-harm-hospitalisations)). Ambulance attendances provide further context to these gender differences. Ambulance attendances capture if the intent of the self-harm was suicidal and therefore can provide information on the extent of these behaviours in the community.

While more males die by suicide compared to females in Australia, females tend to self-harm or attempt at higher rates than males. Overall, across the 5 available states and territories in 2021, the rate of ambulance attendance with self-injury and for suicide attempts is higher for females compared to males. Attendance rates for suicidal ideation are higher in females than males in ACT, Qld and Vic. However, in NSW and Tas, attendances for suicidal ideation for males and females are similar (this can be more clearly observed when viewing the visualisation with the error bars shown).

The interactive data visualisation shows ambulance attendances for suicidal ideation and self-harm behaviours (self-injury and suicide attempt) for 2021. These categories are subdivided by sex (male and female) to show gender variations. Crude rate (per 100,000) or number of attendances can be selected, with an option to show or hide error bars. Selection for different state or territories (ACT, NSW, QLD, TAS, Vic) are also available for viewing.

### Ambulance attendances for suicidal ideation and self-harm behaviours by gender, 2021

New South Wales  
Crude rate (per 100,000)



Select measure:  
 Crude rate (per 100,000)  
 Number

Select state/territory:  
 New South Wales  
 Victoria  
 Queensland  
 Tasmania  
 Australian Capital Territory

Show uncertainty:  
 Hide error bars  
 Show error bars

#### Notes:

1. Data included for NSW, Vic, Qld, Tas and ACT. However, NSW June 2021 data are not available and are not included in totals for 2021.

Supplementary Table: AMB S1

Source: Monash University, Turning Point

[See notes ►](#)

Overall, the largest difference in ambulance attendance rates for males and females are for suicide attempts. During 2021, the annual ambulance attendance rates for suicide attempts per 100,000 population for females were:

- 133 compared to 78 for males, in NSW
- 194 compared to 97 for males, in Vic
- 267 compared to 156 for males, in Qld
- 195 compared to 107 for males, in Tas
- 234 compared to 111 for males, in the ACT.

### Age and gender variations

The interactive data visualisation below illustrates the distribution of self-harm related ambulance attendances separately for males and females by 5-year age groups. For this visualisation, ambulance attendance data for 2021 in NSW, Vic, Qld, Tas and the ACT have been combined.

In general, there were higher numbers of attendances for self-harm behaviours in the younger age groups for both males and females. Attendance numbers generally decreased with increasing age.


During 2021:

- the number of attendances for self-injury were highest for the 15-19 years age group for both males and females (around 1,300 and 3,900 attendances and crude rates of 204 and 671 per 100,000 population, respectively), however attendances for females aged 15-19 years was approximately 3 times the number of attendances for males of the same age
- attendances for suicidal ideation were highest for females in the 15-19 age group (with around 5,900 attendances and a crude rate of 1,010 per 100,000 population)

- attendance for suicidal ideation were highest for males in the 15-19 age group and 20-24 age group, with almost equally high numbers of attendances (around 3,300 attendances each and crude rates 537 and 488 per 100,000 population, respectively)
- attendances for suicide attempts were highest in the 15-19 years age group for females (around 4,800 attendances and a crude rate of 815 per 100,000 population) and in the 20-24 years age group for males (around 1,600 attendances and a crude rate of 229 per 100,000 population); the number of attendances for females aged 15-19 years was approximately three times the number for males of the same age.

The interactive data visualisation shows the distribution of ambulance attendances for suicidal ideation and self-harm behaviours (self-injury and suicide attempt) for 2021. The data is divided according to age (5-year age groups) and gender (male and female) to highlight variations. Crude rate (per 100,000) or number of attendances and attendance type can be selected, with an option to show or hide error bars.

### Ambulance attendances for suicidal ideation and self-harm behaviours by age and gender, 2021



Notes:  
 1. Data included for NSW, Vic, Qld, Tas and ACT. However, NSW June 2021 data are not available and are not included in totals for 2021.  
 Supplementary Table: AMB S2  
 Source: Monash University, Turning Point

[See notes ►](#)

### Ambulance attendances for suicidal ideation, and suicidal and self-harm behaviours over time

Trends in suicidal and self-harm behaviours are a matter of public and policy interest. However, interpretation of trends and changes in rates is complicated by large variations due, in part, to small numbers which produce large confidence intervals.

The following time series visualisations contain monthly data from January 2021 until December 2022 for Vic, Qld and Tas and until September 2022 for NSW and ACT. Data prior to 2021 are based on 1-month per quarter snapshots between March 2018 and December 2020 from NSW, Vic, Tas and the ACT, and between March 2020 and December 2020 for Qld.

Caution is advised when making month to month comparisons, particularly for the 1-month per quarter snapshot data (pre-2021 data). It is advised to compare the same months over a few years to allow for any seasonal effects and variations at different times of year. When comparing changes to estimates over time it is advised to 'Show error bars' on the visualisation. These show the 95% confidence interval for the crude rate which can vary widely in the case of small populations. This means that we are 95% confident that the true number falls within the interval range.

Over the time series in the selected state and territories (see above box for details), the general trend for the crude rate of ambulance attendances:


- increased in NSW for suicidal ideation and suicide attempts from March 2018 to Jan 2021, before decreasing over 2021 and until June 2022. Rates increased, particularly for suicidal ideation, from July to September 2022. However, the rates in September 2022 were similar to the rates in September 2018. Self-injury attendances in NSW followed similar trends to suicide attempt across the time series but with less variation in rates.
- decreased overall in Vic for suicidal ideation from December 2020 until December 2022.
- decreased for suicide attempts in Vic from December 2020 to April 2022, before increasing to just below December 2020 rates. Self-injury attendance rates slightly increased over the time series until August 2021 before steadily declining back to March 2018 levels.

- showed no clear direction in Qld for suicidal ideation and suicide attempt, with rates remaining similar across the time series. Rates of suicidal ideation increased between July and November 2022 before decreasing again in December 2022 to a similar rate to December 2020 and 2021.
- increased, in Tas for self-injury, suicidal ideation and suicide attempts from September 2021 until September 2022, noting that rates vary greatly between months due to smaller numbers.
- decreased overall in the ACT for self-injury and suicidal ideation. Suicide attempts slightly increased overall, noting that rates vary greatly between months due to smaller numbers.

The interactive timeseries visualisation shows ambulance attendances for suicidal ideation and self-harm behaviours. Crude rate or frequency can be chosen, including which state or territory. Quarterly (2018-2020) and monthly data (2021 onwards) is provided, with an option to show error bars. The categories included are self-injury, suicidal ideation, and suicide attempt.

### Ambulance attendances for suicidal ideation and self-harm behaviours

New South Wales  
Crude Rate (per 100,000)



Select data from:  
Jan 2021 to Dec 2022

Select measure:  
 Crude Rate (per 100,000)  
 Number

Select State/Territory:  
New South Wales

Self-injury  
 Suicidal ideation  
 Suicide attempt  
 Hide error bars  
 Show error bars

**Notes:**

1. Suicidal ideation and suicide attempt are mutually exclusive, while self injury is not.
  2. Quarterly snapshot data available for 2018 to 2020. Monthly data available from 2021 onwards.
  3. Data for NSW in June 2021 and NSW and ACT from Oct-Dec 2022 are not currently available.
- Supplementary Table: AMB S3.1 and AMB S3.2  
Source: Monash University, Turning Point

[See notes ►](#)

### Patterns by gender over time

Over the time series in the selected state and territories, the general pattern for the rate of ambulance attendances for:

- self-injury tended to be higher for females compared to males, across NSW, Vic, Qld and ACT
- suicidal ideation were similar for females and males across NSW, Vic, Qld, Tas and ACT
- suicide attempts were higher in females than males in NSW, Vic, and Qld.


From around March to April 2022 in Tas and ACT, suicide attempts attendance rates were distinctly higher in females than males. Prior to this period, no clear difference in attendance rates were observed (which can be more clearly seen when viewing the visualisation with the error bars shown).

The interactive timeseries visualisation shows ambulance attendances for suicidal ideation and self-harm behaviours categorised by gender (females and males) to show patterns over time. Crude rate or frequency can be chosen, with an option to show or hide error bars. Selection for different state or territories and type of suicide behaviour are also available to view.



## Ambulance attendances for suicidal ideation and self-harm behaviours by gender

New South Wales  
Crude Rate (per 100,000)



Select data from:

Jan 2021 to Dec 2022  
and Null values

Select measure:

Crude Rate (per 100,000)  
 Number

Select state/territory:

New South Wales

Select attendance type:

Suicidal ideation

Females

Males

Hide error bars

Show error bars

Notes:

1. Due to small cell numbers, self-injury attendances in Tasmania have been grouped to 'Persons'.
  2. Persons is inclusive of all recorded gender categories (including indeterminate and not stated).
  3. Data for NSW in June 2021 are not currently available.
  4. Data from Oct-Dec 2022 includes Qld, Vic and Tas only
- Supplementary Table: AMB S4.1 and AMB S4.2  
Source: Monash University, Turning Point

[See notes ►](#)

## Patterns by age and gender over time

There is a distinct variation in ambulance attendances for suicidal and self-harm behaviours between age groups. From June 2018 to September 2022, in NSW, Vic, Tas and ACT combined, attendance rates for:


- female self-injury, suicidal ideation, and suicide attempts generally decreased as age increased
- male suicide attempts were highest for the 25-44 years age group
- female self-injury in those aged under 25 years increased between March 2018 and August 2021, followed by an overall decline from November 2021 to September 2022
- female suicide attempts attendances in those aged under 25 years increased between March 2018 and March 2021 before steadily declining until September 2022
- male suicidal ideation decreased from January 2021 onwards for all age groups, particularly those under 65 years of age.
- self-injury were higher in females aged under 24 years compared to males of the same age
- self-injury for those aged 45 years and older were similar across females and males
- suicidal ideation were higher for females compared to males for those aged under 24 years, but generally higher in males compared to females for the 25-44 years age group
- suicide attempts were higher for females compared to males for the under 24 years age group.

Qld data were received from March 2020 onwards. Therefore, to retain the fullest timeseries possible (June 2018-September 2022), Qld data have been excluded from this analysis of patterns by age and gender over time.

The interactive timeseries visualisation shows ambulance attendances for suicidal ideation and self-harm behaviours categorised by age (0 to over 65) and gender (females and males) to show patterns over time. Crude rate or frequency can be chosen, with an option to show or hide error bars. Selection for different state or territories (NSW, Vic, TAS and ACT) and type of suicide behaviour are also available to view.

## Ambulance attendances for suicidal ideation and self-harm behaviours by age and gender

Crude Rate (per 100,000)



Select data from:  
Jan 2021 to Sep 2022

Select measure:  
 Crude Rate (per 100,000)  
 Number

Select attendance type:  
 Suicidal ideation

0-24     45-64  
 25-44     65+

Hide error bars  
 Show error bars

### Notes:

- Selected states and territories include: NSW, Vic, Tas and ACT. QLD is not included in this analysis due to incomplete time series. Once all selected states and territories have available data for October- December 2022, this visualisation will be updated.
  - Suicidal ideation and suicide attempt are mutually exclusive, while self-injury is not.
  - Self-injury attendances 45+ have been suppressed due to small numbers of attendances.
  - June 2021 data are currently unavailable.
- Supplementary Table: AMB S5.1 and AMB S5.2.

[See notes ►](#)

## Ambulance attendances for suicide death and suicide attempts, by modality

Monitoring the modality used in a person's death by suicide or suicide attempts can play an important role in prevention of similar events in the future. These data are provided to inform discussion around restriction of access to means of self-injury as a policy intervention for the prevention of suicide and self-harm.

Please consider your need to read the following information. If this material raises concerns for you or if you need immediate assistance, please contact a [crisis support service](https://www.aihw.gov.au/suicide-self-harm-monitoring/research-information/crisis-support) (<https://www.aihw.gov.au/suicide-self-harm-monitoring/research-information/crisis-support>), available free of charge, 24 hours a day, 7 days a week.

Please consider the [Mindframe guidelines](https://mindframe.org.au/suicide/communicating-about-suicide/mindframe-guidelines) (<https://mindframe.org.au/suicide/communicating-about-suicide/mindframe-guidelines>) if reporting on these statistics.

For each month between January 2021 and September 2022 in NSW, Vic, Qld, Tas and ACT combined, the percentage of ambulance attendances for:

- suicide attempts were highest for attempts by alcohol and other drug (AOD) and lowest for hanging presentations
- suicide deaths were highest for deaths by hanging and lowest for deaths by AOD
- suicide attempts by AOD, hanging, and other modalities showed no particular variation over time
- suicide death by hanging, AOD, and other modalities show some month-to-month fluctuations due to small numbers, but no clear change in trend.

The timeseries visualisation shows the proportion of ambulance attendances for suicide attempt and suicide death from 2021, categorised by modality (AOD, hanging and other).

---

### Content warning:

The data in this visualisation might be distressing to some readers as it contains data on the modality of suicide deaths and attempts. Please consider your need to read the following information. If this material raises concerns for you or if you need immediate assistance, please contact a [crisis support service](#), available free of charge, 24 hours a day, 7 days a week.

Please consider the [Mindframe guidelines](#) if reporting on these statistics.

[Proceed to visualisation](#)

## References

- Queensland Government (2020) *Interstate Ambulance Treatment and Transport Information for Queensland Residents* (<https://www.qld.gov.au/emergency/emergencies-services/interstate-ambulance-treatment>), Queensland Government website, accessed 25 May 2023.
- Tasmanian Government Department of Health (2021) *Ambulance costs* (<https://www.health.tas.gov.au/hospitals/ambulance/ambulance-costs>), Tasmanian Government Department of Health website, accessed 25 May 2023.
- NSW Ambulance (n.d.) *Accounts & Fees* (<https://www.ambulance.nsw.gov.au/our-services/accounts-and-fees>), NSW Ambulance website, accessed 25 May 2023.
- ACT Emergency Services Agency (n.d.) *Fees and charges* (<https://esa.act.gov.au/about-esa-emergency-services/ambulance/fees-and-charges>), ACT Emergency Services Agency website, accessed 25 May 2023.
- Victorian Government Department of Health (2022) *Ambulance fees* (<https://www.health.vic.gov.au/patient-care/ambulance-fees>) Victorian Government Department of Health website, accessed 25 May 2023.
- Lubman DI, Heilbronn C, Ogeil RP, Killian JJ, Matthews S, Smith K, Bosley E, Carney RMcLaughlin K, Wilson A, Eastham M, Shipp C, Witt K, Lloyd B, and Scott D (2020) 'National ambulance surveillance system: A novel method using coded Australian ambulance clinical records to monitor self-harm and mental health-related morbidity'. *PLoS ONE*, 15:e0236344, doi:[org/10.1371/journal.pone.0236344](https://doi.org/10.1371/journal.pone.0236344) (<https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0236344>).

## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on [13 11 14](tel:131114), or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)

## Suicide & self-harm monitoring: Intentional self-harm hospitalisations

If at any point you feel worried about harming yourself while viewing this information - or if you think someone else may be in danger - please stop reading and [seek help](#).

### What is intentional self-harm?

Intentional self-harm is often defined as deliberately injuring or hurting oneself, with or without the intention of dying. Intentional self-harm comes in many forms, and affects people from different backgrounds, ages and lifestyles. The reasons for self-harm are different for each person and are often complex.

The term 'intentional self-harm' in the National Hospital Morbidity Database (NHMD) provides information on patients admitted to hospital for self-poisoning or self-injury, with or without suicidal intent - and therefore includes both suicide attempts and non-suicidal self-harming behaviours.

Most people who self-harm do not go on to end their lives - but previous self-harm is a strong risk factor for suicide. Therefore, monitoring of intentional self-harm is key to suicide prevention.

### What are the sources of data on intentional self-harm?

Understanding the scale of the problem of intentional self-harm in Australia is difficult because many cases of self-harm are unreported unless medical treatment is required.

- Only those patients admitted to hospital for intentional self-harm are currently routinely reported in national data sets.
- Presentations to hospital emergency departments relating to suicide attempts or intentional self-harm cannot be easily identified in the current national emergency department data collection.
- Data collections from general practitioners or mental health services do not routinely capture patients treated for intentional self-harm.
- Data are available from [ambulance attendance records](#) and national population surveys (see below).

### Improving self-harm data

The NHMD is the national source of hospitalisation data in Australia. Data on the patient's diagnosis, interventions and 'external cause' (including intentional self-harm) are reported to the NHMD by all states and territories using the International statistical classification of diseases and related health problems, 10th revision, Australian modification (ICD-10-AM) and the Australian Classification of Health Interventions (ACHI). The World Health Organization's Eleventh revision of the International Classification of Diseases (ICD-11) - yet to be adopted in Australia - has the capability to classify the intent of the external cause of an injury.

In recognition of the need for better data around suicide and self-harm, the AIHW is currently working with key stakeholders, including the Mental Health and Suicide Prevention Data Governance Forum and Emergency Department data custodians to develop a nationally consistent method to identify and collect data on suicide-related ED presentations.

### National survey data

A range of national surveys are conducted to provide information on intentional self-harm. These include, but are not limited to:

- The 2020-21 and 2021-22 National Study of Mental Health and Wellbeing provides lifetime prevalence estimates of mental disorders for Australians aged 16-85. The study collects information on suicidal thoughts and behaviours and self-harm (without suicidal intent).
- The Australian Child and Adolescent Survey of Mental Health and Wellbeing (<https://www.health.gov.au/resources/publications/the-mental-health-of-children-and-adolescents>) collects data on suicidal thoughts and behaviours and self-harm (without suicidal intent) for adolescents aged 12-17.

---

© Australian Institute of Health and Welfare 2024



## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on [13 11 14](tel:131114), or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)

## Intentional self-harm hospitalisations by states & territories

Hospitalisations data for patients with intentional self-harm injuries includes those with and without suicidal intent. For further information see [Technical notes](#).

### How do intentional self-harm hospitalisations vary across states and territories?

In 2021-22:

- there were close to 26,900 hospitalisations due to intentional self-harm in Australia, with the highest proportion in Queensland (31%)
- the rate of intentional self-harm hospitalisations varied between states and territories in 2021-22, with the Northern Territory reporting the highest rate (238 hospitalisations per 100,000 population), which is more than double the national rate (105 hospitalisations per 100,000 population)
- the lowest rate was recorded in New South Wales (68 hospitalisations per 100,000 population).

Reporting is based on a patient's usual residence, not where they received treatment.

Intentional self-harm hospitalisations, by states and territories, Australia, 2008-09 to 2020-22.


The line graph shows rates of intentional self-harm hospitalisations from 2008-09 to 2020-22 for each state and territory and the total for Australia. Users can also choose to view age-specific rate, numbers and proportion of hospitalisations for intentional self-harm by states and territory by sex and specific age groups.

## Intentional self-harm hospitalisations, by states and territories, Australia, 2008-09 to 2021-22

Persons

Age group: Total (All ages)

Age-specific rate (per 100,000)



Note:

- Some data for the ACT and NT are not shown for the age groups 45-64 and 65+ due to small cell sizes.
- Age-specific rates may differ to those previously reported due to re-based estimated residential population.

Source: AIHW National Hospital Morbidity Database

Supplementary Tables: NHMD S1

Latest data: 2021-22 (annual release)

See notes ►

### How have rates of intentional self-harm hospitalisations changed over time by state and territory?

Between 2008-09 to 2021-22, rates of intentional self-harm hospitalisations in Queensland, South Australia and the Northern Territory were consistently higher than that of the national rate.

- Over this period, the highest rates of hospitalisations due to intentional self-harm in Australia were generally in the Northern Territory. Rates increased nearly 1.5 times from 162 hospitalisations per 100,000 population in 2008-09 to 238 in 2021-22.

The most notable changes between 2008-09 and 2021-22 were seen in young females.

- The rate of intentional self-harm hospitalisations for Northern Territory females in the 0-24 age group more than tripled (from 98 hospitalisations per 100,000 population in 2008-09 to 345 in 2021-22).
- In Queensland the rate has more than doubled for females in this age group (158 per 100,000 population in 2008-09 to 344 in 2021-22).

For males aged 24 and below, the rates of intentional self-harm hospitalisations:

- doubled from 90 hospitalisations per 100,000 population in 2008-09 to 189 in 2021-22 in the Northern Territory
- increased from 68 in 2008-09 to 105 in 2021-22 in Queensland, while other jurisdictions remained relatively stable.

Variation in hospital admission policy and practices between states and territories may have contributed to differences in the reporting of hospitalisation data, for further information see the [data quality statement](https://meteor.aihw.gov.au/content/index.phtml/itemId/724188) (https://meteor.aihw.gov.au/content/index.phtml/itemId/724188).

- New South Wales reported an increase in the rate of hospitalisations due to intentional self-harm in 2016-17 (130 per 100,000), before decreasing from 2017-18 to 2021-22 (97 and 68 respectively).
- Queensland reported a similar pattern with an increase in the rate of intentional self-harm hospitalisations between 2008-09 and 2020-21 (133 and 182 per 100,000 hospitalisations), before decreasing in 2021-22 (161).
- Between 2011-12 and 2012-13, Victoria reported a substantial decrease in the rate of hospitalisations due to intentional self-harm from 120 to 78 hospitalisations per 100,000. This may reflect a change in Victoria's emergency department admission policy, for further information see the [data quality statement](https://meteor.aihw.gov.au/content/index.phtml/itemId/724188) (https://meteor.aihw.gov.au/content/index.phtml/itemId/724188).

## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on [13 11 14](tel:131114), or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)

## Intentional self-harm hospitalisations by age groups

Hospitalisations data for patients with intentional self-harm injuries includes those with and without suicidal intent. For further information see [Technical notes](#).

### Rates of hospitalisations for intentional self-harm are higher for females

In 2021-22:

- two thirds of people hospitalised for intentional self-harm injuries were female (67%, or over 18,000 hospitalisations)
- the rate of intentional self-harm hospitalisations was higher for females than males (139 compared with 69 per 100,000 population)
- the rate for females aged 0-14 increased from 41 per 100,000 population in 2019-20 to 72 per 100,000 population in 2021-22.

A higher rate of intentional self-harm among females is the opposite of what is seen in deaths by suicide, where rates are higher for males than for females (see [Deaths by suicide over time](#)). This may, in part, be due to differences between methods used by males and females - with males tending to use more lethal methods than females. In addition, females generally access more health services than males. See [Patterns of health service use in the last year of life among those who died by suicide](https://www.aihw.gov.au/suicide-self-harm-monitoring/data/deaths-by-suicide-in-australia/health-service-use-in-the-last-year-of-life) (<https://www.aihw.gov.au/suicide-self-harm-monitoring/data/deaths-by-suicide-in-australia/health-service-use-in-the-last-year-of-life>) for more details.


Intentional self-harm hospitalisations by age and sex, Australia, 2008-09 to 2020-22.

The bar chart shows the age-specific rates of intentional self-harm hospitalisations for males and females for specific age groups and all ages combined in 2020-21. Users can also view age-specific rate, numbers and the proportion of hospitalisations for intentional self-harm by sex for each age group and year from 2008-09 to 2020-22.



**Intentional self-harm hospitalisations, by age and sex, Australia, 2008–09 to 2021–22**

Age-specific rate (per 100,000)



Source: AIHW National Hospital Morbidity Database  
 Supplementary table: NHMD S2  
 Latest data: 2021–22 (annual)

[See notes ►](#)

**Rates of hospitalisations for intentional self-harm are higher for young people**

Between 2008-09 and 2021-22, the rates of intentional self-harm hospitalisations were consistently high for young people. The highest rates in 2021-22 were recorded for:

- females aged 15-19 (637 per 100,000 population), followed by females aged 20-24 (342 per 100,000 population).

The highest rates for males also occurred in these younger age groups but rates were at least 2-fold lower than those of females. For example, in 2021-22:

- the highest rate of self-harm hospitalisations was 152 per 100,000 population for males aged 15-19, while males aged 20-24 reported 144 per 100,000 population.

During 2008-09 to 2020-21, there was a steady increase in the rates for both males and females aged 15-19, while rates in 2021-22 have declined (see [Intentional self-harm hospitalisations among young people](#)).

## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on [13 11 14](tel:131114), or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)

## Intentional self-harm hospitalisations by method

Understanding the methods used for intentional self-harm can play an important role in its prevention. These data are provided to inform discussion around restriction of access to means as a policy intervention for the prevention of suicide and self-harm.

Please consider your need to read the following information. If this material raises concerns for you or if you need immediate assistance, please contact a [crisis support service](#), available free of charge, 24 hours a day, 7 days a week.

Please consider the [Mindframe guidelines](https://mindframe.org.au/suicide/communicating-about-suicide/mindframe-guidelines) (<https://mindframe.org.au/suicide/communicating-about-suicide/mindframe-guidelines>) if reporting on these statistics.

The classification system used to code hospital admissions data, ICD-10-AM, uses the term 'mechanism' to refer to the external cause of a self-inflicted injury. Throughout *Suicide & self-harm monitoring* 'mechanism' has been used in data visualisations, while the term 'method' has been used in the accompanying text.

Hospitalisations data for patients with intentional self-harm injuries includes those with and without suicidal intent. For further information see [Technical notes](#).

Hospitalisations for intentional self-harm, by age, sex and mechanism, Australia, 2008-09 to 2020-22.

The line graph shows the age-specific rates of intentional self-harm hospitalisations for persons of all ages from 2008-09 to 2020-22 by method of self-harm. Users can also choose to view age-specific rate, numbers, and proportion of hospitalisations for intentional self-harm by sex for each age group.

## Most intentional self-harm hospitalisations are due to poisoning by pharmaceutical drugs

Between 2008-09 and 2021-22, the 2 most common methods of self-harm resulting in hospitalisation were *intentional self-poisoning by anti-epileptic, sedative-hypnotic, antiparkinsonism and psychotropic drugs, not elsewhere classified* (benzodiazepines are included in this category) (X61) and *intentional self-poisoning by nonopioid analgesics, antipyretics and antirheumatics* (X60):

- *Intentional self-poisoning by anti-epileptic, sedative-hypnotic, antiparkinsonism and psychotropic drugs, not elsewhere classified* (X61), which was responsible for 39% of intentional self-harm hospitalisations in 2021-22.
  - In 2021-22, 7,183 females were hospitalised as a result of this method of self-harm, compared to 3,187 males. This is more than 2 times as many hospitalisations among females compared to males in 2021-22.
- *Intentional self-poisoning by nonopioid analgesics, antipyretics and antirheumatics* (X60), which was responsible for 22% of intentional self-harm hospitalisations in 2021-22.
  - This category includes anti-inflammatory drugs, such as ibuprofen, antipyretics (for example, aspirin and paracetamol) and anti-rheumatics (some of which are used to treat arthritis).
  - More than 3 times as many hospitalisations were among females due to this method of self-harm in 2021-22 compared to male hospitalisations (4,756 and 1,161 hospitalisations, respectively).

*Contact with sharp objects (X78)* was another common method of self-harm resulting in hospitalisation.

- This method of self-inflicted injury accounted for 13% of all intentional self-harm hospitalisations in 2021-22, with more hospitalisations among females than males (1,974 and over 1,541 hospitalisations, respectively).

*Hanging (X70)* and *Gas (X67)* were the only methods of intentional self-harm that resulted in more male than female hospitalisations in 2021-22 (457 and 318 hospitalisations, and 100 and 26 hospitalisations, respectively).

---

© Australian Institute of Health and Welfare 2024



## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on [13 11 14](tel:131114), or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)

## Suicide & self-harm monitoring: Populations & age groups

Suicide and self-harm can affect people of all ages (except very young children), races, ethnicities, sexual orientations and occupations. However, a number of subgroups are particularly important to examine in depth because their risk of suicide or self-harm is higher than that of other populations, the impact on the community is different or they have specific requirements for culturally appropriate suicide prevention or postvention services.

- Although deaths by suicide occur more often in older age groups, it is the leading cause of death in Australian children and adolescents. Deaths by suicide at any age have profound effects on the families, friends and communities of those that die, but arguably, these effects are even greater when the person is young (see [Suicide among young people](#)).
- Similarly to employment in general, serving in the Australian Defence Force (ADF) seems to be protective against suicide as rates in both serving and reserve men are lower than that of all Australian men. However, for ex-servicemen suicide rates are higher than the general population (see [Australian Defence Force suicide monitoring](#)).
- The suicide rate in Aboriginal and Torres Strait Islander peoples is twice that of the non-Indigenous population (see [Suicide & Indigenous Australians](#))—although rates vary by community, age group and sex. The high rates experienced by Indigenous Australians are due to multiple, complex and interrelated social, cultural and historical influences, including colonisation, relocation of people to missions and reserves, transgenerational grief and trauma resulting from the removal of children, racism and continued socioeconomic disadvantage. However, it is important to acknowledge that Indigenous Australians may never experience suicidal behaviours or thoughts and aspects unique to their culture can be important protective factors against suicidal or self-harming behaviours.

Understanding differences in numbers and rates of suicide, intentional self-harm and suicidal behaviours in these populations is essential for more effective suicide prevention.

Other population groups identified as priority populations for suicide prevention in Australia include lesbian, gay, bisexual, transgender or intersex (LGBTI) populations and culturally and linguistically diverse (CALD) communities. It is currently not possible to discern these groups in the available suicide and intentional self-harm data sets; however, through the National Suicide and Self-harm Monitoring Project the AIHW is looking to expand data collection on these, and other population groups (see [About](#) for information on the project).

---

## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on [13 11 14](tel:131114), or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)

## Deaths by suicide among young people

Suicide is the leading cause of death among Australians aged 15-24 (see [Deaths in Australia](#)). The proportion of all deaths that are deaths by suicide is relatively high among children and young people. This is because people in these age groups do not tend to die from other causes. Data are presented by year of registration, which is not necessarily the same as the year the death occurred. For more information, see [Technical notes](#).

In 2022:


- 304 Australian young people (aged 18-24 years) took their own lives
- 77 deaths by suicide occurred among children and adolescents (aged 17 and below) with the majority occurring in those aged 15-17 (83.1%)
- deaths by suicide represented 30.9% of all deaths in young people aged 15-17 years and 32.4% of all deaths in those aged 18-24 years—up from 16.5% and 23.9% respectively of all deaths in these age groups in 2001. In children aged 14 and below, the proportion of all deaths that are deaths by suicide is low compared with the two older age groups. In 2022 deaths by suicide represented 0.9% of all deaths in children aged 14 and below.

Suicide deaths of children and young people, Australia, 2001 to 2022.

The line graph shows the age-specific rates of suicide for children and young people aged 14 and below, 15–17 and 18–24 from 2001 to 2022. Users can also choose to view the number of deaths by suicide and deaths by suicide as a proportion of all causes of death for each age group over the period.

## Suicide deaths of children and young people, Australia, 2001 to 2022

Age-specific rate (per 100,000)



Source: AIHW National Mortality Database and ABS Causes of Death, Australia 2023  
 Supplementary table: NMD S6a  
 Latest data: 2022 (annual release)

[See notes ►](#)

### Throughout 2001 to 2022:

- rates of death by suicide were highest among young adults aged 18-24 years (13.6 per 100,000 population in 2022) compared to both adolescents aged 15-17 years (6.9 in 2022), and children aged 14 years and below (0.3 in 2022).
- rates of death by suicide among 18-24 year olds decreased between 2001 and 2009 (15.9 per 100,000 population to 10.3). Then, overall, suicide rates for this age group increased until 2020 (16.7). After which, there was a decrease to 2022 (13.6). However, it is important to note that 2021 and 2022 data are preliminary and subject to change through the revisions process.
- rates of death by suicide among 15-17 year olds ranged between a low of 3.2 per 100,000 population in 2004 and a high of 8.9 in 2021.

## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on [13 11 14](tel:131114), or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)

## Intentional self-harm hospitalisations among young people

Hospitalisations data for patients with intentional self-harm injuries includes those with and without suicidal intent. For further information see the [Technical notes](#).


The data presented here are for children and young people aged below 0 and 24 years, grouped into 3 age ranges: 14 years and below, 15-19 years and 20-24 years. For children, especially those aged under 10 years, it is difficult to determine whether a self-inflicted injury was done with intent to self-harm.

Intentional self-harm hospitalisations in young people, 2008-09 to 2021-22.

The line graph shows age-specific rates of intentional self-harm hospitalisations for young people aged 14 and below, 15-19 and 20-24 from 2008-09 to 2021-22. Users can also choose to view age-specific rate, numbers and proportion of hospitalisations for intentional self-harm by sex for each age group.

## Intentional self-harm hospitalisations in young people, Australia, 2008-09 to 2021-22

Age-specific rate (per 100,000)  
Persons



Note: Age filter applies to proportion only; Sex filter applies to age specific rate and number only.

Source: AIHW National Morbidity Database  
Supplementary Table: NHMD 54  
Latest data: 2021-22 (annual release)

[See notes ►](#)

### Young people have the highest rates of hospitalisation for intentional self-harm

In 2021-22:

- the rate for young people aged 15-19 was 389 hospitalisations per 100,000 population, the highest of all age groups (including adults, not in this visualisation)
- the age-specific hospitalisation rate due to intentional self-harm was slightly lower among people aged 20-24 years (242 per 100,000), and the lowest was for children aged 14 and below (40 per 100,000 population)
- the rate for females aged 0-14 increased from 41 hospitalisations per 100,000 population in 2019-20 to 72 in 2021-22
- the age and sex-specific rate was highest for females aged 15-19 (637 hospitalisations per 100,000 population), followed by females aged 20-24 (342 per 100,000 population)
- rates for young males were lower compared to females across all ages. The lowest rate for males aged 0-14 was 9.1 hospitalisations per 100,000 population, followed by 144 among males aged 20-24 years. Similar to females, males aged 15-19 had the highest rate among young males at 152 hospitalisations per 100,000 population.

### Rates of intentional self-harm hospitalisations for girls and young females are rising

From 2008-09 to 2021-22:

- there has been a greater than 3-fold increase to the rate of intentional self-harm hospitalisations in females aged 14 and below (from 19 hospitalisations per 100,000 population to 72)
- the rate of intentional self-harm hospitalisations in females aged 15-19 has risen from 374 hospitalisations per 100,000 population to 637, and the rate for females aged 20-24 has increased from 295 to 342 hospitalisations per 100,000 population
- rates of intentional self-harm hospitalisations for males have also changed over this period but not to the same extent as those of females. An increase was seen in the 15-19 age group (from 124 hospitalisations per 100,000 population to 152) while males aged 20-24 experienced a minor decrease from 2008-09 to 2021-22 (149 and 144 per 100,000 respectively)



## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on [13 11 14](tel:131114), or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)

## Deaths by suicide among First Nations people

If at any point you feel worried about harming yourself while viewing the information on this website - or if you think someone else may be in danger - please stop reading and [seek help](#).

For further information about the wellbeing, mental health and suicide prevention of Aboriginal and Torres Strait Islander (First Nations) people, see the [Indigenous Mental Health & Suicide Prevention Clearinghouse](https://www.indigenoumshspc.gov.au/) (<https://www.indigenoumshspc.gov.au/>) managed by the AIHW. This website was developed in consultation with experts in First Nations mental health and suicide prevention, practitioners and policy makers. It brings together key research to improve the evidence base on Indigenous mental health and suicide prevention.

Since 2001, age-standardised suicide rates among First Nations people have been higher than those of non-Indigenous Australians. Preliminary 2022 data from the Australian Bureau of Statistics (ABS) Causes of Death (CoD) indicate that the rate of suicide deaths is approximately two and a half times as high among First Nations people compared with non-Indigenous Australians. Reducing deaths by suicide and suicidal behaviour among First Nations people is an issue of major concern for many First Nations communities and a public health priority for all Australian governments. By understanding the factors involved in suicide deaths among First Nations people, and how they may be different to non-Indigenous Australians, prevention strategies can be better targeted to reduce suicide deaths.

Deaths by suicide are reported for New South Wales, Queensland, Western Australia, South Australia and the Northern Territory only. Data for Victoria, Tasmania and the Australian Capital Territory have been excluded (see [Technical notes](#) for further information). All totals on this page, are the sum of these selected jurisdictions only (New South Wales, Queensland, Western Australia, South Australia and the Northern Territory). No national totals are reported for deaths among First Nations people or non-Indigenous Australians, on this page.

## Suicide rates among First Nations people

In 2022:

- suicide accounted for 4.6% of all deaths among First Nations people, with males experiencing a greater proportion of suicide deaths out of all causes of deaths compared with females (6.5% and 2.5% respectively).
- almost one quarter (24.5%) of all deaths by suicide among First Nations people were of females, while males accounted for just over three-quarters (75.5%) of all deaths by suicide among First Nations people.
- First Nations males experienced 3.3 times the rate of suicide deaths compared with First Nations females (46.3 and 14.0 suicide deaths per 100,000 population, respectively).
- both First Nations males and females experienced suicide deaths at a higher rate than their non-Indigenous counterparts. The rate of death by suicide for First Nations males was 2.6 times that of non-Indigenous males. The suicide rate for First Nations females was 2.5 times that of non-Indigenous females.

The line graph shows the age-standardised rates of suicide for First Nations people and non-Indigenous people from 2001 to 2022. Users can also choose to view age-standardised rates, numbers of deaths by suicide and deaths by suicide as a proportion of all causes of death for Indigenous and non-Indigenous people by sex.

## Suicide deaths by Indigenous status and sex, selected states and territories, Australia, 2001 to 2022



In 2022 Aboriginal and Torres Strait Islander deaths are influenced by the use of information from the Medical Certificate of Cause of Death (MCCD) for the first time for deriving the indigenous status of deaths registered in New South Wales

n.p. Not published

Note: Only data from NSW, Qld, WA, SA and NT are included.

Source: AIHW National Mortality Database and ABS Causes of Death, Australia 2023

Supplementary Table: NMD S7

Latest data: 2022 (annual release)

See notes ►

Over the last decade, between 2013 and 2022, age standardised rates of suicide among First Nations people ranged between 22.4 and 29.9 deaths per 100,000 population. Understanding the complexity and interrelationship between protective and risk factors for suicide among First Nations communities is important in designing culturally appropriate and relevant suicide prevention programs and policies. This is because concepts of wellbeing among First Nations people are shaped by factors such as historical events, community values and cultural beliefs, all of which interact and protect from or increase the risk of suicide. See [Protective and risk factors for suicide among Indigenous Australians](https://www.indigenoumhspsc.gov.au/publications/protective-and-risk-factors) (<https://www.indigenoumhspsc.gov.au/publications/protective-and-risk-factors>) for more information.

Caution should be exercised when analysing trends in deaths by suicide for First Nations people due to data quality issues, including the under-identification of First Nations people in deaths data and the uncertainties in estimating and projecting the size and structure of the First Nations population over time. The data may also be impacted by the willingness of an individual to identify as Aboriginal and/or Torres Strait Islander and how this willingness may change over time due to a variety of factors. For information about these data quality issues see: [Investigating enhancements of Indigenous data in suicide-relevant data sets](https://www.indigenoumhspsc.gov.au/publications/enhancements) (<https://www.indigenoumhspsc.gov.au/publications/enhancements>). It is also important to remember that age-standardised rates based on only a small number of deaths by suicide will exhibit a large amount of variation and that increases in numbers of deaths by suicide and rates should be treated with caution as improvements in identifying Indigenous status among deaths data may (at least in part) account for the rise in case numbers and rates. Caution is also advised when making comparisons to 2022 data due to an improvement in methodology for deriving Indigenous status for deaths registered in New South Wales. The use of a secondary source for determining Indigenous status of the deceased has introduced a break in time series of data related to deaths and First Nations people (for further information please visit [here](https://www.abs.gov.au/methodologies/causes-death-australia-methodology/2022#technical-note-the-impact-of-using-two-sources-for-deriving-the-indigenous-status-of-deaths-in-nsw-in-2022) (<https://www.abs.gov.au/methodologies/causes-death-australia-methodology/2022#technical-note-the-impact-of-using-two-sources-for-deriving-the-indigenous-status-of-deaths-in-nsw-in-2022>)).

### Death by suicide across age groups among First Nations people

Suicide contributes to premature mortality in First Nations people, especially in younger age groups. Data from the National Mortality Database and the ABS CoD from 2018-2022 showed:

- The percentage of all deaths, among First Nations people, that were deaths by suicide was highest for younger age groups. For those aged 0-24 years, almost a quarter of all deaths (22.0%) were by suicide. For those aged 25-44 years, 19.2% of all deaths were by suicide. For 45-64 year olds, it was 2.3% and for those aged 65 years and older, 0.2% of deaths were by suicide.
- The rate of suicide deaths per 100,000 population among First Nations people was highest among 25-44 year olds (50.0 deaths per 100,000 population). This was followed by 45-64-year-olds (23.4), 0-24 year olds (16.0) and those aged 65 and over (9.9).
- Among First Nations people aged 0-24 and 25-44, suicide rates were more than 3 times as high (3.1 each) compared to non-Indigenous Australians.
- Non-Indigenous Australians aged 65 years and over experienced a higher suicide rate compared to First Nations people of the same age (13.2 and 9.9 deaths per 100,000 population, respectively).


The proportion of all deaths that were deaths by suicide was higher among First Nations people aged 0-24 years compared to non-Indigenous Australians aged 0-24 years (22.0% and 16.8% respectively). However, non-Indigenous Australians experienced higher proportions of deaths by suicide within every other age-group compared with First Nations people.

Suicide deaths by Indigenous status and age groups, selected states and territories, 2018-2022.

This bar chart shows the death by suicide crude rates (per 100,000), number and per cent of all cause of deaths for Indigenous and non-Indigenous, by age group, from 2018-2022. Users can also choose to view by 5-year aggregates from 2001-2005 to 2018-2022.

### Suicide deaths by Indigenous status and age groups, 2018-2022

Age-specific rates (per 100,000)



In 2022 Aboriginal and Torres Strait Islander deaths are influenced by the use of information from the Medical Certificate of Cause of Death (MCCD) for the first time for deriving the indigenous status of deaths registered in New South Wales.

Note: Only data for NSW, Qld, WA, SA and NT are included.  
 Source: ABS Causes of Death, Australia 2023  
 Supplementary Table: NMD S15  
 Latest data: 2022 (annual release)

[See notes ►](#)

### Death by suicide across age groups and selected states and territories among First Nations people


Between 2018-2022, the highest suicide rate among First Nations people was in Western Australia for those aged 25-34 (87.0 suicide deaths per 100,000 population). Within the Northern Territory, the highest rate of death by suicide was also for those aged 25-34 years (46.9). Compared to other age groups, New South Wales, Queensland and South Australia had their highest rates among 35-44-year-olds (48.6, 54.9 and 43.2 respectively). The lowest suicide rate was in those aged 0-24 in New South Wales (8.5). This was similarly observed in Western Australia (17.9) and South Australia (15.0), while those aged 45 and over had the lowest rates in Queensland (16.2) and Northern Territory (22.3).

Deaths from suicide, by Indigenous status and age, selected states and territories, 2018-2022.

This bar chart shows the age-specific rates (per 100,000) for deaths from suicide, for First Nations people and non-Indigenous people by age group, from 2018-2022. Users can choose to view by 5-year aggregates from 2001-2005 to 2018-2022. Users can also choose to view by New South Wales, Queensland, Western Australia, South Australia, Northern Territory and the total of these states and territories.

## Suicide deaths by Indigenous status and age groups, 2018-2022

Age-specific rate (per 100,000)



Note: Only data for NSW, Qld, WA, SA and NT are included.

Source: ABS Causes of Death, Australia 2023

Supplementary Table: NMD S17

Latest data: 2022 (annual release)

[See notes ►](#)

### Death by suicide across sex and selected states and territories among First Nations people

Across selected states and territories between 2018-2022, First Nations people experienced the highest age-standardised rate (per 100,000 population) in Western Australia (38.1 suicide deaths per 100,000), followed by the Northern Territory (31.6) and Queensland (28.1). These were all greater than the rate summed across selected states and territories, for First Nations people (27.6 deaths per 100,000 population). A similar pattern was seen among First Nations males and females with Western Australia having the highest rate (56.3 and 20.0 respectively). Summed across selected states and territories, First Nations male suicide rates were almost three times higher (2.9) than First Nations females.

Summed across selected states and territories, the rate of death by suicide between 2018-2022, was 2.3 times greater for First Nations people compared to non-Indigenous Australians. The difference, in rates of suicide death among First Nations people and non-Indigenous Australians, was particularly stark in Western Australia. In Western Australia, the rate of death by suicide among First Nations males was 2.9 times greater than among non-Indigenous males, and First Nations females was 3.0 times that non-Indigenous females.


Please note, First Nations suicide rates in Western Australia vary between different regions. To guide government, non-government organisations and communities in preventing suicide in Western Australia, the Western Australia Mental Health Commission developed the [Western Australian Suicide Prevention Framework 2021-2025](https://www.mhc.wa.gov.au/media/2718/draft-suicide-prevention-action-plan-2021-2025.pdf) (Government of Western Australia Mental Health Commission 2020).

Deaths from suicide, by Indigenous status, sex and selected states and territories, 2018-2022.

This bar chart shows the age-specific rates (per 100,000) for death by suicide among First Nations people and non-Indigenous people, by selected states and territories, from 2018-2022. Users can choose to view by 5-year aggregates from 2001-2005 to 2018-2022. Users can also choose to view by NSW, Qld, WA, SA, NT and the total of these selected states and territories.

## Suicide deaths by Indigenous status across select states and territories, 2018–2022

Age-standardised rate (per 100,000)



In 2022 Aboriginal and Torres Strait Islander deaths are influenced by the use of information from the Medical Certificate of Cause of Death (MCCD) for the first time for deriving the indigenous status of deaths registered in New South Wales

Note: Only data for NSW, Qld, WA, SA and NT are included.

Source: ABS Causes of Death, Australia 2023

Supplementary Table: NMD S16

Latest data: 2022 (annual release)

See notes ►

### References

Australian Institute of Health and Welfare (2022) *Protective and risk factors for suicide among Indigenous Australians* (<https://www.indigenoushmhspc.gov.au/getattachment/2a25cdd8-d8a7-4373-938f-2aa51a0b4128/aihw-2022-protective-and-risk-factors.pdf?v=1260>). Catalogue number IMH 11, AIHW, Australian Government.

Government of Western Australia Mental Health Commission 2020. *Western Australian Suicide Prevention Framework 2021-2025* ([https://www.mhc.wa.gov.au/media/2718/draft-suicide-prevention-action-plan-2021\\_2025.pdf](https://www.mhc.wa.gov.au/media/2718/draft-suicide-prevention-action-plan-2021_2025.pdf)). Perth: Mental Health Commission





## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on [13 11 14](tel:131114), or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)

## Intentional self-harm hospitalisations among First Nations people

The AIHW uses 'First Nations people' to refer to Aboriginal and/or Torres Strait Islander people in this report.

Hospitalisations data for patients with intentional self-harm injuries includes those with and without suicidal intent. For further information see the [Technical notes](https://www.aihw.gov.au/suicide-self-harm-monitoring/data/technical-notes) (<https://www.aihw.gov.au/suicide-self-harm-monitoring/data/technical-notes>).


The quality of the hospital data provided for First Nations status varies between states and territories. For further information, see the [data quality statement](https://meteor.aihw.gov.au/content/index.phtml/itemId/724188) (<https://meteor.aihw.gov.au/content/index.phtml/itemId/724188>) and the [Technical notes](https://www.aihw.gov.au/suicide-self-harm-monitoring/data/technical-notes) (<https://www.aihw.gov.au/suicide-self-harm-monitoring/data/technical-notes>).

Intentional self-harm hospitalisations, by age, sex and Indigenous status, Australia, 2008-09 to 2020-22.

The line graph shows age-specific rates of hospitalisations for intentional self-harm by age, sex and persons for Indigenous Australians. Users can choose to view age-specific rate, numbers and proportion of hospitalisations for intentional self-harm by sex and Indigenous status for each age group.

## Intentional self-harm hospitalisations, by age, sex and First Nations status, Australia, 2008-09 to 2021-22

Persons | First Nations people  
Age-specific rate (per 100,000)



Source: AIHW National Hospital Morbidity Database  
Supplementary table: NHMD S5  
Latest data: 2021-22 (annual release)

See notes ►

### Hospitalisations for intentional self-harm among First Nations people

In 2021-22, the rate of intentional self-harm hospitalisations for First Nations people (326 hospitalisations per 100,000 population) was over 3 times that of non-Indigenous Australians (96 per 100,000 population).

During 2021-22:

- the highest rate of hospitalised intentional self-harm among First Nations people was in the 15-19 age group (710 hospitalisations per 100,000 population). The highest rate of hospitalised intentional self-harm among non-Indigenous Australians was also recorded in the 15-19 age group (366 hospitalisations per 100,000 population), though it was almost half that of First Nations Australians aged 15-19.
- First Nations females aged 15-19 years recorded the highest rate of intentional self-harm hospitalisations (1,127 hospitalisations per 100,000 population), followed by those aged 20-24 years (756 hospitalisations per 100,000 population).
- the highest rate of hospitalised intentional self-harm among First Nations males was in the 30-34 year old age group (486 hospitalisations per 100,000 population), followed by those aged 40-44 years (467 per 100,000 population) and 20-24 years (449 per 100,000 population).

### How have rates of intentional self-harm hospitalisations changed for First Nations people?

From 2008-09 to 2021-22:

- the overall rate of hospitalised intentional self-harm for First Nations people rose steadily (from 203 to 326 hospitalisations per 100,000 population)
- the rate of intentional self-harm hospitalisations for non-Indigenous Australians slightly increased from 114 hospitalisations per 100,000 population in 2008-09 to 127 hospitalisations per 100,000 population in 2016-17, before falling to 96 in 2021-22.

Over a similar period (2008-2021), the rate of death by suicide among First Nations people also increased (see, [Suicide & Indigenous Australians](#)).

Rates of hospitalisation for intentional self-harm increased from 2008-09 to 2021-22 for both First Nations females and First Nations males.

From 2008-09 to 2021-22:

- rates of hospitalised intentional self-harm among First Nations females increased from 235 to 415 hospitalisations per 100,000 population
- rates for First Nations males increased from 170 to 236 hospitalisations per 100,000 population
- the largest increase in rates of hospitalised intentional self-harm was among First Nations females aged 15-19 years. For this group, the rate of hospitalisations more than doubled, from 455 to 1,127 hospitalisations per 100,000 population
- rates of hospitalised intentional self-harm also increased more than 1.6 times among non-Indigenous females aged 15-19 years old, from 365 to 602 hospitalisations per 100,000 population

- rates of hospitalised intentional self-harm increased markedly among First Nations females aged 20-24 years (425 to 756 hospitalisations per 100,000 population) and First Nations females aged 25-29 years (381 to 593 hospitalisations per 100,000 population).
- 





## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on 13 11 14, or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)

## Australian Defence Force suicide monitoring

Suicide rates remain a concern in the Australian community and a priority for the Australian Government. Historically, ex-serving ADF members have faced an increased risk of suicide.

To increase understanding on the complex issue of suicide in serving and ex-serving ADF members, the AIHW provides annual updates to monitor the incidence of suicide in permanent, reserve and ex-serving ADF members (see Box 1). This work has been commissioned by the Department of Veterans' Affairs.

### Rate of suicide by service status and sex, 1997-1999 to 2019-2021

Select which data you wish to view below and hover over a data point for detailed information.

Select service category

- Permanent males
- Reserve males
- Ex-serving males
- Ex-serving females

Compare to

- Permanent males
- Reserve males
- Ex-serving males
- Ex-serving females

Show confidence interval?

- Yes
- No



Source: AIHW analysis of linked Defence historical personnel data-PMKeyS-NDI data 1985-2021.

<http://www.aihw.gov.au>

*Tool tip shows suicide rate per 100,000 population per year, confidence interval and significant difference for each 3-year period when the mouse is hovered over a data point.*

### Key findings

Serving permanent males and reserve males were less likely to die by suicide than the general Australian population. However, ex-serving males and females were more likely to die by suicide than the general Australian population.

Compared with the Australian population, suicide rates (after adjusting for age) between 1997 and 2021 were: 49% lower for male permanent ADF members; 45% lower for reserve ADF males; 26% higher for ex-serving ADF males; and 107% (or 2.07 times) higher for ex-serving ADF females. The rate of suicide for ex-serving ADF females was lower than the rate for ex-serving ADF males.

For voluntarily separated ex-serving males who died between 1997 and 2021, the rate of suicide was 21.5 per 100,000 population per year, which is similar to the Australian male population. However, the rate of suicide for involuntarily medically separated ex-serving males was much higher at 67.1.

For more information see *Serving and ex-serving Australian Defence Force members who have served since 1985: suicide monitoring 1997 to 2021*. Further information is presented by service status, age, sex, service, rank, length of service, time since separation and reason for separation. This year, this monitoring report includes a special in-focus chapter on DVA clients. The chapter provides a profile of ADF members who have had interactions with DVA from 2002 to 2021 and deaths by suicide are reported.

### **Box 1: Who is included in this report?**

**Permanent:** ADF members serving in a full-time capacity in the Royal Australian Navy (Navy), Australian Army (Army) or the Royal Australian Air Force (Air Force) on or after 1 January 1985, and serving in a permanent capacity on 31 December 2021 or on the date they died.

**Reserve:** ADF members who were in the reserve forces for the Navy, Army, or the Air Force on or after 1 January 1985, and were in the reserve forces on 31 December 2021 or when they died. Many members leaving full-time service transition to the reserves for a minimum of five years. The service status 'reserve' includes members with a wide range of relationships to the ADF. It includes personnel who have transitioned from full time service as well as both those who joined and have served solely in reserve capacity. Some reserve members may serve with enduring regular employment (active reserves), while others may not render service in any capacity (standby reserves).

**Ex-serving:** ADF members who were in the permanent or reserve services between 1 January 1985 and 31 December 2021, who subsequently transitioned from Defence.

### **Data sources**

In addition to the NMD, the Australian Defence Force (ADF) suicide monitoring analysis used the following data sources:

#### **National Death Index (NDI)**

The NDI is managed by the AIHW and contains person-level records of all deaths in Australia since 1980 obtained from the Registrars of Births, Deaths and Marriage in each state and territory. Its use is confined to data linkage studies approved by the AIHW Ethics Committee for health and medical research. NDI records are supplemented with cause of death information from the NMD. In this study, the NDI is linked with Defence payroll data to create the linked Defence payroll-NDI data set used in analysis of suicide in the ADF population.

#### **Department of Defence personnel system data**

The Department of Defence compiled a file of current and historical Defence personnel systems covering ADF members who have served since 1 January 1985. This combines PMKeyS, Core HR system, D1, CENRESPAY (for reservists), ADFPAY (for permanent members) and other historical payment systems. The Department of Defence and AIHW assessed the resulting file for completeness and duplicates. Comparisons were made with records from Department of Defence annual reports and other sources to validate the list. Data from the National Archives was also investigated for its suitability in validation, however as the majority of records are electronic files based on photos of paper records, this was not usable.

For further information see Technical Notes of *Serving and ex-serving Australian Defence Force members who have served since 1985: suicide monitoring 1997 to 2021*.

## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on [13 11 14](tel:131114), or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)

## Deaths by suicide among people who used disability services

Deaths by suicide among people who used disability services and the general population data (who were not disability service users) are sourced from the AIHW report: [Mortality patterns among people using disability support services: 1 July 2013 to 30 June 2018](#). These data include details about deaths for people who accessed disability specific support services, funded under the National Disability Agreement (NDA), from 1 July 2013 to 30 June 2018, herein referred to as 'people who used disability services'. Deaths reported are those that occurred between 1 July 2013 to 30 June 2018. It should, therefore, be noted that the data below are not representative of all deaths of people with disability, but rather those who had access to and were successful in applying for NDA funded support services over the 5-year study period. In this reporting, the general population is the Australian population aged under 65 years, less people who used disability services. For further information on the methods of this study please see the [Technical Report](https://www.aihw.gov.au/reports/disability-services/mortality-patterns-of-people-using-disability-serv/contents/technical-report) (<https://www.aihw.gov.au/reports/disability-services/mortality-patterns-of-people-using-disability-serv/contents/technical-report>). The National Disability Insurance Scheme (NDIS) has largely replaced the disability services currently provided by states and territories to people with disability under the NDA.

People living with disability are one of the Australian government's priority populations for suicide prevention due to the high rates of suicide and self-harm among those with disability (Cth of Australia, 2022). As such, analysis and visualisation of data from the above report has been incorporated into the National Suicide and Self-harm Monitoring System.

Further, we acknowledge those who have died by suicide and those who are bereaved and affected by suicide and self-harm. Suicide and self-harm are preventable. If you, or you believe somebody you know, is experiencing suicidality, please seek [help](#) (<https://www.aihw.gov.au/suicide-self-harm-monitoring/research-information/crisis-support>).

## Deaths by suicide among people who used disability services and the general population

People aged under 65 years, who used disability services between 1 July 2013 to 30 June 2018, died by suicide at a rate three times greater than the general population of the same age (34 and 11 per 100,000 population respectively). See notes in the visualisation below for more information on age groups.


## Deaths by suicide, by sex

The rate of death by suicide for all males who used disability services (38 per 100,000 population) was 1.4 times greater than females who used disability services (27 per 100,000 population). The difference in rate of death by suicide between males and females is substantially smaller among those who used disability services when compared to the general population. Among the general population, the rate of death by suicide for males (17 per 100,000 population) was 2.8 times greater than females (6 per 100,000 population).

The data visualisation compares rates of deaths by suicide between the general population and disability support service users from 2013 to 2018. It is categorised by sex (females, males, persons).

## Comparison of rates of suicide (per 100,000) among people who used disability services and the general population, by sex, 2013–2018

Rate of deaths by suicide (per 100,000)  
Disability service users



Select population:

- Disability service users  
 General population

Notes: Data presented above includes people aged under 20 years to 64 years.

Data source: Mortality patterns among people using disability support services: 1 July 2013 to June 2018.  
Supplementary table: S1

[See notes ►](#)

### Deaths by suicide, by sex and age group

For both males and females, and across each of the age groups, those who used disability services had higher rates of death by suicide compared to the general population.

The rate of deaths by suicide for females aged 20–34 years who used disability services (35 per 100,000 population) was over five times greater than the rate among females aged 20–34 in the general population (6.0 per 100,000 population). Among males aged 20–34 years, the rate of suicide (53 per 100,000) was more than double the rate among in the general population of the same age (21 per 100,000 population).

For females who used disability services, the highest rates of suicide were within the 35–49 years age group (38 per 100,000 population respectively). The rate of death by suicide among females who used disability services within the 35–49 years age group was almost five times greater than for females in the general population of the same age (8.0 per 100,000 population).


The highest rate of death by suicide, across all gender and age groups, was among men who used disability services aged 35–49 years (62 per 100,000 population). For men in the general population, the highest rate of death by suicide was also among those aged 35–49 years of age (26 per 100,000 population).

For woman aged 50–64 years of age, the rate of death by suicide among those who used disability services (28 per 100,000 population) was almost four times greater than among the general population (7.5 per 100,000 population). For men aged 50–64 years of age, the rate of death by suicide among those who used disability services (47 per 100,000 population) was approximately two times greater than among the general population (23 per 100,000 population).

The interactive data visualisation compares rates of deaths by suicide amongst the general population and those who used a disability support service from 2013 to 2018. The two population groups are divided according to age group from persons aged 20 to over 65 years of age. Sex can be selected (females, males, persons).

## Comparison of rates of deaths by suicide among people who used disability services and the general population, by sex and age group, 2013–2018

Rate of death by suicide (per 100,000)  
Persons



Notes: People aged under 20 years are not presented due to small numbers. 'All persons aged under 65' includes those aged under 20 years.  
Data source: Mortality patterns among people using disability support services: 1 July 2013 to June 2018.  
Supplementary table: S1


See notes ►

Among both disability service users and the general population, and across each of the age groups, males comprise a substantially higher percentage of all deaths by suicide compared females. However, across each of the age groups, the percentage of all deaths by suicide that are female is higher among those who used disability services compared to the general population. For example, 37% of deaths by suicide among disability service users aged 50-64 years were female. Whereas 25% of deaths by suicide among the general population aged 50-64 years were female.

The data visualisation shows the percentage of males and females who died by suicide, from 2013 to 2018. It is divided by age groups and range from people aged 20 to all persons under 65. Viewing for the general population or those who used a disability support service can be selected.

## Percentage (%) of females and males who died by suicide by age group, 2013–2018

Percentage (%) of each age group  
Disability service users



Notes: People under the age of 20 years were removed due to small numbers of deaths by suicide among people who used disability services when disaggregated by sex. The 'all persons under 65' category includes people aged under 20 years.  
Data source: Mortality patterns among people using disability support services: 1 July 2013 to June 2018.  
Supplementary table: S1

See notes ►

## Deaths by suicide as a percentage of deaths by all causes

Death by suicide (for all persons under 65 years) accounted for 5.2% of deaths by all causes among disability service users, making it the fourth leading cause of death for this population. Among the general population, death by suicide accounted for 8.9% of deaths by all causes and was the number one leading cause of death.

Particularly among the younger age groups, deaths by suicide accounted for a larger percentage of all deaths occurring within the general population as compared to those occurring within those who used disability services. Among the 20-34 years age group, deaths by suicide accounted for 31% of all deaths occurring within the general population and 15% of all deaths occurring within those who used disability services. Even so, death by suicide was the number one leading cause of death for the 20-34 years age group among both the general population and those who used disability services.

Deaths by suicide as a percentage of deaths by all causes can be viewed by hovering the mouse over the data points included within 'Comparison of rates of deaths by suicide among people who used disability services and the general population, by sex and age group, 2013-2018' and 'Comparison of rates of suicide (per 100,000) among people who used disability services and the general population, by sex, 2013-2018' visualisations above.

## Deaths by suicide by primary disability


In this reporting the concept of 'primary disability' is that the type of disability which most clearly reflects the person's experience of disability and causes them the most difficulty in everyday life. Primary disability groups reported here are those used by the *Mortality patterns among people using disability support services: 1 July 2013 to June 2018* (<https://www.aihw.gov.au/getmedia/de0fc029-4574-4e7b-899c-9818fa482966/aihw-dis-76-summary.pdf.aspx?inline=true>) report. It is important to note that people who live with disability can experience multiple types of disability at any point in time. The people whose data are reported here may have experienced different types of disability.

People using disability services whose primary disability was 'psychosocial disability' had substantially higher rates of death by suicide (101 per 100,000 population) compared to disability service users with all other primary disabilities and compared to the general population. The rate of death by suicide for disability service users with 'psychosocial disability' was approximately nine times greater than for the general population (11 per 100,000 population).

The rate of death by suicide for disability service users (for people aged under 65) with 'psychosocial disability' was 1.7 times greater than among those with 'acquired brain injury' (56 per 100,000 population) as a primary disability. The rate of death by suicide for those with 'psychosocial disability' was four times greater than among those with a 'physical disability' (25 per 100,000 population) as their primary disabilities. Among those with a primary 'psychosocial disability' the rate of death by suicide was more than six times greater than for those with hearing disability, more than nine times greater than for those with learning specific/ADD disability, and more than 25 times greater than for those with autism as their primary disability.

The data visualisation shows rates of death by suicide for those who used a disability support service, by primary disability, from 2013 to 2018. Primary disability is categorised into 6 types (Autism, specific learning/ADD, hearing disability, physical disability, acquired brain injury, psychosocial disability) and data includes people aged under 20.

**Rates of suicide (per 100,000), among people who used disability services, by primary disability, 2013-2018**



Notes: Data presented above includes people aged under 20.  
Data source: Mortality patterns among people using disability support services: 1 July 2013 to June 2018.  
Supplementary table: S1


See notes ►

For those using disability services whose primary disability was ‘psychosocial disability’, suicide deaths accounted for 20.7% of all deaths occurring within this group during the study period. Suicide deaths accounted for the highest percentage of all cause deaths for those with psychosocial disability compared to all other primary disabilities. The percentage of all deaths accounted for by suicide was second highest among those with a primary disability of autism. Suicide deaths accounted for 6.0% of all deaths among those with a primary disability of autism. The percentage of all deaths accounted for by suicide among people with psychosocial disability was around four times greater than those with autism.

The interactive data visualisation shows deaths by primary disability, population and cause of death for all persons aged under 65 between 2013 to 2018. Selection for mortality rate or number of deaths, cause of death (all cause or suicide) and population type are all selectable features.

### Deaths by primary disability, population and cause of death, all persons under 65, 2013–2018

Number of deaths



Notes: Data presented above includes people aged under 20.  
 Data source: Mortality patterns among people using disability support services: 1 July 2013 to June 2018.  
 Supplementary table: S1


[See notes ►](#)

The rate of death by suicide among people whose primary disability was psychosocial disability was highest for those aged 20-34 years (118 per 100,000 population). However, among all people who used disability services the highest rate of death by suicide was for those aged 35-49 years (51 per 100,000 population).

The data visualisation shows rates of death by suicide from 2013 to 2018, for the general population, disability service users and disability service users with a psychosocial disability. These categories are subdivided by age group, ranging from 20 to all persons under 65 years of age.

**Rates of death by suicide among disability service users with psychosocial disability, people who used disability services, and the general population, by age group, 2013–2018**

Rate of death by suicide (per 100,000)



Notes: People under the age of 20 years were removed due to small numbers of deaths by suicide among people who used disability services when disaggregated by sex. The 'all persons under 65' category includes people aged under 20 years. 'Disability service users' includes people with psychosocial disability. Disability service users were removed from the 'general population'.

Data source: Mortality patterns among people using disability support services: 1 July 2013 to June 2018. Supplementary table: S1

See notes ►

Among those whose primary disability was psychosocial disability, rates of death by suicide were highest for the 20-34 year age group (118 per 100 000 population) and lowest for the 50-64 years age group (86 per 100 000 population). Even so, the rate of death by suicide among those with psychosocial disability aged 50-64 years, was considerably higher than the peak rates of death by suicide for all disability service users (51 per 100,000 population among those aged 35-49 years) and the general population (17 per 100,000 population among those aged 35-49 years).

Rates of suicide by all those who used disability services was curved across age groups, peaking in the 35-49 year age group.

**References**

The Commonwealth of Australia (Cth of Australia) (2022) *National Mental Health and Suicide Prevention Agreement* (<https://federalfinancialrelations.gov.au/agreements/mental-health-suicide-prevention-agreement>), The Federal Financial Relations website, accessed 3 March 2023.



## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on [13 11 14](tel:131114), or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)

## LGBTIQ+ Australians: suicidal thoughts and behaviours and self-harm

If at any point you feel worried about harming yourself while viewing this information—or if you think someone else may be in danger—please stop reading and seek [help](#). You can access [LGBTIQ+ resources online](https://qlife.org.au/resources) (<https://qlife.org.au/resources>), and Qlife (trained LGBTIQ+ peer support): Telephone [1800 184 527](tel:1800184527) (3:00pm - midnight everyday 7 days a week) or by [webchat](https://qlife.org.au/resources/chat) (<https://qlife.org.au/resources/chat>).

The acronym LGBTIQ+ is used here as an umbrella term to refer to lesbian, gay, bisexual, trans/transgender, intersex, queer and other sexuality, gender and bodily diverse people and communities.

The data presented on these webpages are from the 2019 Private Lives 3 (PL3) and Writing Themselves In 4 (WTI4) surveys. Whilst these surveys included participants with an intersex variation/s, the data are not able to be disaggregated by this category and, therefore, the acronyms LGBTQ+ or LGBTQA+ are used when referring to the PL3 and WTI4 results. LGBTIQ+ is used when referring to communities more generally and different acronyms may be used throughout these pages, depending on how communities are represented within the different data sources discussed.

For more information on terminology relating to LGBTIQ+ people and communities, see the [Australian Institute of Family Studies' \(AIFS\) LGBTIQ+ glossary of common terms \(AIFS 2022\)](https://aifs.gov.au/resources/resource-sheets/lgbtqi-glossary-common-terms) (<https://aifs.gov.au/resources/resource-sheets/lgbtqi-glossary-common-terms>).

LGBTIQ+ communities have been identified as priority populations under *The National Mental Health and Suicide Prevention Agreement* (Cth of Australia, 2022) and for data development as part of the National Suicide and Self-harm Monitoring System. Under the agreement, governments have a responsibility to support priority populations, who may be at higher risk of mental ill health and suicide due to vulnerability caused by social, economic, and environmental circumstances.

### Data on suicide and self-harm among LGBTQ+ people from the Private Lives 3 and Writing Themselves In 4 surveys

The Australian Research Centre in Sex, Health and Society (ARCSHS) at La Trobe University runs Australia's two largest targeted surveys of LGBTQ+ adults and LGBTQA+ young people, the Private Lives and Writing Themselves In surveys, respectively (Hill et al. 2020, 2021). The most recent iterations of these surveys, Private Lives 3 (PL3) and Writing Themselves In 4 (WTI4) were undertaken in 2019. The PL3 and WTI4 datasets are the largest and most comprehensive available for the LGBTQ+ population in Australia and include a diverse sample of participants from all states and territories and demographic groups (Hill et al. 2020, 2021).

ARCSHS has provided the AIHW with existing data on suicide and self-harm from PL3 and WTI4, aggregated by state/territory, age-group, gender and sexual orientation. In addition, the AIHW has engaged ARCSHS to undertake secondary analysis of the data from PL3 and WTI4, including:

- Types of gender affirmation accessed by trans and gender diverse adults and association with health and wellbeing outcomes (PL3).
- Types of gender affirmation accessed by trans and gender diverse young people and association with mental health outcomes and suicidality (WTI4).
- The role of relationship status and gender of relationship partner in shaping health and wellbeing outcomes among multigender attracted (bisexual+) adults (PL3).

Data on suicidal thoughts and suicide attempt among LGBTQ+ adults from the PL3 survey and LGBTQ+ young people from the WT14 survey are presented in the following sections. This is the second tranche of data from PL3 and WT14 to be published on the AIHW Suicide and self-harm monitoring website. The results of the secondary analysis of PL3 and WT14 will be published in 2024.

The findings of PL3 and WT14 are consistent with evidence from Australia and overseas, which indicate that LGBTQ+ communities experience higher levels of mental ill health, suicidality and self-harm, compared with the general population (Hill et al. 2020, 2021, Marchi et al. 2022, Swannell et al. 2016, Zwickl et al. 2021). Within the LGBTQ+ research, trans and gender diverse participants appear to experience a greater risk of suicidal thoughts and behaviours, compared with cis-gendered participants. For instance, among PL3 participants:

- The lifetime prevalence of suicidal thoughts ranged from 64% among cisgender men to 90% among non-binary participants and 91% among trans men.
- More than half of trans men reported having attempted suicide in their lifetimes (53%), in contrast to around one-fifth of cisgender men (22%).

Other Australian studies of trans people have found that a large proportion of participants (ranging from 43 to 48%) have attempted to take their own lives at some point (Zwickl et al. 2021, Bretherton et al. 2021, Strauss et al. 2017).

A limitation of PL3, WT14 and other targeted, community surveys of LGBTQ+ people is that they tend not to be based on probability sampling and, as a result, it is not possible to conclude that they provide representative data for the LGBTQ+ population. However, these surveys do provide important information about the survey respondents, which can inform the work of LGBTQ+ researchers and advocates, and policy makers.

### What other national suicide and self-harm data are available for LGBTQ+ communities in Australia?

There are currently no reliable national data on rates of suicide and self-harm among LGBTQ+ communities in Australia. The two key administrative datasets used by the AIHW to report on rates of suicide and hospitalised self-harm, the National Mortality Database (NMD) and the National Hospital Morbidity Database (NHMD) do not include information on LGBTQ+ status. LGBTQ+ status is not available in any national linked administrative datasets and has not been enumerated in the Census of Population and Housing. Data gaps could be improved by the broader inclusion of the [Standard for Sex, Gender, Variations of Sex Characteristics and Sexual Orientation Variables, 2020](https://www.abs.gov.au/statistics/standards/standard-sex-gender-variations-sex-characteristics-and-sexual-orientation-variables/2020) (https://www.abs.gov.au/statistics/standards/standard-sex-gender-variations-sex-characteristics-and-sexual-orientation-variables/2020) (ABS 2020) in national collections.

State and territory suicide registers include variables on sexual orientation and gender, however, LGBTQ+ status tends to be underreported in these surveillance systems (CCOV 2022, Leske et al. 2022). To date, the Victorian Suicide Register (VSR) and the Queensland Suicide Register (QSR) are the only state suicide registers to publish data on suicide deaths among LGBTQ+ people. In each state, the numbers are too small to disaggregate by gender and sexual orientation (CCOV 2022, Leske et al. 2022).

The population representative, National Study of Mental Health and Wellbeing (2020-22), conducted by the Australian Bureau of Statistics (ABS) collected information on suicidality and self-harm and was the first ABS collection to use the [Standard for Sex, Gender, Variations of Sex Characteristics and Sexual Orientation Variables, 2020](https://www.abs.gov.au/statistics/standards/standard-sex-gender-variations-sex-characteristics-and-sexual-orientation-variables/2020) (https://www.abs.gov.au/statistics/standards/standard-sex-gender-variations-sex-characteristics-and-sexual-orientation-variables/2020) (ABS 2020, 2020-22b). The summary statistics for the study were published by the ABS on 5 October 2023 and include results for lived experience of suicide and self-harm by sex at birth (male/female) (ABS 2020-22a), see [Australian prevalence estimates of suicidal behaviour](#). Data were collected from people whose sex at birth was recorded as something other than male or female, however the sample size was too small to report separately on this group. The data for these people are included in the results for total persons but not in the male or female categories (ABS 2020-22b). The AIHW is investigating whether further disaggregation of the suicide and self-harm data by gender (*Cis, Trans and gender diverse*) and sexual orientation (*Heterosexual, Gay, or Lesbian, Bisexual and Different term*) is possible.

### Prevalence of suicidal behaviour and thoughts among LGBTQ+ adults - data from the 2019 Private Lives 3 (PL3) survey

We acknowledge those from the LGBTQ+ community who have died by suicide and those bereaved by suicide. You can access [LGBTQ+ resources online](https://qlife.org.au/resources) (https://qlife.org.au/resources), and Qlife (trained LGBTQ+ peer support): Telephone 1800 184 527 (3:00pm - midnight everyday 7 days a week) or by [webchat](https://qlife.org.au/resources/chat) (https://qlife.org.au/resources/chat).

PL3 is the third iteration of national surveys investigating the health and wellbeing of lesbian, gay, bisexual, trans and gender diverse and queer (LGBTQ) adults in Australia. The PL3 survey is managed by the Australian Research Centre in Sex, Health and Society (ARCSHS) at La Trobe University in Melbourne and was developed by ARCSHS in consultation with an Expert Advisory Group comprising representatives from the states and territories and LGBTQ+ groups. The sample was recruited via paid advertising on social media and promotion through relevant professional networks and LGBTQ+ organisations. The survey was open to people aged 18 years and over, from 24 July 2019 until 1 October 2019 and could be completed online or in paper form if requested. A sample of 6,835 participants was achieved, whose ages ranged from 18-88 years. The PL3 survey included questions on suicide attempt and suicidal thoughts in the past 12 months and lifetime. For more information, view the [PL3 national report](https://www.latrobe.edu.au/arcshs/work/private-lives-3) (https://www.latrobe.edu.au/arcshs/work/private-lives-3).

People with an intersex variation/s were specifically targeted for participation in PL3 but the sample achieved (n=47) was too small to provide statistically meaningful comparisons. Therefore, data for participants with an intersex variation/s are not reported as a separate group and the acronym 'LGBTQ+' is used when discussing the PL3 results below. The data for PL3 participants who reported having

intersex variation/s are included in the other categories presented, according to their responses to gender and sexual orientation questions. View the [PL3 national report](https://www.latrobe.edu.au/arcshs/work/private-lives-3) (https://www.latrobe.edu.au/arcshs/work/private-lives-3) for more information about participants with an intersex variation/s.

The PL3 results relating to suicide attempt and suicidal thoughts are depicted in the visualisations below. As PL3 uses a non-probability convenience sample, the results may not be representative of the Australian LGBTQ+ population and cannot be generalised to this population group. However, they provide valuable insights into the experiences of close to 7000 people from this population group and highlight where further work is needed to obtain better data and improve outcomes for at-risk communities. Importantly, the PL3 sample allows for disaggregation of data by gender and sexual orientation, which illustrates the wide variations in experiences of suicidal thoughts and behaviours between the different gender and sexual orientation groups in the sample. These results are consistent with other studies that show considerable variation in the prevalence of suicidal thoughts and behaviours between sub-groups under the LGBTQ+ umbrella (e.g. Kirakosian et al. 2023, Marchi et al. 2022, Stinchcombe & Hammond 2021, Swannell et al. 2016).

The results of PL3 are not directly comparable with those for the general population from national population surveys, such as the ABS National Study of Mental Health and Wellbeing (2020-21) and the ABS National Survey of Mental Health and Wellbeing (2007). The ABS surveys used probability sample designs as well as different recruitment methods, instruments, and modes of administration (see ABS 2020-21b for information on methodology). Both types of survey designs have limitations regarding sampling LGBTQ+ communities. Targeted surveys, such as PL3, may be biased towards people with stronger attachment to the LGBTQ+ community, while population surveys may underrepresent LGBTQ+ people (Hottes et al. 2016) and obtain insufficient samples to report results by gender and sexual orientation. A meta-analysis of lifetime prevalence of suicide attempt among lesbian, gay and bisexual (LGB) people by Hottes et al. (2016) found that targeted community surveys reported higher prevalence of lifetime suicide attempt among LGB people, compared with results for LGB people from population surveys. Even so, LGB people reported higher prevalence of lifetime suicide attempt compared with heterosexual people, regardless of the survey type (Hottes et al. 2016).

In relation to the PL3 data, the term 'suicidal thoughts' is being used rather than 'suicidal ideation'. This is because suicidal ideation is defined in national population data as 'serious thoughts about taking one's own life', whereas the PL3 data item is 'thoughts about suicide, wanting to die, or about ending your life'.

As the PL3 was a voluntary online survey, participants could leave questions blank if they wished. In these cases, the PL3 results reported below are the proportions (percentages) of those who answered the relevant question.

### **Suicide attempts and suicidal thoughts among PL3 participants**

The visualisation includes PL3 data for suicidal thoughts and attempts broken down by sexual orientation, gender, state and territory of residence, or age group.

#### **Suicidal thoughts and attempts among LGBTQA+ PL3 (2019) participants by personal characteristics**

The visualisation includes PL3 data for suicidal thoughts and attempts broken down by sexual orientation, gender, state and territory of residence, or age group.

Choose a variable:  Sexual orientation  Gender  Age group  State or territory


Select one:  Suicidal thoughts  Suicide attempt

Select response:  Ever  Last 12 months  Never  Prefer not to say

Label:  On  Off

Axis range:  Fixed  Automatic

**Sexual orientation | Suicidal thoughts | Ever**  
Per cent %



Notes: Results for Northern Territory (NT) should be interpreted with caution due to small numbers of respondents (n=37). n.p. - not presented  
Source: Private Lives 3 (PL3)  
Supplementary table: S1, S2, S3, S4

Nationally, around three-quarters (75%) of PL3 participants had experienced suicidal thoughts and around one third (30%) reported attempting suicide in their lifetimes.

### Results by sexual orientation

The PL3 survey asks participants to select which terms best describe their sexual orientation and then asks them to select the term they would use if they had to choose only one. The options provided were developed in consultation with the PL3 Expert Advisory Group and include: ‘lesbian’, ‘gay’, ‘homosexual’, ‘bisexual’, ‘pansexual’, ‘heterosexual’, ‘queer’, ‘asexual’, ‘prefer not to have a label’, ‘prefer not to answer’, ‘don’t know’ and ‘something different’ (with free text option to describe). In the analysis, participants who selected ‘homosexual’ (due to low numbers), ‘prefer not to have a label’ and ‘something different’ as well as trans and gender diverse participants and those with an intersex variation/s who selected ‘heterosexual’ were combined into the category ‘something else’. For more information on sexual orientation in PL3, please see the [PL3 published report by La Trobe University](https://www.latrobe.edu.au/_data/assets/pdf_file/0009/1185885/Private-Lives-3.pdf) ([https://www.latrobe.edu.au/\\_data/assets/pdf\\_file/0009/1185885/Private-Lives-3.pdf](https://www.latrobe.edu.au/_data/assets/pdf_file/0009/1185885/Private-Lives-3.pdf)).

The results for sexual orientation differed across response categories, apart from gay respondents, who were least likely to have experienced suicidal thoughts or attempt, recent (last 12 months) or in their lifetimes.

- Pansexual and queer participants reported the highest prevalence of lifetime suicidal thoughts (88% and 87%, respectively), followed by bisexual and asexual participants (79.7% and 79.2% respectively), participants categorised as “something else” (75.3%), lesbian participants (73.0%) and gay participants (63.3%).
- A similar pattern was seen for recent suicidal thoughts, reported by 54.7% of queer participants, 54.6% of pansexual participants, 49.1% of bisexual participants, 44.9% of respondents categorised as “something else”, 43.4% of asexual participants, 36.8% of lesbian participants and 30.7% of gay participants.
- Nearly half (46.7%) of pansexual participants reported having attempted suicide in their lifetimes, followed by queer participants (37.1%), “something else” (34.3%), bisexual (31.5%), lesbian (30.1%), asexual (27.0%) and gay (20.5%) participants.
- Participants in the “something else” sexual orientation category were most likely to report recent suicide attempt (9.6% of this group), followed by pansexual (7.8%), bisexual (6.0%), queer (5.1%), asexual and lesbian (4.1%) and gay (3.3%) participants.

There is limited research that includes the sexual orientations “pansexual”, “queer” and “asexual” as specific groups with which to compare these results. Studies that have included lesbian, gay and bisexual people as separate groups have generally reported higher levels of suicidal thoughts and behaviour among bisexual and multi-gender attracted people, compared with gay and lesbian people (e.g. Stinchcombe & Hammond 2020, Marchi et al. 2022).

### Results by gender

When disaggregated by gender, the results show that trans (trans man and trans woman) and non-binary participants were more likely to have experienced recent (last 12 months) and lifetime suicidal thoughts and suicide attempts, compared with cisgendered participants (cisgender man and cisgender woman).

- Lifetime prevalence of suicidal thoughts among PL3 participants ranged from 64.2% of cisgender men to 89.9% of non-binary participants and 90.6% of trans men.
- More than half of trans men reported having attempted suicide in their lifetimes (52.9%), compared with around one fifth of cisgender men (22.3%).

The high levels of suicidal thoughts and behaviour among trans participants are consistent with other studies of trans people in Australia. A 2017-2018 survey of Australian trans adults found that 43% of participants had attempted suicide in their lifetimes (Zwickl et al. 2021, Bretherton et al. 2021). The 2016 *Trans Pathways* survey of Australian trans young people aged 14-25 years reported that 48.1% of participants had ever attempted suicide (Strauss et al. 2017).

## Results by age group

The results for age group show a clear gradient by age, with younger age groups more likely to report lifetime and recent (last 12 months) experience of suicidal thoughts and lifetime suicide attempt.

- Lifetime experience of suicidal thoughts ranged from around half of people aged 65 years and over (50.7%) to 79.6% of people in the 18-24 and 25-34-year age groups.
- Lifetime suicide attempt ranged from 17.5% of people aged 65 years and over to 34.0% of 18-24-year-olds.
- Recent suicide attempt was most likely among participants aged 18-24 years (9.8%), followed by participants aged 35-44 years (4.5%), then those aged 25-34 (3.1%), 45-54 years (2.0%), 55-64 years (1.2%) and over 65 years (0%).

The decline in recent suicidal thoughts and attempts with increasing age is consistent with other studies with trans participants (Zwickl et al 2023). This pattern is also observed with the general population results from the ABS National Study of Mental Health and Wellbeing (2020-22), which show prevalence of lifetime and recent suicidal ideation and suicide attempt is highest among the youngest age group (16-34 years) and decreases with increasing age (ABS 2020-22a). [Ambulance data](#) from the National Ambulance Surveillance System (NASS) also show higher rates of attendances for suicidal ideation and attempt among younger age groups.

The PL3 results for younger people may be influenced by the greater proportions of younger people categorised as trans, gender diverse, bisexual, pansexual and queer, relative to those in the older age groups (Hill et al. 2020). Participants in each of these categories are more likely to experience poor mental health outcomes, as well as discrimination and stigma, when compared with cisgendered, gay and lesbian participants (Hill et al. 2020).

## Results by state and territory

The prevalence of suicidal thoughts and attempt among PL3 participants was similar across states and territories.

- Lifetime prevalence of suicidal thoughts ranged from 67.6% in the Northern Territory (NT) to 80.6% in the Australian Capital Territory (ACT).
- Recent (last 12 months) prevalence of suicidal thoughts ranged from 35.1% in the NT to 47.3% in Tasmania (Tas).
- Participants from Tas and Queensland (Qld) were most likely to report having attempted suicide in their lifetimes (35.5% and 34.7%, respectively), followed by those in South Australia (SA) (33.1%), Western Australia (WA) (32.3%), ACT (30.7%), Vic (28.1%), New South Wales (NSW) (28.0%) and the NT (21.4%).
- Recent suicide attempt was also more likely to be reported by participants from Tas (7.8%) and Qld (7.1%), followed by WA (6.2%) and ACT (5.7%), Vic (4.5%), SA (4.2%) and NSW (4.1%).
- Recent suicide attempt is not reported for participants from the NT due to the small number of NT participants in the survey.

Caution should be used in interpreting the results by state and territory, as they may be affected by sampling and recruitment bias, in particular for the NT, where the sample size was only 37. Relative to the general population, PL3 oversampled people from Vic and the ACT, and under-sampled people from NSW and Qld (Hill et al. 2020). There may also be confounding due to differences in the age, gender, and sexual orientation distributions of LGBTQ+ people by state and territory.

## Suicide attempt, suicidal thoughts and disability or long-term health conditions in PL3

The PL3 survey asks participants whether they have a disability or long-term health condition (defined as one that has lasted or is expected to last 6 months or longer). If participants answer “yes” to this question, they are then asked a series of questions taken from the AIHW’s Standardised Disability Flag Module (SDFM). The SDFM identifies people who may be living with disability and/or long-term health conditions and the impact these conditions have on their day-to-day living (none, mild, moderate, and severe). For further information about how the SDFM was used and the limitations it may have on the data presented please see the [PL3 national report](#) ([http://www.latrobe.edu.au/\\_data/assets/pdf\\_file/0009/1185885/Private-Lives-3.pdf](http://www.latrobe.edu.au/_data/assets/pdf_file/0009/1185885/Private-Lives-3.pdf)).

The visualisation below includes two charts related to disability or long-term health condition. The left-hand chart shows the prevalence of suicidal thoughts and attempts among PL3 participants according to their disability or long-term health condition. The right-hand chart shows suicidal thoughts and attempts among PL3 participants who reported having a disability or long-term health condition, according to the impact on their day to day living (disability severity).

### Suicidal thoughts and attempts among LGBTQA+ PL3 (2019) participants by disability and/or long-term health condition status.

The visualisation includes two charts related to disability or long-term health condition. The left-hand chart shows the prevalence of suicidal thoughts and attempts among PL3 participants according to their disability or long-term health condition. The right-hand chart shows suicidal thoughts and attempts among PL3 participants who reported having a disability or long-term health condition, according to the impact on their day to day living (disability severity).

Select one:  
 Suicidal thoughts  
 Suicide attempt


Select response:  
 Ever  
 Last 12 months  
 Never  
 Prefer not to say

Label:  
 On  
 Off

Axis range:  
 Fixed  
 Automatic

### Suicidal thoughts | Ever

Per cent %



Notes: Disability and long-term health conditions were identified through the use of the Australian Institute of Health and Welfare's Standardised Disability Flag Module. For an overview of how this may be affected the data, please see the original PL3 report.

Source: Private Lives 3 (PL3)

Supplementary table: S5, S6

### Results by disability or long-term health condition status

PL3 participants who reported they have a disability or were unsure whether they have a disability (84.8%) or were unsure whether they have a disability (84.7%) were most likely to have experienced suicidal thoughts in their lifetimes, compared with PL3 participants who did not have a disability (63.9%).

PL3 participants who reported having a disability or being unsure about having a disability also reported the highest prevalence of recent (last 12 months) suicidal thoughts (54.2% and 53.0%, respectively), compared with those without a disability (28.6%).

Suicide attempt was more common among PL3 participants who reported having a disability, with 40.4% having attempted suicide in their lifetimes. This is twice the proportion of PL3 participants without a disability (19.6%) who reported lifetime suicide attempt.

### Results by disability or long-term health condition impact on day-to-day activities

The results for lifetime and recent (last 12 months) suicidal ideation and suicide attempt show a clear gradient by the severity of activity limitation.

- Among PL3 participants with a disability or long-term health condition who reported severe limitations, 90.9% reported having suicidal thoughts in their lifetimes and 50.1% reported attempting suicide in their lifetimes. More than two thirds (68.0%) of these participants reported recent suicidal thoughts.
- For PL3 participants with a disability or long-term health condition who had no activity limitation, 71.2% reported lifetime suicidal thoughts and 26.4% reported lifetime suicide attempt.
- Lifetime and recent suicidal thoughts and attempt for PL3 participants with a disability or long-term health condition who had no activity limitation were similar to those without or unsure of whether they have a disability or long-term health condition. This suggests that activity limitation due to disability/long-term health conditions are associated with greater risk of suicidal thoughts and attempts.

### Suicide attempt, suicidal thoughts by age, sexual orientation, and gender in PL3


The visualisation below illustrates the percentage of PL3 participants, who reported experiencing suicidal thoughts and attempting suicide, categorised by age, sexual orientation, and gender, throughout their lifetime and over the last 12 months.

#### Suicidal thoughts and attempts among LGBTQA+ PL3 (2019) participants by age, sexual orientation, and gender.

This visualisation illustrates the percentages of PL3 participants, who reported experiencing suicidal thoughts and attempting suicide, categorised by age, sexual orientation, and gender, throughout their lifetime and over the last 12 months.

### Sexual orientation | Suicidal thoughts | Ever

Per cent %



Notes: Caution should be exercised when interpreting the results of asexual or trans woman participants 45 years and over, due to low numbers in the PL3 survey.

Source: Private Lives 3 (PL3)

Supplementary table:

## Results by sexual orientation and age

While the results for lifetime suicide thoughts and suicide attempts are mixed, recent (last 12 months) suicidal thoughts were found to decrease with age.

- Lesbian participants reported a decrease in the likelihood of lifetime suicidal thoughts with age from 85.5% for 18-24-year-olds to 62.7% for those aged 45 years and over. For all other sexual orientations there was no obvious trend.
- Lesbian participants reported a decrease in the likelihood of lifetime suicide attempt with age (41.5% of 18-24-year-olds, 22.0% of those aged 45 years and over). Gay and queer participants also show an overall decrease with age.
- Participants of all sexual orientations, except those who are asexual, showed a decrease in recent suicidal thoughts with age. While a similar relationship may exist for suicide attempts it is difficult to determine due to small numbers.

## Results by gender and age

- Cisgender woman participants reported that lifetime suicidal thoughts (80.1% for 18-24-year-olds, 64.1% of those aged 45 years and over) and suicide attempt (31.0% of 18-24-year-olds, 19.6% of those aged 45 years and over) decreased with age. Cisgender man and non-binary participants may also demonstrate a decrease in lifetime suicidal thoughts and suicide attempt with age.
- Cisgender man, cisgender woman and non-binary participants reported that recent suicidal thoughts decreased with age. Trans man and trans woman participants also show that recent suicidal thoughts trend downwards with age. It is difficult to analyse recent (last 12 months) suicide attempt due to low small numbers.

## Prevalence of suicidal behaviour and thoughts and self-harm among LGBTQA+ young people - data from the 2019 Writing Themselves In 4 (WTI4) survey

WTI4 is the fourth iteration of national surveys investigating the health and wellbeing of lesbian, gay, bisexual, trans and gender diverse, queer and Asexual (LGBTQA) young people, aged 14 to 21 years, in Australia. The WTI4 was conducted by the Australian Research Centre in Sex, Health and Society (ARCSHS) at La Trobe University in Melbourne in 2019. The sample was recruited via paid advertising on social media and promotion through relevant professional networks and LGBTQA+ organisations and achieved a sample of 6,418 participants. The WTI4 survey included questions on suicidal thoughts, suicide plans and attempts and self-harm in the past 12 months and lifetime. For more information, view the [WTI4 report](https://www.latrobe.edu.au/data/assets/pdf_file/0010/1198945/Writing-Themselves-In-4-National-report.pdf) (https://www.latrobe.edu.au/data/assets/pdf\_file/0010/1198945/Writing-Themselves-In-4-National-report.pdf). As with PL3, the results of WTI4 may not be representative of young LGBTQA+ Australians and cannot be compared with the results for young people in the general population from national population surveys.

The visualisation below illustrates WTI4 participants who have experienced suicidal thoughts, suicide plan, suicide attempt, or engaged in self-harm, throughout their lifetime and over the last 12 months, categorised by sexual orientation, gender and state or territory of residence.

### Suicidal thoughts, attempts, plan, and self-harm among LGBTQA+ WTI4 (2019) participants by personal characteristics.

The visualisation illustrates WTI4 participants who have experienced suicidal thoughts, suicide plan, suicide attempt, or engaged in self-harm, throughout their lifetime and over the last 12 months, categorised by sexual orientation, gender and state or territory.

Choose a variable:  Sexual orientation  Gender  State or territory

Select one:  Suicidal thoughts  Suicide plan  Suicide attempt  Self-harm


Select response:  Ever  Last 12 months  Never  Prefer not to say

Label:  On  Off

Axis range:  Fixed  Automatic

### Sexual orientation | Suicidal thoughts | Ever

Per cent %



Notes: Caution should be exercised when interpreting the results of trans woman participants due to low numbers in the WT14 survey; n.p. - not published.

Source: Writing Themselves In 4 (WT14).

## Results by sexual orientation

The results for sexual orientation for suicidal thoughts, suicide plan, suicide attempt, and self-harm in the lifetime and last 12 months, show little difference between different sexual orientations except for gay participants who consistently had the lowest prevalences of these behaviours. Pansexual, queer, and lesbian participants often had the highest percentages of suicidal and self-harming behaviour, however the differences between sexual orientation categories were often small.

- Pansexual, queer, lesbian and bisexual participants and those classified as ‘something else’ reported the highest percentages of lifetime suicidal thoughts. 84.8% of pansexual participants, 83.1% of queer participants, 81.5% of lesbian participants, 79.3% of bisexual participants and 78.8% of participants classified as ‘something else’ reported having thoughts about suicide, wanting to die, or ending their own life at some point in their lifetimes, followed by asexual and gay participants (75.4% and 68.8%, respectively).
- A similar pattern was seen with recent (last 12 months) suicidal thoughts, which were most likely to be reported by pansexual participants (67.4%) and least likely among gay participants (47.3%).
- A similar pattern of responses was evident for lifetime and recent (last 12 months) suicide plan with the highest proportions among pansexual participants (57.2% and 31.2% respectively) and the lowest among gay participants (37.6% and 17.8% respectively)
- The results for lifetime suicide attempt were similar to suicidal thoughts and suicide plans, from highest to lowest prevalence: pansexual (35.1%), lesbian (30.0%), queer (30.0%), something else (25.6%), bisexual (23.5%), asexual (21.1%), and gay (19.3%) participants.
- For recent suicide attempt, lesbian participants reported the highest percentage (14.1%) followed by pansexual (13.4%) and queer (11.6%) participants, with gay participants the lowest (7.8%)
- Lifetime and recent self-harm followed similar pattern to suicidal thoughts and suicide plan.

Comparing the differences in the order of sexual orientation between the WT14 and PL3 surveys for suicidal thoughts and suicide attempt, lesbian participants rank in the highest three in WT14 survey results but second lowest in the PL3 survey results. This indicates that younger WT14 lesbian participants may be more at risk of suicidal thoughts and suicide attempts than older PL3 lesbian participants.

## Results by gender

Trans man, trans woman, and non-binary participants more likely to have experienced lifetime and recent (last 12 months) suicidal thoughts, suicide plan, suicide attempt, and self-harm than cisgender (man and woman) participants.

- Trans man participants were more likely to experience lifetime suicidal thoughts (92.1%) and suicide attempt (46.9%) than trans woman (90.7% and 40.0% respectively).
- However, trans woman participants were more likely to have experienced recent suicidal thoughts (77.3%) and suicide attempt (20.0%) than trans man participants (73.1% and 16.7% respectively).

## Results by state and territory

The visualisation below illustrates WT14 participants who have experienced suicidal thoughts, suicide plan, suicide attempt, or engaged in self-harm, throughout their lifetime and over the last 12 months, categorised by sexual orientation, gender and state or territory.



There was little difference in terms of likelihood of suicide thoughts, suicide plan, suicide attempt, and self-harm over the lifetime or in the last 12 months by state/territory of participant. However, Tasmania and the Northern Territory (NT) often had the highest percentages compared to the other states and territories.

- Participants from Tasmania and NT were more likely to experience lifetime suicidal thoughts (86% and 83.7%, respectively). Other states ranged from 76.2% in the Australian Capital Territory to 79.6% in New South Wales.
- Lifetime suicide plan ranged from 45.3% for Victoria to 54.0% for Tasmania. Tasmania was also the highest for recent (last 12 months) suicide plan (30.2%). Other states ranged from 22.9% in Western Australia (WA) to 28.6% in the NT for recent suicide plan.
- Lifetime suicide attempts ranged from 22.5% of NT participants to 30.4% of Tasmanian participants. However, NT participants had the highest proportion of recent (last 12 months) suicide attempt at 15.0% (in other states recent suicide attempt ranged 9.4% for Victoria to 11.0% for Queensland).


In the PL3 survey Tasmania was consistently the highest or among the highest states or territories, in terms of likelihood of suicidal thoughts and suicide attempt. It was difficult to assess comparisons with NT due to low NT participant numbers in the PL3 survey.

## Results by age

The visualisation below compares the likelihood of suicidal thoughts, attempts and self-harm among WT14 participants aged 14 to 17 and 18 to 21 years, throughout their lifetime and over the last 12 months.

### Suicidal thoughts, plan, attempts and self-harm among LGBTQA+ WT14 (2019) participants by age.

The visualisation compares the likelihood of suicidal thoughts, attempts and self-harm among WT14 participants aged 14 to 17 and 18 to 21 years, throughout their lifetime and over the last 12 months



There was little difference between 14 to 17-year-olds and 18 to 21-year-olds in terms of likelihood of suicidal thoughts, suicide plan, suicide attempt, and self-harm over the lifetime. However, over the last 12 months, 14 to 17-year-olds were more likely than 18 to 21-year-olds to experience suicidal thoughts (60.3% compared with 55.3%), suicide plan (27.3% compared with 20.2%) suicide attempt (12.0% compared with 7.4%) and self-harm (44.7% compared with 33.5%).

## Results by disability

The visualisation below illustrates disability and disability type of WT14 participants who have experienced suicidal thoughts, suicide plan, suicide attempt, or engaged in self-harm, throughout their lifetime and over the last 12 months.

### Suicide and self-harm support access among LGBTQA+ WT14 (2019) participants by disability status and type.

The visualisation illustrates disability and disability type of WT14 participants who have experienced suicidal thoughts, suicide plan, suicide attempt, or engaged in self-harm, throughout their lifetime and over the last 12 months.

Participants with disability compared to those without were more likely to experience lifetime suicidal thoughts (89.3% compared with 69.4%), suicide plan (64.9% compared with 33.9%), suicide attempt (39.4% compared with 15.7%) and self-harm (78.1% compared with 48.6%). This is also reflected in recent suicidal thoughts (70.5% compared with 47.9%), suicide plan (34.5% compared with 16.6%), suicide attempt (15.7% compared with 6.0%) and self-harm (53.6% compared with 28.1%).


There was little difference between disability types in terms of likelihood of lifetime and recent (last 12 months) suicidal thoughts, suicide plan, suicide attempt and self-harm. Intellectual disability showed the highest likelihood of lifetime and recent (last 12 months) suicide plan and suicide attempt, however the differences from the other disability types were small.

## Results by support service use

The visualisation below illustrates the proportions of WT14 participants who had accessed professional support services in relation to suicide or self-harm, including type of service, by sexual orientation, gender, disability and state or territory.

### Suicide and self-harm support access among LGBTQA+ WT14 (2019) participants by personal characteristics.

The visualisation illustrates the proportions of WT14 participants who had accessed professional support services in relation to suicide or self-harm, including type of service, by sexual orientation, gender, disability and state or territory.



Accessing support services in relation to suicide or self-harm, was related to likelihood of lifetime suicide thoughts, suicide attempt, and self-harm and accessibility of the services themselves.

### Support service use and sexual orientation

- Pansexual and queer participants and those classified as ‘something else’ were most likely to access professional support services in relation to suicide or self-harm (61.6%, 59.4% and 54.1%, respectively), followed by lesbian, bisexual, asexual and gay participants (52.9%, 50.0%, 47.2% and 41.1%, respectively). This order generally reflects the likelihood of lifetime suicidal thoughts, suicide attempt, and self-harm.
- Support services that were not in-person were most likely to be accessed by queer participants - 12.6% of queer participants reported accessing professional telephone support services and 17.1% reported accessing professional text or webchat support services. After queer participants, pansexual and lesbian participants were next most likely to access professional telephone support services (11.8% and 10.5%, respectively), while lesbian and pansexual participants were next most likely to access professional text or webchat support service (15.1% and 14.4%, respectively).

### Support service use and gender

- The results for accessing professional support services in relation to concerns about suicide or self-harm by gender also reflect the likelihood of lifetime suicidal thoughts, suicide attempt, and self-harm. Trans man, trans woman, non-binary, and cisgender woman participants were more likely to report having accessed professional support services (71.1%, 63.8%, 56.7% and 50.8%, respectively), followed by cisgender man participants (38.6%).

## Support service use and disability

- Participants with a disability were more likely to access professional support service in relation to suicide or self-harm, than participants without a disability (68.5%, compared with 38.9%).

## Support service use and state or territory


- Participants in the Australian Capital Territory (ACT) were more likely to access professional support services in relation to suicide or self-harm, with participants from the Northern Territory (NT) the least likely (65.0% of ACT participants compared with 41.7% of those from the NT).
- Access to professional telephone support services was highest in the NT (16.7%). Telephone services may be used more in the NT due to greater challenges in accessing other forms of support. Other states and territories ranged from 8.5% for Queensland to 14.6% for the ACT.

## Suicide attempt, suicidal thoughts by age, sexual orientation, and gender in WT14

The visualisation below illustrates the percentages of WT14 participants aged 14 to 17 and 18 to 21 years, who have experienced suicidal thoughts, attempted suicide, or engaged in self-harm, categorised by sexual orientation and gender, throughout their lifetime and over the last 12 months.

### Suicidal thoughts and attempts among LGBTQA+ WT14 (2019) participants by age, sexual orientation, and gender.

This visualisation illustrates the percentages of WT14 participants aged 14 to 17 and 18 to 21 years, who have experienced suicidal thoughts, attempted suicide, or engaged in self-harm, categorised by sexual orientation and gender, throughout their lifetime and over the last 12 months.



### Results by sexual orientation by age group

Among sexual orientation classifications there was little difference between participants aged 14 to 17 years and 18 to 21 years in the likelihood of lifetime suicidal thoughts, suicide attempt, and self-harm. However, participants aged 14 to 17 years reported an increased likelihood of recent (last 12 months) suicidal thoughts, suicide attempt, and self-harm compared to participants aged 18 to 21 years.

- Participants of all sexual orientations, except bisexual, reported higher likelihood of recent suicidal thoughts in 14 to 17 years age-group compared to the 18 to 21 years age-group, although the differences were small.
- Participants of all sexual orientations, except asexual, reported higher likelihood of recent suicide attempt and self-harm in the 14 to 17 years age-group compared with the 18 to 21 years age group. Compared with suicidal thoughts, the relative differences for suicide attempt and self-harm were larger between the two age groups.

### Results by gender by age group

- Among gender classifications there was little difference between participants aged 14 to 17 years and 18 to 21 years in reported lifetime suicidal thoughts, suicide attempt, and self-harm except for trans woman.

- Trans women participants aged 14 to 17 years were about half as likely as those aged 18 to 21 years to experience lifetime suicide attempt (27.3% to 50.0% respectively) and lifetime self-harm (48.5% to 83.3% respectively).
- All genders, except trans woman, reported higher likelihood of recent (last 12 months) suicidal thoughts, suicide attempt and self-harm in the 14 to 17 years age-group, compared to the 18 to 21 years age-group. As per sexual orientation, the relative differences were larger between the two age groups for suicidal attempt and self-harm than suicidal thoughts.
- Trans women participants aged 14 to 17 years were less likely than those aged 18 to 21 years to experience recent suicide attempt (15.2% to 23.8% respectively) and about half as likely to experience recent self-harm (33.3% to 59.5% respectively).
- Caution should be exercised when interpreting the results for trans woman participants due to low numbers in the WT14 survey, with 33 participants aged 14 to 17 years and 42 aged 18 to 21 years.

## References

- ABS (Australian Bureau of Statistics) (2020) *Standard for Sex, Gender, Variations of Sex Characteristics and Sexual Orientation Variables* (<https://www.abs.gov.au/statistics/standards/standard-sex-gender-variations-sex-characteristics-and-sexual-orientation-variables/2020>), ABS website, accessed 4 July 2023.
- ABS (2020-22a) *National Study of Mental Health and Wellbeing* (<https://www.abs.gov.au/statistics/health/mental-health/national-study-mental-health-and-wellbeing/2020-2022#key-statistics>), ABS Website, accessed 25 October 2023.
- ABS (Australian Bureau of Statistics) (2020-22b) *National Study of Mental Health and Wellbeing methodology* (<https://www.abs.gov.au/statistics/health/mental-health/national-study-mental-health-and-wellbeing/2020-2022#methodology>), ABS website, accessed 25 October 2023.
- AIFS (Australian Institute of Family Studies) (2022) *LGBTIQ+ glossary of common terms* (<https://aifs.gov.au/resources/resource-sheets/lgbtiqa-glossary-common-terms>), AIFS website, accessed 4 July 2023.
- Bretherton I, Thrower E, Zwickl S, Wong A, Chetcuti D, Grossmann M, Zajac JD, Cheung AS 2021, *The Health and Well-Being of Transgender Australians: A National Community Survey* (<https://www.liebertpub.com/doi/10.1089/lgbt.2020.0178>), *LGBT Health*, 8(1):42-49, doi:10.1089/lgbt.2020.0178.
- CCOV (Coroners Court of Victoria) (2022) *Suicide among LGBTIQ+ people*, CCOV website, accessed 2 June 2023.
- Hill AO, Bourne A, McNair R, Carman M & Lyons A (2020) *Private Lives 3: The health and wellbeing of LGBTIQ people in Australia* ([https://www.latrobe.edu.au/\\_data/assets/pdf\\_file/0009/1185885/Private-Lives-3.pdf](https://www.latrobe.edu.au/_data/assets/pdf_file/0009/1185885/Private-Lives-3.pdf)), ARCSHS Monograph Series No. 122. Melbourne, Australia: Australian Research Centre in Sex, Health and Society, La Trobe University.
- Hill AO, Lyons A, Jones J, McGowan I, Carman M, Parsons M, Power J, Bourne A (2021) *Writing Themselves In 4: The health and wellbeing of LGBTIQ+ young people in Australia. National report* ([https://www.latrobe.edu.au/\\_data/assets/pdf\\_file/0010/1198945/Writing-Themselves-In-4-National-report.pdf](https://www.latrobe.edu.au/_data/assets/pdf_file/0010/1198945/Writing-Themselves-In-4-National-report.pdf)), ARCSHS monograph series number 124. Melbourne: Australian Research Centre in Sex, Health and Society, La Trobe University.
- Hottes TS, Bogaert L, Rhodes AE, Brennan DJ & Gesink D (2016) *Lifetime Prevalence of Suicide Attempts Among Sexual Minority Adults by Study Sampling Strategies: A Systematic Review and Meta-Analysis* (<https://doi.org/10.2105/AJPH.2016.303088>), *American journal of public health*, 106(5), e1-e12, doi:org/10.2105/AJPH.2016.303088.
- Kirakosian N, Stanton AM, McKetchnie SM, King D, Dolotina B, O'Cleirigh C, Grasso C, Potter J, Mayer KH, Batchelder AW (2023) *Suicidal Ideation Disparities Among Transgender and Gender Diverse Compared to Cisgender Community Health Patients* (<https://doi.org/10.1007/s11606-022-07996-2>). *J Journal of General Internal Medicine*, 38(6):1357-1365, doi.org/10.1007/s11606-022-07996-2.
- Leske S, Adam G, Catakovic A, Weir B & Kölves K (2022) *Suicide in Queensland: Annual Report 2022* ([https://www.griffith.edu.au/\\_data/assets/pdf\\_file/0033/1639473/AISRAP-Annual-Report-2022.pdf](https://www.griffith.edu.au/_data/assets/pdf_file/0033/1639473/AISRAP-Annual-Report-2022.pdf)), Australian Institute for Suicide Research and Prevention, World Health Organization Collaborating Centre for Research and Training in Suicide Prevention, School of Applied Psychology, Griffith University, Brisbane, Queensland, Australia.
- LGBTIQ+ Health Australia 2021, *Snapshot of Mental Health and Suicide Prevention Statistics for LGBTIQ+ People* ([https://assets.nationbuilder.com/lgbtihealth/pages/549/attachments/original/1648014801/24.10.21\\_Snapshot\\_of\\_MHSP\\_Statistics\\_for\\_LGBTIQ\\_People\\_-\\_Revised.pdf?1648014801](https://assets.nationbuilder.com/lgbtihealth/pages/549/attachments/original/1648014801/24.10.21_Snapshot_of_MHSP_Statistics_for_LGBTIQ_People_-_Revised.pdf?1648014801)), LGBTIQ+ Health Australia website, accessed 2 June 2023.
- Stinchcombe A & Hammond NG (2021) *Sexual orientation as a social determinant of suicidal ideation: A study of the adult life span* (<https://doi.org/10.1111/sltb.12754>), *Suicide and Life-Threatening Behavior*, 51(5):864-871, doi.org/10.1111/sltb.12754.
- The Commonwealth of Australia (Cth of Australia) (2022) *National Mental Health and Suicide Prevention Agreement* (<https://federalfinancialrelations.gov.au/agreements/mental-health-suicide-prevention-agreement>), The Federal Financial Relations website, accessed 15 June 2023.
- Marchi M, Arcolin E, Fiore G, Travascio A, Uberti D, Amaddeo F, Converti M, Fiorillo A, Mirandola M, Pinna F, Ventriglio A, Galeazzi, GM & Italian Working Group on LGBTIQ Mental Health (2022) *Self-harm and suicidality among LGBTIQ people: a systematic review and meta-analysis* (<https://doi.org/10.1080/09540261.2022.2053070>), *International Review of Psychiatry*, 34:3-4, 240-256, doi.org/10.1080/09540261.2022.2053070.

Strauss P, Cook A, Winter S, Watson V, Wright Toussaint D, Lin A (2017) *Trans Pathways: the mental health experiences and care pathways of trans young people. Summary of results* (<https://www.telethonkids.org.au/globalassets/media/documents/brain--behaviour/trans-pathwayreport-web.pdf>). Telethon Kids Institute, Perth, Australia.

Swannell S, Martin G, Page A 2016 *Suicidal ideation, suicide attempts and non-suicidal self-injury among lesbian, gay, bisexual and heterosexual adults: Findings from an Australian national study* (<https://doi.org/10.1177/0004867415615949>), *Australian & New Zealand Journal of Psychiatry*, 50(2):145-53, doi.org/10.1177/0004867415615949.

Zwickl S, Wong AFQ, Dowers E, Leemaqz YN, Bretherton I, Cook T, Zajac JD, Yip PSF & Cheung AS (2021) *Factors associated with suicide attempts among Australian transgender adults* (<https://pubmed.ncbi.nlm.nih.gov/33557793/>), *BMC Psychiatry* 21:81 doi.org/10.1186/s12888-021-03084-7.

Zwickl S et al. (2023) Depression, self-harm, and suicidality in the Australian trans community during the COVID-19 Pandemic. [Manuscript in preparation].

---

© Australian Institute of Health and Welfare 2024



## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on [13 11 14](tel:131114), or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)

## Culturally and Linguistically Diverse Australians: Suicide among refugee and humanitarian entrants and other permanent migrants

'Culturally and linguistically diverse' (CALD) is a broad term describing the cultural and linguistic diversity of multicultural populations living in Australia (ABS 1999). Refugee and humanitarian entrants, and other permanent migrant Australians are part of the larger CALD communities within Australia.

In 1999, The Australian Bureau of Statistics (ABS) introduced the 'Standards for Statistics on Cultural and Language Diversity' (the Standards). The Standards were designed to replace the use of 'non-English speaking background' as the sole indicator of cultural and linguistic diversity. They provide a more holistic, accurate, and consistent measurement of cultural and linguistic diversity in Australia. While the ABS has not revised the Standards since the initial publication, text and formatting were refreshed in 2022 (ABS 2022a).

The Standards comprise indicators related to country of birth (of an individual and their parents), year of arrival in Australia, language(s) spoken, ancestry, religious affiliation, and First Nations status. The AIHW uses 'First Nations people' to refer to Aboriginal and/or Torres Strait Islander people in this publication. While CALD terminology as outlined by the Standards is widely adopted within Australia, there is no universally accepted definition of CALD (Pham et al. 2021).

CALD communities and refugees are identified as priority populations under the *National Mental Health and Suicide Prevention Agreement* (the Agreement) (Cth of Australia 2022). Under the Agreement, Commonwealth and State and Territory governments have a shared responsibility to support priority populations who may be at higher risk of mental ill health and suicide due to vulnerability caused by social, economic, and environmental circumstances. While CALD communities have varied experiences, they may also have some shared experiences that contribute to suicide risk factors. These include difficulties adjusting to a new culture, experiences of stigma, and changes in social and family networks a result of migration (Bowden et al. 2020). CALD Australians who are refugees or humanitarian entrants may experience additional or more pronounced challenges due to past experiences of persecution or human rights abuses within their country of origin, or trauma associated with war or their refugee journey (FASSTT 2017).

Although First Nations people are diverse in language and culture, their experiences as First Nations people are unique. Furthermore, the *National Mental Health and Suicide Prevention Agreement* (Cth of Australia 2022) identifies 'Aboriginal and Torres Strait Islander peoples' as priority populations separate to the identification of 'culturally and linguistically diverse communities and refugee' priority populations. As such, First Nations people are considered distinct from CALD terminology used throughout this report. Though it is acknowledged that a person may both identify as a First Nations person and as a person of Cultural and Linguistic Diversity.

### Data on suicide among refugee and humanitarian entrants and other permanent migrants

Data presented in this report are drawn from a [larger project](#) investigating the health and welfare of Australia's refugee and humanitarian entrant populations (AIHW 2023a). This larger project was funded by the Department of Home Affairs and involved linking the Settlement Database (Department of Home Affairs 2019) with other datasets available in the Person-level Integrated Data Asset (PLIDA); formally known as the Multi Agency Data Integration Project (MADIP) (ABS n.d.). Linking the Settlement Database to the PLIDA enabled the identification and analysis of migrant status for deidentified individuals appearing within other PLIDA datasets.

The key datasets used in the analysis presented in this publication are the Settlement Database (Department of Home Affairs 2019) and the Causes of Death (ABS 2023) dataset. For full details see [technical notes for the ‘Health of refugees and humanitarian entrants in Australia’ report](#), which was undertaken as part of the larger Department of Home Affairs funded project for details.

Information derived from analysis of linked Settlement Database and Causes of Death dataset data, is limited to the experience of people who have moved to Australia from another country (first-generation migrant Australians). Refugee and humanitarian, and other permanent migrants are part of the broader CALD communities within Australia. Migration status and year of arrival capture a limited number of indicators within ABS ‘Standards for Statistics on Cultural and Language Diversity’. Therefore, data presented should not be considered representative of broader CALD communities within Australia. Instead, this analysis provides robust information about deaths by suicide among first-generation permanent migrant Australians: refugee and humanitarian entrants, and other permanent migrants.


### First generation ‘Humanitarian entrants’ and ‘Other permanent migrants’ experienced lower rates of suicide compared to the ‘Rest of the Australian population’.

The interactive data visualisation below displays deaths by suicide between 2007-2020, by sex and migration group. The age-standardised rates show that:

- Both ‘Humanitarian entrants’ and ‘Other permanent migrants’ each experienced lower rates of suicide when compared to the ‘Rest of the Australian population’.
- ‘Humanitarian entrants’ experienced 1.7 times the rate of suicide compared to ‘Other permanent migrants’.
- Deaths by suicide were higher for males than females across all three cohorts, with the largest difference in the ‘Humanitarian entrants’ cohort. Among the ‘Humanitarian entrants’ cohort, the age-standardised suicide rate for males (11 per 100,000 population) was more than 3.5 times higher than for females (3.0 per 100,000 population). Among the ‘Rest of the Australian population’ cohort, the age-standardised suicide rate for males (18.8 per 100,000 population) was approximately three times higher than for females (6.1 per 100,000 population).

This bar chart shows the age-standardised suicide rate (per 100,000), crude suicide rate (per 100,000 population) and number of deaths by suicide among by sex and migration group (other permanent migrants and humanitarian entrants), over the years 2007–2020. Migration groups are compared to the rest of the Australian population.

#### Deaths by suicide by sex and migration group arriving on or after 2000, 2007–2020



Supplementary Table: Table Permanent Migrants S1  
Source: PLIDA-linked datasets: Settlement Database and the Causes of Death dataset.  
Latest data: 2020

[See Notes ▶](#)

### Rates of death by suicide for first generation ‘Humanitarian entrants’ and ‘Other permanent migrants’ increased with time since arriving in Australia

The interactive data visualisation below displays deaths by suicide between 2007 and 2020, disaggregated by time since arrival in Australia and migration group. Rates were calculated using person years, which account for the different lengths of time individuals have been living in Australia (and at risk of death by suicide). This enables examination of how these migrant groups are affected by suicide at different times after their arrival in Australia.


The age-standardised rates show that:

- Deaths by suicide were higher for ‘Humanitarian entrants’ compared to ‘Other permanent migrants’ at each period since arrival in Australia. Although, particularly with error bars displayed, the differences between these groups are small.

- Deaths by suicide increased for both ‘Humanitarian entrants’ and ‘Other permanent migrants’ as time since arrival in Australia increased.

This bar chart shows the age-standardised suicide rate (per 100,000), crude suicide rate (per 100,000 population) and number of deaths by suicide among by migration group (other permanent migrants and humanitarian entrants) and age group, over the years 2007–2020. The age group includes ‘<5 years’, ‘5-10 years’ and ‘>10 years’.

### Deaths by suicide by time since arrival and migration group, 2007–2020



Supplementary Table: Table Permanent Migrants S2

Source: PLIDA-linked datasets: Settlement Database and the Causes of Death dataset.

Latest data: 2020

[See Notes ►](#)

### What other national suicide and self-harm data available for CALD and refugee communities in Australia?

The AIHW uses the [National Mortality Database](#) (AIHW 2023b) and the [National Hospital Morbidity Database](#) (AIHW 2023c) to report on key suicide and self-harm statistics in Australia. Information relevant to CALD communities within these databases is limited to country of birth.

The most recent [ABS Causes of Death publication \(ABS 2023\)](#) (<https://www.abs.gov.au/statistics/health/causes-death/causes-death-australia/latest-release/intentional-self-harm-deaths-suicide-in-australia>), which presents National Mortality Database data, includes information about suicide by country of birth. This publication found that, between 2018-2022, those born in Croatia, New Zealand and Scotland had a higher age-standardised suicide rate than those born in Australia (ABS 2023). There is no information specific to refugee status within with the National Mortality Database or the National Hospital Morbidity Database.

Linking datasets that contain more comprehensive information about members of CALD communities and/or refugee status and administrative datasets, can provide insight into suicide and self-harm among these communities. For example, while not focused on suicide and self-harm, the AIHW (2022) report “[Reporting on the health of culturally and linguistically diverse populations in Australia: An exploratory paper](#)”, investigates the use of PLIDA linked data to report on the health of CALD populations. Linking the Settlement Database to the PLIDA, made the analysis presented within this release possible. Migration status and time since arrival in Australia indicators drawn from the Settlement Database provide additional information regarding the experience of first-generation permanent migrant Australians: refugee and humanitarian entrants, and other permanent migrants.

The National Study of Mental Health and Wellbeing (ABS 2022b) collects information both about cultural and linguistic diversity, and about participants’ lived experience of suicide and self-harm. The ABS conducts this nationally representative survey on an irregular basis, most recently during 2020-2022. Analysis of this survey data may be informative for future work.

### References

ABS (Australian Bureau of Statistics) (n.d.) [Multi-agency data integration project \(MADIP\)](#) (<https://www.abs.gov.au/about/data-services/data-integration/integrated-data/person-level-integrated-data-asset-plida>), ABS website, accessed 1 August 2023.

ABS (1999). [1289.0 - Standards for statistics on cultural and language diversity](#) (<https://www.abs.gov.au/ausstats/abs@.nsf/mf/1289.0>), ABS website, accessed 23 October 2023.

ABS (2022a). [Standards for statistics on cultural and language diversity](#). Australian Bureau of Statistics (<https://www.abs.gov.au/statistics/standards/standards-statistics-cultural-and-language-diversity/latest-release>), ABS website, accessed 23 October 2023.



ABS (2022b). *National study of mental health and wellbeing methodology*. Australian Bureau of Statistics (<https://www.abs.gov.au/methodologies/national-study-mental-health-and-wellbeing-methodology/2020-21>), ABS website, accessed 23 October 2023.

ABS (2023). *Causes of death, Australia, 2022* (<https://www.abs.gov.au/statistics/health/causes-death/causes-death-australia/latest-release>), ABS website, accessed 23 October 2023.

AIHW (Australian Institute of Health and Welfare) (2022) *Reporting on the health of culturally and linguistically diverse populations in Australia: An exploratory paper*, AIHW, Australian Government, accessed 23 October 2023.

AIHW (2023a) *Culturally and linguistically diverse Australians. Refugee and humanitarian entrant health*, AIHW, Australian Government, accessed 3 November 2023.

AIHW (2023b) *National Mortality Database (NMD)*, AIHW, accessed 25 October 2023.

AIHW (2023c) *National Hospitals Data Collection*, AIHW, accessed 25 October 2023.

Bowden M, McCoy A, and Reavley N (2020) 'Suicidality and suicide prevention in culturally and linguistically diverse (CALD) communities: A systematic review', *International Journal of Mental Health*, 49(4):293-320, doi: [10.1080/00207411.2019.1694204](https://doi.org/10.1080/00207411.2019.1694204) (<https://doi.org/10.1080/00207411.2019.1694204>)

Cth of Australia (The Commonwealth of Australia) (2022). *National mental health and suicide prevention agreement* (<https://federalfinancialrelations.gov.au/agreements/mental-health-suicide-prevention-agreement>), Federal Financial Relations, accessed 23 October 2023.

Department of Home Affairs (2019). *Settlement reports* (<https://immi.homeaffairs.gov.au/settling-in-australia/settlement-reports>). Australian Department of Home Affairs, Immigration and Citizenship, accessed 23 October 2023.

FASSTT (The Forum of Australian Services for Survivors of Torture and Trauma) (2017), *Never turning away Australia's world-leading program of assistance to survivors of torture and trauma (PASTT)* (<https://www.fasstt.org.au/publications/>), FASSTT, accessed 23 October 2023.

Pham TTL, Berecki-Gisolf J, Clapperton A, O'Brien KS, Liu S and Gibson K (2021) 'Definitions of culturally and linguistically diverse (CALD): A Literature review of epidemiological research in Australia' *International Journal of Environmental Research and Public Health*, 18(2):737, doi: [10.3390/ijerph18020737](http://dx.doi.org/10.3390/ijerph18020737) (<http://dx.doi.org/10.3390/ijerph18020737>)



---

## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on [13 11 14](tel:131114), or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)

## The use of mental health services, psychological distress, loneliness, suicide, ambulance attendances and COVID-19

While there has been a rise in the use of mental health and crisis services during the COVID-19 pandemic, the pandemic was not associated with a rise in suspected deaths by suicide in 2020 and 2021, see [Suspected deaths by suicide](#) for details. Preliminary national mortality data published by the Australian Bureau of Statistics for 2020 to 2021 show that the rate of death by suicide in Australia was lower in 2020 (12.1 per 100,000 population) and 2021 (12.0) than in 2019 (13.1), see [Deaths by suicide over time](#).

Deaths referred to the coroner where initial investigation points to suicide are referred to as 'suspected deaths by suicide'. In some cases it can take a number of years for the coronial process to determine if suicide was the cause of a specific death. These data are not directly comparable with coroner-certified deaths as reported in [Deaths by suicide in Australia](#) or published by the Australian Bureau of Statistics as 'Causes of Death, Australia ([https://www.abs.gov.au/statistics/health/causes-death#:~:text=Provisional%20Mortality%20Statistics&text=The%20age%20standardised%20death%20rate,in%20November%20\(see%20article\).](https://www.abs.gov.au/statistics/health/causes-death#:~:text=Provisional%20Mortality%20Statistics&text=The%20age%20standardised%20death%20rate,in%20November%20(see%20article).))'.

---

### Use of mental health services

Since April 2020, the AIHW has compiled and reported mental health-related data. This reporting commenced as part of the National Suicide and Self-harm Monitoring Project. Data from the Medicare Benefits Schedule (MBS), Pharmaceutical Benefits Scheme (PBS), and crisis and support organisations (Lifeline, Beyond Blue, Kids Helpline) were collected weekly in 2020, fortnightly in 2021 and monthly in 2022. These data are shared within government to inform the mental health response to the COVID-19 pandemic. The most current data are available at [Mental health services in Australia](#).

### Increased demand for mental health services and crisis and support organisations between 2020 and 2022

These data show that since the onset of the COVID-19 pandemic, overall, there has been a substantial increase in the use of crisis and support organisations (as measured by the number of calls or other contacts, such as webchat or email) and mental health-related services (as measured by MBS and PBS claims processed). The extent to which this increase in demand has been driven by a rise in psychological distress (rather than an increase in people seeking assistance for other reasons, such as loneliness or concern about contracting COVID-19) is unclear. However, given a range of survey data indicate that the average level of psychological distress rose in Australia in 2020 and 2021 from pre-pandemic levels (see psychological distress below), increased demand for mental health-related services and crisis and support organisations is almost certainly indicative of an increase in the need for mental health support and assistance as a result of the pandemic.

---

### Psychological distress

Psychological distress is commonly measured using the Kessler Psychological Distress Scale—10 items (K10). The K10 questionnaire was developed to yield a global measure of psychological distress, based on questions about people's level of nervousness, agitation, psychological fatigue and depression in the past 4 weeks. The Kessler 6 Scale is an abbreviated version of K10.

There is an association between high levels of psychological distress and serious mental health disorders. As a result, instruments such as K10 and K6 can be used in representative sample surveys as a broad indicator of the level of these disorders in the Australian population (Slade, Grove & Burgess 2011). This is important, as there is an association between mental health issues and deaths by suicide. Data from the Queensland Suicide Register for 2016-2018 based on police and coroners reports, show that 53% of people who died from suicide

reportedly had a mental health condition (Leske et al. 2022). Leske et al. (2022) note that this is likely to be an underestimate. Australia wide, just under 63% of people who died by suicide in 2021 had a recorded mental and behavioural disorder as an associated cause of death (ABS 2022a).

There are several ways to gain insights into the level of psychological distress in the community and monitor trends over time.

One way is to look at trends in severe levels of psychological distress through the use of mental health and crisis support services. The AIHW has compiled data on the use of mental health services and crisis and support organisations during the COVID-19 pandemic. The most current data are available at [MHSA](#). However, while this approach is useful, it is not a direct measure of the level of psychological distress in the population. It also does not capture those who choose not to seek assistance or those who cannot access mental health services or crisis and support organisations.

Another way to analyse trends in the level of psychological distress since the onset of the pandemic is to use sample surveys. This approach has been challenging since the onset of COVID-19 due to the fact that face-to-face surveys are very difficult to undertake at this time and pose a potential health and safety risk to interviewers and interviewees. This has led to a number of online surveys being conducted but many of these surveys are not based on probability sampling. In some cases, samples are drawn by inviting all members of the public above a certain age to respond, with unknown response rates. Other samples are drawn from panels where individuals opt-in online. While this sort of approach can provide some useful information, results are unlikely to be representative of the Australian population and therefore cannot be used, even with reweighting, to derive estimates for the Australian population. A major report on online panels for the American Association for Public Opinion Research (AAPOR 2010) noted that:

Researchers should avoid nonprobability online panels when one of the research objectives is to accurately estimate population values. There currently is no generally accepted theoretical basis from which to claim that survey results using samples from nonprobability online panels are projectable to the general population. Thus, claims of “representativeness” should be avoided when using these sample sources. Given the need for representative data, the AIHW collaborated with the Centre for Social Research and Methods at the Australian National University to include questions on loneliness and levels of psychological distress in the ANUpoll surveys, which collect data using the Life in Australia™ Panel. Importantly, this panel uses random probability-based sampling methods and covers both online and offline populations (that is, people who do and do not have access to the internet). In addition, ANUPoll data collected between January 2020 and January 2021 were used to develop a longitudinal dataset. This longitudinal data set enables changes in the experience of an individual to be measured across this period (Biddle 2021a). Data on psychological distress during the COVID-19 pandemic have been collect in April, May, August, October and November 2020, January, April, August and October 2021, January, April, August, October 2022, and January 2023.

## Pre COVID-19 snapshot

To understand how COVID-19 may have affected Australians’ levels of psychological distress, it is important to look at data from before the pandemic. It is particularly important to consider any existing trends prior to the pandemic—for example, if psychological distress was generally increasing among Australians in the years before the pandemic.

This is possible using results from the Australian Bureau of Statistics’ National Health Survey (NHS), which is conducted approximately every 3 years. [Tables 1 to 3](#) show the proportion of males, females and people with high or very high levels of psychological distress, as measured by the Kessler 10 Scale, from 2004-05 to 2017-18. While the results vary by age, there is no consistent trend over this period. It is worth noting, however, that young women aged 18-24 generally have higher levels of psychological distress than other age groups. Overall, between 2011-12 to 2017-18 there appears to have been small increases in the proportion of both males and females with high or very high levels of psychological distress.

The National Drug Strategy Household Survey (NDSHS) also showed an increase in the proportion of people reporting high or very high levels of psychological distress; from 10% in 2010 to 14% in 2019 (AIHW 2020).

Table 1: Proportion of persons with high/very high psychological distress, by age group and year

|         | 18-24 | 25-34 | 35-44 | 45-54 | 55-64 | 65-74 | 75+  | Total |
|---------|-------|-------|-------|-------|-------|-------|------|-------|
| 2004-05 | 15.5  | 11.8  | 14.9  | 13.7  | 12.2  | 11.2  | 10.7 | 13.0  |
| 2007-08 | 11.8  | 12.3  | 11.5  | 13.2  | 13.2  | 9.0   | 10.8 | 12.0  |
| 2011-12 | 11.8  | 10.9  | 10.9  | 11.4  | 10.4  | 9.3   | 9.3  | 10.8  |
| 2014-15 | 15.4  | 10.9  | 11.9  | 12.4  | 11.0  | 10.1  | 9.7  | 11.7  |
| 2017-18 | 15.2  | 13.1  | 11.8  | 14.3  | 14.4  | 10.7  | 8.8  | 13.0  |

Source: ABS 4364.0.55.001 - National Health Survey

Table 2: Proportion of males with high/very high psychological distress, by age group and year

|         | 18-24 | 25-34 | 35-44 | 45-54 | 55-64 | 65-74 | 75+  | Total |
|---------|-------|-------|-------|-------|-------|-------|------|-------|
| 2004-05 | 12.4  | 9.3   | 11.4  | 11.0  | 11.3  | 9.8   | 10.8 | 10.8  |

|         |      |      |      |      |      |      |     |      |
|---------|------|------|------|------|------|------|-----|------|
| 2007-08 | 6.5  | 10.9 | 8.8  | 11.1 | 11.6 | 7.4  | 8.8 | 9.6  |
| 2011-12 | 10.6 | 8.2  | 9.1  | 8.9  | 8.8  | 7.0  | 7.8 | 8.8  |
| 2014-15 | 11.1 | 9.8  | 10.0 | 9.2  | 9.8  | 9.7  | 9.0 | 9.9  |
| 2017-18 | 12.4 | 11.8 | 10.3 | 13.2 | 12.1 | 10.0 | 7.5 | 11.3 |

Source: ABS 4364.0.55.001 - National Health Survey

Table 3: Proportion of females with high/very high psychological distress, by age group and year

|         | 18-24 | 25-34 | 35-44 | 45-54 | 55-64 | 65-74 | 75+  | Total |
|---------|-------|-------|-------|-------|-------|-------|------|-------|
| 2004-05 | 18.7  | 14.3  | 16.6  | 16.3  | 13.1  | 12.5  | 10.7 | 15.1  |
| 2007-08 | 17.4  | 13.9  | 14.2  | 15.3  | 14.7  | 10.6  | 12.3 | 14.3  |
| 2011-12 | 13.0  | 13.6  | 12.7  | 13.8  | 11.9  | 11.6  | 10.5 | 12.7  |
| 2014-15 | 20.0  | 11.7  | 13.7  | 15.4  | 12.3  | 10.3  | 10.6 | 13.5  |
| 2017-18 | 18.4  | 14.7  | 13.7  | 15.3  | 16.9  | 11.3  | 9.7  | 14.5  |

Source: ABS 4364.0.55.001 - National Health Survey

### Psychological distress during COVID-19

The Australian Bureau of Statistics' National Study of Mental Health and Wellbeing, 2020-21 showed that 15% of Australians aged 16-85 experienced high or very high levels of psychological distress as measured by the Kessler 10 scale in 2020-21 (ABS 2022b). However, being from a different survey, these results are not directly comparable with the pre-COVID-19 results from the NHS or NDSHS.

The ANUpoll collected data on psychological distress using the K6 scale prior to and during the COVID-19 pandemic. Data were collected from 2,500 respondents in 2017 (February). Data were collected from over 3,000 respondents for each of the ANUpoll's administered during 2020 (April, August, October and November), 2021 (January, April, August and October), 2022 (January, April, August and October), and 2023 January.


### Prevalence of 'severe' psychological distress from 2020 to 2023

The K6 scale can be used to categorise those survey respondents experiencing 'severe' psychological distress, consistent with having a 'probable serious mental illness' according to their K6 score (Prochaska et al. 2012). Figure 1, below, shows the proportion of people experiencing severe psychological distress for each of wave of the ANUpoll during the COVID-19 pandemic as well as prior to the pandemic in February 2017.

Results of the surveys show that the proportion of the population experiencing severe psychological distress rose from 8.4% (of people aged 18+) in February 2017 to 10.6% in April 2020 (Biddle et al. 2020b) with a subsequent fall to 9.7% in May 2020 followed by an increase to 10.9% in October 2020, and a substantial decrease to 9.4% in January 2021 (Biddle & Gray 2021b). The proportion of people experiencing severe psychological distress in August 2021 (10.1%) was similar to April 2021 (9.7%) and continued to be significantly higher than in February 2017 (8.4%) (Biddle et al. 2021c). Between August and October 2021, there was a large and statistically significant increase in the proportion of Australians experiencing severe psychological distress from 10.1% to 12.5%. At 12.5% in October this proportion was at its highest level recorded since the onset of the COVID-19 pandemic (Biddle et al. 2021d). The proportion of people experiencing severe psychological distress declined to 11.0% in January 2022, increased slightly to 11.6% in April 2022 (Biddle 2022a,b), before declining again in August 2022 to 9.7%. (Biddle, McAllister and Sheppard 2022). The proportion of Australians experiencing severe psychological distress again increased to 10.8% in October 2022 (Biddle 2022c). In January 2023, 10.9% of Australian's experienced severe psychological distress (Biddle & Gray 2023).

While the percentage changes in the proportion of Australians experiencing severe psychological distress between 2020 and January 2023 may appear to be small, a change of 1% point represents a change in the experience of approximately 200,000 people. This calculation is based on the June 2021 estimated resident adult Australian population of about 20,000,000 people (ABS 2022c).

**Figure 1: Proportion of Australians aged 18 years and over experiencing severe psychological distress, by survey month from February 2017 to January 2023**



### Average psychological distress (K6 score) in 2020, 2021 and 2022

Another way of analysing trends in psychological distress is to look at how the average K6 score has changed over time. The data show some notable changes in the average level of psychological distress between 2020 and 2023, with marked differences by age and some variation by gender and jurisdiction. The K6 measure of psychological distress used in the analysis prepared by the ANU has been constructed to have a minimum value of 6 and a maximum value of 30 (Biddle et al. 2020c). Higher scores indicate higher average levels of distress. People with a sum of 11-18 out of a possible maximum of 30 are categorised as experiencing 'moderate' psychological distress. This group can be considered to be struggling with mental distress worthy of mental health support but are not at risk of clinical levels of mental health problems (Prochaska et al. 2012). Those with a K6 sum of 19 or higher out of a possible maximum of 30 are categorised as experiencing 'severe' psychological distress consistent with having a 'probable serious mental illness'.


In February 2017, the average K6 score was 11.2. In April 2020, it increased to 11.9. Between April and May 2020 there was a significant reduction in psychological distress (to an average score of 11.5); however, the score remained above the pre-COVID-19 level of February 2017. The average score rose from 11.5 in May 2020 to 11.7 in August 2020 but showed very little change from August to October 2020 (11.8) (Biddle & Gray 2020). This was followed by a large and statistically significant fall in the average K6 score from October to November 2020 (11.4) (Biddle et al. 2020e). While the average score in November 2020 was quite a bit lower than it was in April 2020 it was higher than it was prior to the onset of COVID-19 in February 2017.

From November 2020 to January 2021, the average K6 score fell from 11.4 to 11.2 and continued to decline to 11.1 in April 2021 (Biddle & Gray 2021b). In August 2021, following the most recent outbreak of COVID-19, the average K6 score increased (worsened) to 11.3 but remained lower than scores recorded in April to October 2020. The increase in psychological distress between April and August 2021 was greater for women than men (after controlling for levels of psychological distress in April 2021). Also, after controlling for other factors, women had higher levels of psychological distress in August 2021 compared with men (Biddle & Gray 2021c).

The average level of psychological distress in August 2021 (11.3) was not significantly different to the pre-pandemic level (11.2) observed in February 2017 (Biddle & Gray 2021c). In other words, the average level of psychological distress had returned to pre-pandemic levels. That said, however, there were differences by age (Figure 3).

Between August and October 2021, the average level of psychological distress increased significantly from 11.3 to 11.7 but remained lower than the peaks April and October 2020 (11.9 and 11.8, respectively) (Biddle & Gray 2021d). The worsening of psychological distress between August and October 2021 was due to increases among people aged 18 to 44 years and 75 years and older (Figure 2). Between October 2021 and January 2022, there was a slight but not statistically significant decline in average psychological distress from 11.7 to 11.6 (Biddle & Gray 2022). Average psychological distress remained at 11.6 in April 2022 (Biddle 2022b). There may have then been a small decline in average psychological distress to 11.3 during August 2022 (Biddle, McAllister and Sheppard 2022). Average psychological distress remained stable at 11.3 in October 2022 and January 2023 (Biddle 2022c; Biddle & Gray 2023). Since August 2022, average psychological distress has no longer been significantly higher than the pre-COVID-19 level of February 2017 (p10., Biddle & Gray 2023).

**Figure 2: Average psychological distress of Australians aged 18 years, by survey month from February 2017 to January 2023**




### Psychological distress by age

Figure 3 shows average K6 scores by age groups. The chart shows a clear gradient with younger people experiencing higher average levels of psychological distress than people in older age groups (Biddle et al. 2020c,d; Biddle & Gray 2020; Biddle & Gray 2021a,b,c,d). The chart also shows a distinct pattern over time. For younger people (18-44), average levels of psychological distress were higher in 2020, 2021 and January and April 2022 than they were before the pandemic, especially for those aged 18-24. Those aged 45 and above experienced either little change or improvements in their level of psychological distress over the same period. As an example, the average levels of psychological distress among those aged 18-24, 25-34 and 35-44 were significantly higher in April 2020 than in February 2017 (Biddle et al. 2020c). However, it is worth noting there were improvements during the course of 2020. For example, the level of psychological distress among those aged 18-24 showed a significant improvement from October to November 2020 (Biddle et al. 2020e). Furthermore, average levels of distress for people within the younger age groups (18-24, 25-34 and 35-44 years) appear to have either lowered or remained relatively stable between April 2022 and January 2023 (Biddle 2022c; Biddle & Gray 2023; Biddle, McAllister and Sheppard 2022).

### Figure 3: K6 measure of psychological distress, by age group and selected survey months from February 2017 to January 2023

This figure shows the average K-6 measure of psychological distress by age group for February 2017, April 2020, October 2021 and April 2022. For young Australians (18 to 24 years old in particular, but also all those aged under 45), there was a worsening in psychological distress between February 2017 and April 2020. Although there were improvements in psychological distress in these age groups, psychological distress remained higher in October 2021 and April 2022 than it was in February 2017. For those aged 55 years and over, average levels of psychological distress were slightly lower in April 2022 than in February 2017

Average Kessler-6 score



Time series view ►

See notes ►

### Demographic factors associated with psychological distress during the COVID-19 pandemic

An advantage of the ANUpoll data is that longitudinal data are available for a proportion of respondents and therefore, the impacts of the COVID-19 pandemic and associated factors can be measured in the same respondents over time. Biddle et al. constructed a statistical (regression) model to investigate the possible impacts of the COVID-19 pandemic on psychological distress, as measured by the K6, from April 2020 to April 2022 (Biddle et al. 2022). The model also included measures of mental health, other than the K6, from the February 2020 ANUpoll to control for pre-pandemic levels of psychological distress. It was necessary to use other mental health measures because, prior to April 2020, the most recent ANUpoll to include the K6 was conducted in April 2017 and the remaining longitudinal sample from this time is insufficient as a baseline for the present regression analysis. The analysis showed that over the experience of the pandemic between April 2020 and April 2022 (Biddle et al. 2022):

- females had significantly higher levels of psychological distress than males
- Aboriginal and Torres Strait Islander Australians had higher levels of psychological distress than non-Indigenous Australians, although the difference was not statistically significant
- Victorians had higher levels of psychological distress than residents of New South Wales (the base case).
- household income was associated with psychological distress, with higher levels of psychological distress experienced by those living in lower income households.

### What contributed to increased levels of psychological distress?

The heightened level of psychological distress in April 2020 coincided with the first wave of COVID-19 infections in Australia and the initial lockdown period, while the improvement from April to May 2020 coincided with the loosening of restrictions. The increase in the level of psychological distress between May and August 2020 coincided with the second wave of COVID-19 in Victoria and the associated lockdown - with much of the worsening in the average K6 score over this period reflecting changes in Victoria (Biddle & Gray 2020).

To test whether outcomes worsened in Victoria relative to the rest of the country after the reintroduction of lockdowns in July 2020, Biddle et al. conducted a difference-in-difference analysis using linked data for May and August (that is, data across these months for the same people) (Biddle et al. 2020d). This showed a significant worsening in Victoria relative to the rest of the country on several outcomes including: psychological distress, loneliness, life satisfaction, satisfaction with direction of country, likely to be infected by COVID-19 and hours worked). On the other hand, the decrease in levels of psychological distress from October to November 2020 coincided with improvements in Victoria. According to Biddle et al. (2020e):

There has also been a continued convergence in psychological distress between Victoria and the rest of Australia. In October 2020, just as lockdown conditions had started to be eased, psychological distress in Victoria was more than 1-point higher in Victoria compared to the rest of Australia (12.67 compared to 11.52). By November 2020, however, this difference had declined to less than half of one point - 11.73 compared to 11.32.

In 2021, the worsening of psychological distress between April and August was greater for residents of Sydney and Melbourne, than those living in the rest of Australia (Biddle & Gray 2021c). The increase in psychological distress from August to October did not show a clear geographic pattern.

To investigate the factors that appeared to be contributing to the rises in psychological distress Biddle et al. undertook regression modelling of the ANUpoll longitudinal subset (Biddle et al. 2020c). Modelling of ANUpoll data collected from the same respondents in February 2017, April 2020 and May 2020 showed that the strongest predictor of psychological distress (K6 score) was 'increased (worsened) stress' (this is not surprising as stress is a key predictor of poor mental health outcomes) (Biddle et al. 2020c). 'Increased loneliness' was also a strong predictor of K6 scores even when other factors like changes in employment status were controlled for. This suggests that increased loneliness during the pandemic is of concern and that increases in psychological distress are not only being driven by job loss.

Job loss itself was a predictor of K6 scores in the modelling (Biddle et al. 2020c). Controlling for other factors, people who were employed in February 2020 but not in May 2020 had higher levels of psychological distress than those who were employed. In all the models, those living outside capital cities had lower rates of psychological distress than those living in capital cities, after controlling for other factors (Biddle et al. 2020c). This is probably a reflection of the fact that infection rates and the economic impacts of lockdowns have been higher in major cities than they have been in regional or remote areas.

After controlling for factors such as 'relationships worsening', 'increased stress' and 'loss of employment', there was no significant difference in K6 scores between young people (18-24) and older people. This suggests that these factors were the drivers of higher levels of psychological distress among young people.

One final point worth noting is that in the regression analysis of K6 scores, previous K6 scores in February 2017 had a significant predictive effect on K6 scores for May 2020 (Biddle et al. 2020c). This shows that people who are already experiencing high levels of psychological distress can be particularly vulnerable when the situation worsens.

As the COVID-19 pandemic continues, it becomes less clear whether pandemic related factors or other socioeconomic factors may be primarily responsible for any community level change in psychological distress. In the August and October 2022 and January 2023 ANUpolls, respondents were asked whether they 'felt anxious or worried for the safety of yourself, close family members or friends, due to COVID-19'. Responses to this question provide some information as to whether distress experienced is related to the COVID-19 pandemic or to other factors. In August 2022, 48.3% of respondents reported feeling anxious or worried due to COVID-19. In October 2022, this dropped to 29.8% of respondents, and in January 2023 34.3% of respondents reported feeling anxious or worried due to COVID-19 (Biddle 2022c; Biddle & Gray 2023; Biddle, McAllister and Sheppard 2022).

## Loneliness

The ANUpoll also asked respondents whether 'In the past week, how often have you felt lonely?' Analysis summarised in Biddle et al. (2020c) shows that those who experienced loneliness had higher rates of psychological distress than those who did not.

Between April 2020 and May 2020, there was a significant decline in experiences of loneliness overall, with 36.1% of the sample saying that they experienced loneliness at least some of the time in May 2020, compared with 45.8% in April 2020 (Biddle et al. 2020c). Declines in loneliness were consistent by age and sex, with the exception of young people (aged 18-24) who did not have a statistically significant reduction in the level of loneliness from April to May 2020—despite the fact that they had the highest proportion of respondents saying that they felt lonely at least some of the time in April 2020 (63.3%) (Biddle et al. 2020c).

There was a rise in reported loneliness from 36.1% in May 2020 to 40.5% in August 2020 (Biddle et al. 2020d). However, this rise only occurred in Victoria where the proportion of the population who were lonely at least some of the time increased from 35.7% in May 2020 to 44.5% in August 2020; in 'the other seven States and Territories, there was no significant difference between loneliness in May 2020 (37.1%) and August 2020 (38.8%)' (Biddle et al. 2020d). The increase in loneliness from May 2020 to August 2020 in Victoria coincided with the lockdown associated with the second wave of COVID-19 infections.

Females were more likely to report experiencing loneliness than males (44.8% of females compared with 35.7% of males in August 2020). The proportion of young people aged 18-24 years who were experiencing loneliness in August 2020 was also higher than for other ages groups (Biddle et al. 2020d).

The proportion of Australians who said that they had experienced loneliness at least some of the time declined from 40.5% in August 2020 to 35.2% in November 2020 (Biddle et al. 2020e). This is the lowest value observed over the pandemic period so far. In addition, according to Biddle et al (2020e):

There has been a very large decline in the proportion of Australians who said that they 'never met socially with friends, relatives or work colleagues' since the early days of the pandemic. In April 2020, 49.4% of Australians said they 'never met socially'. This declined to 26.5% by May 2020, and even further to 6.8% in November 2020. While this is a dramatic change over a reasonably short period of time, the level of social isolation in November 2020 is still above the pre-pandemic level of 2.0 per cent [recorded in February 2020].

The proportion of Australians who said that they had experienced loneliness at least some of the time remained consistent from November 2020 (35.2%) to January 2021 (36.1%) and April 2021 (35.5%) and increased slightly in August 2021 (37.6%) (Biddle & Gray 2021a & c).

However, the increase in loneliness in August 2021 was mainly due to a large increase in Sydney, with the proportion of Sydney residents experiencing loneliness at least some of the time increasing from 35.3% in April 2021 to 44.3% in August 2021, while the rest of Australia reported little change in this period (34.6% in August 2021 compared with 34.0% in April 2021). At the time of the August 2021 survey, Sydney had been in lockdown for the longest amount of time during the COVID-19 Delta strain outbreak. The proportion of Australians who said that they had experienced loneliness at least some of the time increased in October 2021 (39.2%) (Biddle 2021b) and then decreased in January 2022 (38.2%) and April 2022 (36.5%) (Biddle 2022a,b) but remained above the lows from November 2020 to April 2021.



Loneliness has a clear impact on levels of psychological distress and life satisfaction. In a regression analysis of data from the November 2020 ANUpoll (that controlled for psychological distress in April 2020) those who felt lonely 'some', 'occasionally' or 'most' of the time all had significantly higher levels of psychological distress than others (Biddle et al 2020e). This suggests that reductions in loneliness may contribute to reductions in levels of psychological distress. Similar results are evident for life satisfaction - that is, after controlling for life satisfaction in April 2020, people who reported feeling lonely at least some of the time had significantly lower levels of life satisfaction than others.

As with average levels of psychological distress over the pandemic period, loneliness also differed by population group. Biddle et al. calculated an average loneliness score across the 10 ANUpoll survey waves from April 2020 to April 2022 (Biddle et al. 2022). Regression analysis was then undertaken to examine the factors associated with average loneliness over the pandemic period and differences between population groups. The regression model included a measure of social interaction from the February 2020 ANUpoll to control for pre-pandemic loneliness. The results of the model showed that during the pandemic period (Biddle et al. 2022):

- females were slightly more likely to experience loneliness than males, although the difference was not statistically significant
- respondents aged 45 and over had substantially lower levels of loneliness than those aged under 45
- those in the lowest income households had the highest levels of loneliness, compared with those in higher income households
- Victorians seemed to experience similar levels of loneliness as residents of New South Wales and South Australians experienced lower levels of loneliness compared with New South Wales (the base case)

The most recent ANUpolls show that the proportion of Australians' who were lonely at least some of the time during August 2022, October 2022, and January 2023 was 35.6%, 35.9%, and 35.4% respectively (Biddle 2022c; Biddle & Gray 2023; Biddle, McAllister and Sheppard 2022).

## Life satisfaction

Another way of tracking wellbeing is to analyse changes in life satisfaction. In the ANUpoll surveys life satisfaction is measured on a scale of 1 to 10, with higher scores indicating higher levels of satisfaction. Average life satisfaction scores fell substantially during the early stages of the pandemic from 6.9 in January 2020 to 6.5 in April 2020, before rising to 6.8 in May 2020 as infection rates fell and lockdown conditions started to be eased (Biddle et al. 2020d). The average level of satisfaction then fell to 6.6 in August 2020. However, between October and November 2020, life satisfaction improved substantially from an average score of 6.7 to 7.0. The average life satisfaction score was no longer significantly different to that recorded in October 2019 (7.1), and was slightly higher than that recorded during the Black Summer Bushfire crisis (6.9 in January 2020) (Biddle et al. 2020e).

Although the overall level of life satisfaction in November 2020 had returned to pre-pandemic levels (October 2019) there was a substantial reduction in life satisfaction scores during 2020. A regression analysis conducted in November 2020 using the longitudinal nature of the ANUpoll data, suggested that after controlling for the level of life satisfaction in January 2020, the total loss of life satisfaction over 2020 was:

- significantly higher for people living in Victoria compared with the rest of the Australian population
- lower for those aged 55 years and over compared with those aged 35-44
- lower for those who living outside the capital cities (Biddle et al. 2020e).

This is consistent with what you would expect given the greater impact of, among other things, lockdowns (through, for example, their impact on employment) for younger people, people in Victoria and people living in capital cities.

Nationally, the average life satisfaction score showed little change from November 2020 (7.0) to January 2021 (7.0), and then declined slightly to 6.9 in April 2021, although the difference was not statistically significant (Biddle & Gray 2021a,b).

In August 2021, the average life satisfaction score in Australia decreased substantially to 6.5, to a level similar to that reported in April 2020 at the peak of the first wave of COVID-19 in Australia and was lower than all other time points measured by ANUpoll (Biddle & Gray 2021c). In order to demonstrate the magnitude of the decline in life satisfaction between April and August 2021, Biddle & Gray (2021c) converted the decline into income equivalents, based on the relationship between life satisfaction and household income prior to the pandemic. This model estimated that the decrease in life satisfaction reported between April and August 2021 was equivalent to a loss of \$827 in average weekly household income (Biddle & Gray 2021c).

There was a slight increase in life satisfaction between August 2021 and October 2021, from 6.5 to 6.6. The October ANUpoll was conducted in mid-October when COVID-19 restrictions were beginning to be lifted. Life satisfaction remained at 6.6 in January 2022 and increased slightly to 6.7 in April 2022, still lower than the levels observed pre-pandemic and from November 2020 to April 2021 (Biddle 2022a,b, Biddle & Gray 2021d). Unlike psychological distress and loneliness, life satisfaction was included in the January 2020 ANUpoll, which can be used as a pre-pandemic baseline. Average life satisfaction over the pandemic period has mostly been lower than before the pandemic in January 2020 (noting that life satisfaction was lower in January 2020 when bushfires were affecting large parts of Eastern Australia than it was in October 2019) (Biddle et al. 2022). Biddle et al. calculated average loss of life satisfaction during the pandemic period for respondents to the April 2022 ANUpoll, using the January 2020 ANUpoll as the baseline and data from the ANUpoll surveys conducted between April 2020 and April 2022 (Biddle et al. 2022). The calculation assumes that had the COVID-19 pandemic not occurred, life satisfaction would have remained at January 2020 levels. The average level of 'lost life satisfaction' due to the pandemic for respondents to the April 2022 ANUpoll was 0.216, which when converted to income equivalents was roughly equivalent to a halving in income (Biddle et al. 2022).

The impact of the pandemic on life satisfaction differed by age, with the greatest declines among those aged 18 to 24 years, moderate declines for those aged 25 to 54, no significant declines for those aged 55 and over and slight increases for those aged 75 and over (Biddle et al. 2022).

Regression modelling was used to estimate the association between other demographic variables and loss of life satisfaction due to the pandemic. In this model, greater declines in life satisfaction were observed in respondents who had not completed year 12 compared with those who had as well as Victorians compared with residents of New South Wales. Respondents in the 4th and 5th income quintiles (the two highest income groups) had the smallest loss in life satisfaction, compared with other income quintiles (Biddle et al. 2022)

The most recent ANUpolls show that average Australian life satisfaction during August 2022, October 2022, and January 2023 was 6.8, 6.7, and 6.8 respectively (Biddle 2022c; Biddle & Gray 2023; Biddle, McAllister and Sheppard 2022). While life satisfaction has steadily increased since August 2021, it remains lower than November 2020 and pre-COVID-19 during October 2019 (p6., Biddle & Gray 2023).

## Data on deaths by suicide

There has been considerable commentary since the start of the pandemic on its potential to impact on the incidence of deaths by suicide. Much of this commentary has been based on modelling using previous experience including the relationship between unemployment and deaths by suicide. However, evidence to-date does not indicate an increase in suicide deaths in Australia during the pandemic. National mortality data published by the Australian Bureau of Statistics show that the rate of death by suicide in Australia was lower in 2020 (12.1 per 100,000 population) and 2021 (12.0) than in 2019 (13.1), see [Deaths by suicide over time](https://www.aihw.gov.au/suicide-self-harm/monitoring/data/deaths-by-suicide-in-australia/suicide-deaths-over-time) (<https://www.aihw.gov.au/suicide-self-harm/monitoring/data/deaths-by-suicide-in-australia/suicide-deaths-over-time>). That said, Australian Bureau of Statistics coding of psychosocial risk factors associated with deaths by suicide in 2020 determined that 3.2% of these deaths had the pandemic mentioned in either a police or pathology report or a coronial finding. In 2021, the percentage of suicide deaths where the pandemic was mentioned decreased to 2.6%. In most of these cases, other risk factors for suicide were also present. In 2021, the pandemic appeared to impact on people in different ways, including through job loss and financial insecurity as well as general concern or anxiety about societal changes or contacting the virus (ABS 2022a). For more information, visit [Psychosocial risk factors and deaths by suicide](#).

Internationally, a 'living systematic review' (John et al. 2021) based on evidence until 19 October 2020, has concluded that:

There was no consistent evidence of a rise in suicide but many studies noted adverse economic effects were evolving. There was evidence of a rise in community distress, fall in hospital presentation for suicidal behaviour and early evidence of an increased frequency of suicidal thoughts in those who had become infected with COVID-19. This living review provides a regular synthesis of the most up-to-date research evidence to guide public health and clinical policy to mitigate the impact of COVID-19 on suicide risk as the longer term impacts of the pandemic on suicide risk are researched.

A study investigating trends in suicide deaths between January 2017 and August 2020, using data from the Queensland, New South Wales, and Victorian suicide registers concluded that (Clapperton et al., 2021):

Although our analysis found no evidence of an overall increase in suicides after the pandemic began, the picture is complex. The identified increase in suicide in young men indicates that the impact of the pandemic is likely unevenly distributed across populations. The increase in suicides in the context of unemployment reinforces the vital need for mitigation measures during COVID-19, and for ongoing monitoring of suicide as the pandemic continues.

Since 2020, suicide registers in Victoria and New South Wales have regularly published data on suspected deaths by suicide for 2019, 2020 and 2021. The *Suicide in Queensland: Annual Report 2022* (Leske et al. 2022) included data on suspected deaths by suicide from the interim Queensland Suicide Register (iQSR) from 2019 to 2021. The number of suspected deaths by suicide recorded in suicide registers fluctuates from year to year and, while there have been increases and decreases in some jurisdictions in 2020 and 2021 compared with 2019, there is no evidence to-date that the pandemic has affected suicide rates. For more information see [Suspected deaths by suicide](#).

While the pandemic does not appear to have affected suicide rates in Queensland compared with previous reports, examination of police reports by the iQSR indicates that between 29 January 2020 and 31 December 2021 COVID-19 appeared to be a contributing factor in 86 of the 1,539 suspected deaths by suicide (5.6%) (Leske et al. 2022).

It is true that some key risk factors associated with deaths by suicide did worsen following the onset of the pandemic. For example, there were considerable job losses and rises in the level of psychological distress. On the other hand, it is possible that a general sense of 'we are all in this together' could have a protective impact. From February to April 2020 there were rises in the level of trust in others and in governments in Australia (Biddle et al. 2020a). In addition, the vast majority of people who experience unemployment or high levels of psychological distress or mental health issues will never experience a suicide attempt. That said, it is very important to monitor trends in risk factors and trends in deaths by suicide in real time.

A study undertaken by Leske et al (2021) used Queensland Suicide Register data to compare rates of suspected suicide before the COVID-19 pandemic (2015-2019) to rates of suspected suicide during the earlier months of the pandemic (February 2020-August 2020). Leske et al. (2021) found there was no overall change in the rates of suspected suicide during the first seven months of the pandemic in Queensland. Nonetheless, that COVID-19 had been a contributing factor for some individual suspected suicide deaths that occurred during the period.

Another factor that should be considered is the impact of both JobKeeper and the JobSeeker supplement. This is important given the association between the risk of dying by suicide and socioeconomic outcomes. Modelling undertaken by the ANU suggests that not only were levels of poverty and housing stress lower than they otherwise would have been as a result of these payments, they were also lower than they were prior to the spread of COVID-19 (Philips et al. 2020). Households who mainly relied on the JobSeeker payment prior to the pandemic and the introduction of the JobSeeker supplement saw their poverty rate fall from 67% prior to COVID-19 to 6.8% (Phillips et al. 2020). On a similar note Biddle et al. (2020d) found that real incomes actually rose for those in the bottom decile of the income distribution from February to August 2020. Using data from the Taking the Pulse of the Nation Survey, Botha et al. (2020) have shown that the level of

psychological distress among the unemployed declined after May 2020. The ANU modelling suggests that the protective impact of JobKeeper and the JobSeeker supplement on housing stress and poverty were reduced somewhat by the changes to these payments announced in July 2020 (Phillips et al. 2020).

## Ambulance attendances


A key part of the National Suicide and Self-harm Monitoring Project is the compilation and coding of data from ambulance attendances. The National Ambulance Surveillance System (NASS), established in 2020, provides data on ambulance attendances for suicidal and self-harm behaviours for New South Wales, Victoria, Queensland, Tasmania and the Australian Capital Territory. The NASS is a partnership between Turning Point, Monash University and jurisdictional ambulance services across Australia. Comprehensive data from the NASS are reported in [Ambulance attendances: suicidal and self-harm behaviours](#).

In 2020, prior to the establishment of the NASS, the AIHW requested Turning Point prioritise the coding of data for Victoria. Victoria was chosen as monthly data on ambulance attendances were already being compiled for the Victorian Government and there were concerns about the impact of the Melbourne lockdown on suicide and self-harm behaviours.

Monthly data on the number of ambulance attendances related to suicide attempts in Victoria from 2018 to 2020 are shown in Figure 4. As the figure shows, there is no clear difference from 2018 to 2020. The total number of ambulance attendances related to suicide attempts in Victoria in 2020 was 4% lower than in 2019 but 8% higher than in 2018.

In 2020, there was an 11% increase in the total number of ambulance attendances in Victoria relating to suicidal ideation (thinking about suicide), compared with 2019, with the increase more pronounced in the second half of 2020. There was also an increase in the total number of mental health attendances (16% higher in 2020 than in 2019). This is consistent with the overall greater use of mental health services in 2020 that is evident in other data. This highlights the fact that greater use of, and need for, mental health services does not necessarily equate to trends in the number of suicide attempts. The vast bulk of people who use mental health services will never have a suicide attempt but timely access to mental health services may reduce the number of deaths by suicide. The total number of ambulance attendances for self-injury in Victoria in 2020 was considerably higher (33%) than in 2019. This highlights the fact that self-injury and suicide attempts are not the same thing.

**Figure 4: Monthly ambulance attendances for suicide attempts, Victoria, 2018 to 2020**



## References

AAPOR (American Association for Public Opinion Research) (2010) *Report on Online Panels* (<https://www.aapor.org/Education-Resources/Reports/Report-on-Online-Panels.aspx>), accessed 16 November 2022.

ABS (Australian Bureau of Statistics) (2022a) *Causes of Death, Australia 2021* (<https://www.abs.gov.au/statistics/health/causes-death/causes-death-australia/latest-release>), ABS website, accessed 4 November 2022.

ABS (2022b) *National Study of Mental Health and Wellbeing* (<https://www.abs.gov.au/statistics/health/mental-health/national-study-mental-health-and-wellbeing/latest-release>), ABS website, accessed 15 February 2023.

ABS (2022c) *National, State and Territory Population* (<https://www.abs.gov.au/statistics/people/population/national-state-and-territory-population/jun-2022>), ABS website, accessed 16 February 2023.

ABS (2021) *First insights from the National Study of Mental Health and Wellbeing, 2020-21* ([https://www.abs.gov.au/articles/first-insights-national-study-mental-health-and-wellbeing-2020-21#:~:text=In%202020%2D21%2C%2015%25,high%20levels%20of%20psychological%20distress.&text=one%20in%20five%20\(20%25\),%2D85%20years%20\(9%25\).](https://www.abs.gov.au/articles/first-insights-national-study-mental-health-and-wellbeing-2020-21#:~:text=In%202020%2D21%2C%2015%25,high%20levels%20of%20psychological%20distress.&text=one%20in%20five%20(20%25),%2D85%20years%20(9%25).)), ABS website, accessed 16 November 2022.

AIHW (Australian Institute of Health and Welfare) (2020) *National Drug Strategy Household Survey 2019*, AIHW, Australian Government, accessed 16 November 2022.

Biddle, N (2021a) *ANU Poll 2020-2021 longitudinal* (<https://dataverse.ada.edu.au/dataset.xhtml?persistentId=doi:10.26193/JGW1LD>) [dataset], dataverse.ada.edu.au, doi:10.26193/JGW1LD, accessed 23 March 2023.

Biddle N (2021b) *ANU Poll 48 (October 2021): Mental health and economic stress during COVID-19* (<https://dataverse.ada.edu.au/dataset.xhtml?persistentId=doi:10.26193/THF1VZ>) [data set], dataverse.ada.edu.au, accessed 16 November 2022.

Biddle N (2022a) *ANU Poll 49 (January 2022): Experiences with COVID-19* (<https://dataverse.ada.edu.au/dataset.xhtml?persistentId=doi:10.26193/2MX3D0>) [data set], dataverse.ada.edu.au, accessed 16 November 2022.

Biddle N (2022b) *ANU Poll 50 (April 2022): Volunteering, aged care, policy priorities and experiences with COVID-19* (<https://dataverse.ada.edu.au/dataset.xhtml?persistentId=doi:10.26193/AXQPSE>) [data set], dataverse.ada.edu.au, accessed 16 November 2022.

Biddle N (2022c) *ANU Poll 53 (Oct 2022): COVID-19, mental health, employment, data privacy and support for randomised controlled trials* (<https://dataverse.ada.edu.au/dataset.xhtml?persistentId=doi:10.26193/WBJE1K>) [data set], dataverse.ada.edu.au, accessed 23 February 2023.

Biddle N and Gray M (2020) *Tracking outcomes during the COVID-19 pandemic (October 2020) - Reconvergence* (<https://csm.cass.anu.edu.au/research/publications/tracking-outcomes-during-covid-19-pandemic-october-2020-reconvergence>), COVID-19 Briefing Paper, ANU Centre for Social Research and Methods, Australian National University, Canberra.

Biddle N and Gray M (2021a) *Tracking outcomes during the COVID-19 pandemic (January 2021) - Cautious Optimism* (<https://csm.cass.anu.edu.au/research/publications/tracking-outcomes-during-covid-19-pandemic-january-2021-cautious-optimism>), COVID-19 Briefing Paper, ANU Centre for Social Research and Methods, Australian National University, Canberra.

Biddle N and Gray M (2021b) *Tracking wellbeing outcomes during the COVID-19 pandemic (April 2021) - Continued social and economic recovery and resilience* (<https://csm.cass.anu.edu.au/research/publications/tracking-wellbeing-outcomes-during-covid-19-pandemic-april-2021-continued>), COVID-19 Briefing Paper, ANU Centre for Social Research and Methods, Australian National University, Canberra.

Biddle N and Gray M (2021c) *Tracking wellbeing outcomes during the COVID-19 pandemic (August 2021) - Lockdown blues* (<https://csm.cass.anu.edu.au/research/publications/tracking-wellbeing-outcomes-during-covid-19-pandemic-august-2021-lockdown>), COVID-19 Briefing Paper, ANU Centre for Social Research and Methods, Australian National University, Canberra.

Biddle N and Gray M (2021d) *Tracking wellbeing outcomes during the COVID-19 pandemic (October 2021) - Putting the worst behind us?* (<https://csm.cass.anu.edu.au/research/publications/tracking-wellbeing-outcomes-during-covid-19-pandemic-october-2021-putting>), COVID-19 Briefing Paper, ANU Centre for Social Research and Methods, Australian National University, Canberra.

Biddle N and Gray M (2022) *Tracking wellbeing outcomes during the COVID-19 pandemic (January 2022) - Riding the Omicron wave* (<https://csm.cass.anu.edu.au/research/publications/tracking-wellbeing-outcomes-during-covid-19-pandemic-january-2022-riding>), COVID-19 Briefing Paper, ANU Centre for Social Research and Methods, Australian National University, Canberra.

Biddle N and Gray M (2023) *Taking stock: Wellbeing and political attitudes in Australia at the start of the post-COVID era, January 2023* (<https://csm.cass.anu.edu.au/research/publications/taking-stock-wellbeing-and-political-attitudes-australia-start-post-covid-era>), COVID-19 Briefing Paper, ANU Centre for Social Research and Methods, Australian National University, Canberra.

Biddle N, Gray M and Rehill P (2022) *Mental health and wellbeing during the COVID-19 period in Australia* (<https://csm.cass.anu.edu.au/research/publications/mental-health-and-wellbeing-during-covid-19-period-australia>), ANU Centre for Social Research and Methods, Australian National University, Canberra.

Biddle N, Edwards B, Gray M and Sollis K (2020a) *Hardship, distress, and resilience: The initial impacts of COVID-19 in Australia* (<https://csm.cass.anu.edu.au/research/publications/hardship-distress-and-resilience-initial-impacts-covid-19-australia-1>), ANU Centre for Social Research and Methods, Australian National University, Canberra.

Biddle N, Edwards B, Gray M and Sollis K (2020b) *Initial impacts of COVID-19 on mental health in Australia (22 May 2020)* ([https://csm.cass.anu.edu.au/sites/default/files/docs/2020/6/Mental health before and during the COVID crisis.pdf](https://csm.cass.anu.edu.au/sites/default/files/docs/2020/6/Mental%20health%20before%20and%20during%20the%20COVID%20crisis.pdf)), COVID-19 Briefing Paper, ANU Centre for Social Research and Methods, Australian National University, Canberra.

Biddle N, Edwards B, Gray M and Sollis K (2020c) *Mental health and relationships during the COVID-19 pandemic (July 2020)* (<https://csm.cass.anu.edu.au/research/publications/mental-health-and-relationships-during-covid-19-pandemic>), COVID-19 Briefing Paper, ANU Centre for Social Research and Methods, Australian National University, Canberra.

Biddle N, Edwards B, Gray M and Sollis K (2020d) *Tracking outcomes during the COVID-19 pandemic (August 2020) - Divergence within Australia* (<https://csm.cass.anu.edu.au/research/publications/tracking-outcomes-during-covid-19-pandemic-august-2020-divergence-within>), COVID-19 Briefing Paper, ANU Centre for Social Research and Methods, Australian National University, Canberra.

Biddle N, Edwards B, Gray M and Sollis K (2020e) *Tracking outcomes during the COVID-19 pandemic (November 2020) - Counting the Cost of the COVID-recession* (<https://csm.cass.anu.edu.au/research/publications/tracking-outcomes-during-covid-19-pandemic-november-2020-counting-costs-covid>), COVID-19 Briefing Paper, ANU Centre for Social Research and Methods, Australian National University, Canberra.

Biddle, N, McAllister, I, and Sheppard, J (2022) *ANU Poll 52 (August 2022): COVID-19, mental health, population issues, data privacy and coercive control* (<https://dataverse.ada.edu.au/dataset.xhtml?persistentId=doi:10.26193/FCZGOK>) [dataset], dataverse.ada.edu.au, doi:10.26193/FCZGOK, accessed 23 March 2023.

Botha F, Butterworth P and Wilkins R (2020) *Mental distress in Australia over the course of the first wave of the COVID-19 Pandemic* (<https://melbourneinstitute.unimelb.edu.au/publications/research-insights/search/result?paper=3491818>), Melbourne Institute Research Insights 24/20, Melbourne Institute: Applied Economic & Social Research, Melbourne.

Clapperton, A., Spittal, M. J., Dwyer, J., Garrett, A., Kölves, K., Leske, S., Millar, C., Edwards, B., Stojcevski, V., Crompton, D. R., & Pirkis, J. (2021). 'Patterns of suicide in the context of COVID-19: Evidence from three Australian States. *Frontiers in Psychiatry*, 12. <https://www.frontiersin.org/articles/10.3389/fpsy.2021.797601>.

John A, Eyles E, Webb RT, Okolie C, Schmidt L, Arensman E, Hawton K, O'Connor RC, Kapur N, Moran P, O'Neill S, McGuinness LA, Olorisade BK, Dekel D, Macleod-Hall C, Cheng HY, Higgins JPT and Gunnell D (2021) 'The impact of the COVID-19 pandemic on self-harm and suicidal behaviour: update of a living systematic review' [version 2; peer review: 1 approved, 2 approved with reservations] *F1000Research*, 9:1097, doi: 10.12688/f1000research.25522.2.

Leske S, Adam G, Catakovic A, Weir B, and Kölves K (2022) *Suicide in Queensland: Annual Report 2022* ([https://www.griffith.edu.au/\\_data/assets/pdf\\_file/0033/1639473/AISRAP-Annual-Report-2022.pdf](https://www.griffith.edu.au/_data/assets/pdf_file/0033/1639473/AISRAP-Annual-Report-2022.pdf)). Australian Institute for Suicide Research and Prevention, Griffith University, Brisbane.

Leske, S., Kölves, K., Crompton, D., Arensman, E., & de Leo, D. (2021). Real-time suicide mortality data from police reports in Queensland, Australia, during the COVID-19 pandemic: An interrupted time-series analysis. *The Lancet Psychiatry*, 8(1), 58-63. [https://doi.org/10.1016/S2215-0366\(20\)30435-1](https://doi.org/10.1016/S2215-0366(20)30435-1).

Phillips B, Gray M and Biddle N 2020 *JobKeeper and JobSeeker impacts on poverty and housing stress under current and alternative economic and policy scenarios* (<https://csm.cass.anu.edu.au/research/publications/covid-19-jobkeeper-and-jobseeker-impacts-poverty-and-housing-stress-under>), ANU Centre for Social Research and Methods, Australian National University, Canberra

Prochaska J, Sung H-Y, Max W, Shi Y and Ong M (2012) 'Validity study of the K6 scales as a measure of moderate mental distress based on mental health treatment need and utilization', *International Journal of Methods in Psychiatric Research*, 21(2): 88-97, doi: 10.1002/mpr.1349.

Slade T, Grove R and Burgess P (2011) 'Kessler Psychological Distress Scale: Normative data from the 2007 Australian National Survey of Mental Health and Wellbeing', *Australian & New Zealand Journal of Psychiatry*, 45(4):308-316, doi:10.3109/00048674.2010.543653.



## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on [13 11 14](tel:131114), or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)

## Suicide & self-harm monitoring: Geography

Reporting deaths by suicide and hospitalisations for intentional self-harm at smaller, more ‘localised’ geographical areas, can reveal information that may be masked by reporting for the whole of Australia or by states and territories—allowing for a better understanding of suicidal behaviours for local communities, policymakers and researchers.

Although suicide has a significant impact on the community, it is a relatively rare cause of death in Australia meaning that depending on the level of geography considered, there may be areas where there are very few—or even no—deaths by suicide recorded in a given year. The number of hospitalisations for intentional self-harm are approximately 10 times that of deaths by suicide; however, further disaggregation (or breakdown) of the data by age or sex reduces the numbers of events able to be reported for each group in each small geographical area in a single year. Strict privacy and confidentiality controls or concerns regarding statistical reliability mean that small numbers (or rates based on them) cannot be publicly reported, thereby reducing the coverage of reportable data as smaller geographical areas are considered.

Numbers and age-standardised rates (where they could be reliably calculated) of deaths by suicide and hospitalisations for intentional self-harm have been reported by PHN area and Statistical Areas level 3 and 4. For the reporting of suicide and hospitalised intentional self-harm data by Statistical Area, the smallest possible geographical area has been used while still allowing for maximum coverage of reportable data across these small geographical areas.

This section also contains global statistics on suicide—intended to provide a broad view of the issue across the world.

---

## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on [13 11 14](tel:131114), or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)

## Australian Youth Self-Harm Atlas

The Australian Youth Self-Harm Atlas study investigated regional variability in suicidality and self-harm, as well as risk and protective factors, for young people aged 12 to 17 years. Aspects of the quantitative component of the study are presented here. The full [Australian Youth Self-Harm Atlas: Summary Report](#) (<https://espace.library.uq.edu.au/view/UQ:2090e31>) (Hielscher et al., 2022), includes a summary of both quantitative and qualitative study components.

### Strengthening suicide prevention

*The National Mental Health and Suicide Prevention Agreement* (Commonwealth of Australia, 2022) identifies the importance of strengthening regional planning and evaluation of suicide prevention initiatives. To do this, detailed regional data are needed.

The Australian Youth Self-Harm Atlas study:

- Is the first national Australian study to estimate the variability of youth self-harm and suicidality, across small areas of geography (Hielscher et al., 2022).
- Distinguishes between self-harm without suicidal intent, suicidal ideation/planning, and suicide attempt. This differentiation has service and program planning implications but is not often available within administrative datasets.
- Data are representative of whole communities, rather than being limited to the experience of those using hospital (or other healthcare) services.

While identifying communities whose residents are not faring as well as others may be seen as stigmatising, the purpose for doing so is to provide evidence upon which community members and decision-makers can rely.

## About the Study

### Data sources

The Australia Youth Self-Harm Atlas study generated synthetic estimates of youth suicidality and self-harm using:

- [Young Minds Matter \(YMM\) survey](#) (<https://youngmindsmatter.telethonkids.org.au/>). A nationally representative household survey about the health and wellbeing of children and young people conducted between 2013 and 2014.
- [2016 Census](https://www.abs.gov.au/websitedbs/censushome.nsf/home/2016) (<https://www.abs.gov.au/websitedbs/censushome.nsf/home/2016>).
- 2019 Australian Bureau of Statistics [Estimated Resident Population](https://www.abs.gov.au/methodologies/national-state-and-territory-population-methodology/sep-2022) (<https://www.abs.gov.au/methodologies/national-state-and-territory-population-methodology/sep-2022>) data.

### Generating synthetic estimates

The Australian Youth Self-Harm Atlas study generated synthetic estimates to enable measurement of suicidality and self-harm prevalence, and related risk and protective factors for small areas.

The Young Minds Matter (YMM) survey data holds information about suicidality and self-harm among young people that completed the survey. While the sampling strategy used for the survey was scientifically robust, not every community across Australia was invited to participate (Hafekost et al., 2016). As such, it is not possible to use YMM data to directly measure suicidality and self-harm among the young people within each community across Australia. To solve this problem, small area estimation methods were used to produce synthetic prevalence estimates of youth suicidality and self-harm for Statistical Area level 1s (SA1) across Australia. SA1s are a standardised measure

of geography and part of the [Main Structure of the Australian Statistical Geography Standard](https://www.abs.gov.au/statistics/standards/australian-statistical-geography-standard-asgs-edition-3/jul2021-jun2026/main-structure-and-greater-capital-city-statistical-areas) (ASGS), developed by the Australian Bureau of Statistics. SA1s generally have a population of 200 to 800 people, and an average population of about 400 people.

The small area estimation undertaken involved linking Young Minds Matter survey data with 2016 Census data. Noting that 2016 Census data are available for all SA1 areas, whereas the survey data are only available for those SA1 areas that were invited to participate. Patterns in responding for those who completed both the Young Minds Matter (YMM) survey and the 2016 Census were then used to extrapolate responses to the youth suicidality and self-harm YMM survey questions for communities that were not actually invited to complete the survey. Data generated in this way, using sophisticated statistical models, are referred to as synthetic estimates. Synthetic estimates generated for SA1 areas were then summed together and presented at broader areas of geography. Synthetic estimates were presented in this publication at SA3, SA4 and Primary Health Network (PHN) areas.

Australian Bureau of Statistics Estimated Resident Population data for 2019 were used to calculate suicidality and self-harm prevalence estimates for geographic areas.

As a means of external validation, synthetic suicidality and self-harm prevalence estimates were compared to rates of death by suicide. At an SA2 level, each of the suicidality and self-harm measures used within the study were positively correlated with the average annual rate of death by suicide between 2010-2019.

Synthetic estimates based on small numbers of young people were suppressed to maintain confidentiality and avoid publishing statistics of low reliability.

#### The Australian Youth Self-Harm Atlas study includes the following suicidality and self-harm outcomes:

- Self-harm (regardless of intent): self-injurious behaviour irrespective of intent or motivation, including behaviours with either suicidal or non-suicidal intent, or where intent is ambiguous. This was the primary outcome of this study (inclusive of non-suicidal self-harm and suicide attempt behaviour).
- Non-suicidal self-harm: self-injurious behaviour for which there is evidence that the person did not intend to kill themselves.
- Suicidal ideation/plans: thoughts of engaging in or planning suicide-related behaviour; without engaging in suicidal behaviour.
- Suicide attempt: non-fatal, self-directed, potentially self-injurious behaviours with an intent to die.
- Suicidality: suicidal thoughts or behaviours, including ideation, plans, and attempts.

#### Study limitations and important data considerations

The information provided by the Australian Youth Self-Harm Atlas Study may be the best available small area data for youth suicidality and self-harm.

Even so, there are important limitations to consider included within the '[Study limitations and important data considerations](https://aihw.gov.au/suicide-self-harm-monitoring/data/geography/youth-self-harm-atlas/limitations-and-interpretation-considerations)' sections of this publication.

#### The study team

The Australian Youth Self-Harm Atlas study was undertaken by a team of researchers and clinicians, and at the heart of the project was a partnership between Queensland Institute of Medical Research (QIMR) Berghofer and Roses in the Ocean. Roses in the Ocean is a lead Australian organisation for lived experience of suicide.

The AIHW has worked in collaboration with Youth Self-Harm Atlas study authors, Dr Emily Hielscher (formerly of Queensland Institute of Medical Research (QIMR) Berghofer) and Professor David Lawrence (Curtin University), to integrate quantitative findings of the study into the AIHW Suicide and Self-Harm Monitoring website.





## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on [13 11 14](tel:131114), or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)

## Deaths by suicide by remoteness areas

About 28% of the Australian population live in regional and remote areas—areas outside Australia’s major cities. There are many positive aspects about living in regional and remote areas, including higher levels of life satisfaction compared with those in urban areas (Wilkins 2015), increased community interconnectedness and social cohesion, and higher levels of community participation, volunteering and informal support from their communities (Ziersch et al. 2009). However, Australians living in these areas face unique challenges due to their geographic isolation, and often have poorer health and welfare outcomes than those living in major cities.

For further information on how the statistics reported here were calculated see [Technical notes](#).


Suicide deaths by remoteness area, Australia, 2010 to 2022.

The line graph shows the age-standardised rates of suicide for Very Remote, Remote, Outer Regional and Inner Regional areas and Major Cities from 2010 to 2022. Users can also choose to view age-standardised rates and numbers of deaths by suicide for remoteness areas by sex.

## Suicide deaths by sex and remoteness areas, Australia, 2001 to 2022

Persons

Age-standardised rate (per 100,000)



Select remoteness area: Major Cities Inner Regional Outer Regional Remote Very Remote

Select measure:

Age-standardised rate (per 100,000)  
Number

Select sex:

Females  
Males  
Persons

n.p. Not published

Source: AIHW National Mortality Database and ABS Causes of Death, Australia 2023

Supplementary table: NMD S8

Latest data: 2022 (annual release)

See notes ►

## Are people in regional and remote areas at greater risk of deaths by suicide?

From 2001 to 2022:

- numbers of deaths by suicide were highest in *Major Cities* and fell as remoteness increased, while age-standardised suicide rates tended to increase with the increasing remoteness
- suicide rates for residents of *Major Cities* were the lowest of all 5 remoteness areas each year and remained relatively stable over the period (ranging from 9.2 deaths per 100,000 population in 2006 to 11.7 in 2017)
- suicide rates in *Very Remote* areas fluctuated between 2001 and 2022, with the lowest rate recorded in 2003 (17.6 deaths per 100,000 population) and the highest in 2007 and 2016 (29.1). The rate of suicide in *Very Remote* areas was 27.3 in 2001 and 24.5 in 2022. Fluctuations in rates are due largely to the small population and small numbers of deaths by suicide in these areas
- suicide rates for residents of *Remote* areas also fluctuated over the period, ranging from 10.6 per 100,000 population in 2009 to 23.7 in 2022
- suicide rates in *Inner Regional* and *Outer Regional* areas were generally higher in the second half of the period, with lows of 11.4 and 12.1 per 100,000 population, respectively, in 2006 and highs of 17.1 and 20.1, respectively, in 2019
- the greatest proportion of deaths by suicide occurred in *Major Cities* and remained relatively stable at 61.3-66.7% over the period.

In 2022:

- the age-standardised suicide rate for residents of *Major Cities* (10.5 deaths per 100,000 population) was lower than the national rate of 12.3 deaths per 100,000 population
- rates for residents of all other remoteness areas were above the national rate
- the rate for residents of *Very Remote* areas (24.5 deaths per 100,000 population) was 2.3 times that of the rate for residents of *Major Cities* (10.5 deaths per 100,000 population); however, numbers of deaths were relatively small (46 deaths in *Very Remote* areas and 2,030 in *Major Cities*)
- the proportion of deaths by suicide occurring in *Major Cities* in 2022 was 63.1%.

## References

Wilkins R 2015. [The Household, Income and Labour Dynamics in Australia Survey: selected findings from waves 1 to 12.](https://melbourneinstitute.unimelb.edu.au/data/assets/pdf_file/0006/2155506/hilda-statreport-2015.pdf)

([https://melbourneinstitute.unimelb.edu.au/data/assets/pdf\\_file/0006/2155506/hilda-statreport-2015.pdf](https://melbourneinstitute.unimelb.edu.au/data/assets/pdf_file/0006/2155506/hilda-statreport-2015.pdf)) Melbourne: Melbourne Institute of Applied Economic and Social Research.

Ziersch A, Baum F, Darmawan I, Kavanagh A & Bentley, R 2009. [Social capital and health in rural and urban communities South Australia.](https://www.ncbi.nlm.nih.gov/pubmed/19236353)

(<https://www.ncbi.nlm.nih.gov/pubmed/19236353>) Australian and New Zealand journal of public health 33:7-16. Adelaide: Flinders University.



## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on [13 11 14](tel:131114), or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)

## Intentional self-harm hospitalisations by remoteness areas

Hospitalisations data for patients with intentional self-harm injuries includes those with and without suicidal intent. For further information see the [Technical notes](#).

Understanding the geographical distribution of hospitalisations due to intentional self-harm based on patients' area of usual residence (see [Technical notes](#) for more information) can help target suicide prevention activities to areas in need.

Intentional self-harm hospitalisations, by age and remoteness areas, 2012-2013 to 2020-22.

The line graph shows age-specific rates of intentional self-harm hospitalisations for Very Remote, Remote, Outer Regional, Inner Regional, Major Cities and Total remoteness areas for all ages combined from 2012-13 to 2020-22. Users can also choose to view age-specific rate, numbers and proportion of hospitalisations for intentional self-harm by remoteness area and specific age groups.

### Are people in regional and remote areas at greater risk of intentional self-harm hospitalisations?

In 2021-22:

- residents of *Very Remote* areas recorded a rate of 193 hospitalisations per 100,000 population, nearly twice that of residents of *Major cities*, which recorded the lowest rate (97 hospitalisations per 100,000 population)
- two-thirds of intentional self-harm hospitalisations were residents of *Major cities* (66%)
- young people aged 15-19 had the highest rates of intentional self-harm hospitalisations in each remoteness area
- the highest rate of intentional self-harm hospitalisations overall was in the 15-19 age group in *Remote* areas (608 hospitalisations per 100,000 population), followed by the same age group in *Outer Regional* areas (566).

A similar pattern was seen with deaths by suicide as age-standardised suicide rates tended to increase with remoteness of place of residence [Suicide by remoteness areas](#).

### How have rates of intentional self-harm hospitalisations changed for remoteness areas?

Between 2012-13 and 2021-22:

- overall rates of intentional self-harm hospitalisations increased in *Very Remote* areas (from 172 to 193 hospitalisations per 100,000 population) and *Remote* areas (from 146 per 100,000 population to 160)
- rates fell in *Inner Regional* areas (from 125 to 103), and *Major Cities* (111 to 97) over this period while rates initially increased in *Outer Regional* areas from 136 hospitalisations per 100,000 population in 2012-13 to 174 in 2016-17, and then steadily decreased to 133 in 2021-22
- the highest increases in rates of intentional self-harm hospitalisations occurred in those aged 15-19 in *Outer Regional*, *Remote* and *Very remote* areas (367 hospitalisations per 100,000 population in 2012-13 to 566 in 2021-22, 465 to 608 and 428 to 538 hospitalisations per 100,000 population, respectively). Outside of the 15-19-year-old age group, the highest increase since 2012-13 was among the 50-54-year-old age group in *Remote* areas (85 hospitalisations per 100,000 population in 2012-13 to 201 in 2021-22)
- the largest decrease in rates of intentional self-harm hospitalisations was among the 35-39-year-old age group in *Outer Regional* areas (from 256 hospitalisations per 100,000 population in 2012-13 to 147 in 2021-22).





## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on [13 11 14](tel:131114), or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)

## Deaths by suicide, by Primary Health Network areas

Where people live can impact on their risk of suicide and also their access to services. Reporting rates or numbers of deaths by suicide at Primary Health Network (PHN) areas allows for more localised information that may provide a better understanding of the incidence of deaths by suicide in the local community and allow clinicians, policymakers and researchers to better plan services or suicide prevention activities.


PHNs are organisations that connect health services across a specific geographic area (PHN areas). There are 31 PHN areas that cover the whole of Australia with the [boundaries defined by the Australian Government Department of Health](#) (<https://www1.health.gov.au/internet/main/publishing.nsf/content/phn-maps-aust>). For further information on how the statistics reported here were calculated see [Technical notes](#).

Suicide deaths by Primary Health Network areas, Australia, 2010 to 2022.

The line graph shows the age-standardised rates of suicide for Australia by Primary Health Network (PHN) area from 2010 to 2022. Users can choose to view age-standardised rates and numbers of deaths by suicide by selected PHN.

## Suicide deaths by Primary Health Network areas, Australia, 2001 to 2022

Age-standardised rate (per 100,000)



\*Age-standardised rates are not shown where there are fewer than 20 deaths in an area due to unstable rates.  
 Source: AIHW National Mortality Database and ABS Causes of Death, Australia 2023  
 Supplementary Table: NMD S9  
 Latest data: 2022 (annual release)

[See notes ►](#)

### How do suicide rates vary among PHN areas?

In 2022:

- Age-standardised rates and numbers of deaths by suicide varied across PHN areas, ranging from 7.4 deaths per 100,000 population in Northern Sydney PHN area to 20.1 in the NT Northern Territory PHN area.
- The greatest number of deaths by suicide occurred in the North Western Melbourne PHN (190).

Data are not published for PHN areas where there are small numbers of deaths by suicide due to privacy and confidentiality concerns or other concerns about the quality of the data (for example, age-standardised rates cannot be published for Western Queensland for most years).



## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on [13 11 14](tel:131114), or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)

## Intentional self-harm hospitalisations by Primary Health Network areas

Hospitalisations data for patients with intentional self-harm injuries includes those with and without suicidal intent. For further information see the [Technical notes](#).

The reporting of rates of intentional self-harm hospitalisations by Primary Health Network (PHN) areas can provide localised information to enable PHNs to identify and investigate areas requiring more coordination of care to patients, by working directly with key primary and secondary health care providers and hospitals.

Intentional self-harm hospitalisations, by age and sex, by Primary Health Network areas, Australia, 2020-22.

The distribution plot shows the age-specific rates of intentional self-harm hospitalisations for males and females by all ages and broad age groups (0-24, 25-44, 45-64, 65 and over) for Primary Health Networks (PHNs) in 2020-22. Users can also choose to view horizontal stacked bar charts showing numbers and proportion of intentional self-harm hospitalisations for PHNs by all ages and age groups by sex.



## Intentional self-harm hospitalisations, by age, sex and Primary Health Network (PHN) areas, Australia, 2021-22

New South Wales  
Age-specific rate (per 100,000)



Note: Some data are not shown for age groups 45-64 and 65+ for selected PHNs due to small cell sizes.  
Source: AIHW National Hospital Morbidity Database  
Supplementary Table: NHMD S7  
Latest data: 2021-22 (annual release)

See notes ►

### How do rates of intentional self-harm hospitalisations vary across PHN areas?

The rates of hospitalisations for intentional self-harm in 2021-22 varied greatly by PHN area:

- the Northern Territory PHN area had the highest rate (238 hospitalisations per 100,000 population), while South Western Sydney PHN area had the lowest rate (44) (Supplementary table: [2021-22 National Hospital Morbidity Database - Intentional self-harm hospitalisations](https://www.aihw.gov.au/suicide-self-harm-monitoring/data/data-downloads) (<https://www.aihw.gov.au/suicide-self-harm-monitoring/data/data-downloads>), NHMD S7)
- The rate per 100,000 population for Australia was 105.

In 2021-22, rates of intentional self-harm hospitalisations for females and males were generally highest in younger age groups (those aged 24 and below and 25-44 respectively).

- the highest rate of intentional self-harm hospitalisations was among females aged 24 and below in the Western Queensland PHN area (507 per 100,000 population; 53 hospitalisations). This was also followed by those aged 24 and below, in the Country SA PHN (393 per 100,000 population; 273 hospitalisations)
- rates of intentional self-harm hospitalisation for males tended to be highest in those aged 25-44. The Northern Territory PHN area reported the highest rate for males in the 25-44 age group (242 per 100,000 population; 102 hospitalisations) followed by the Western Queensland PHN (241 per 100,000 population; 21 hospitalisations).

## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on [13 11 14](tel:131114), or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)

## Deaths by suicide, by local areas

Suicide incidence data in local communities provide insight into small populations and the variability of suicide rates across Australia. This is particularly pertinent for suicide prevention activities.

Deaths by suicide data have been aggregated (pooled) across five 5-year periods (2014-18, 2015-2019, 2016-2020, 2017-2021, and 2018-2022). Data are provided at Statistical Area Level 3 (SA3s) and Statistical Area Level 4 (SA4s). Data at SA4s are further disaggregated by sex.

SA3s and SA4s are a standardised measure of geography and part of the [Main Structure of the Australian Statistical Geography Standard](https://www.abs.gov.au/statistics/standards/australian-statistical-geography-standard-asgs-edition-3/jul2021-jun2026/main-structure-and-greater-capital-city-statistical-areas) (<https://www.abs.gov.au/statistics/standards/australian-statistical-geography-standard-asgs-edition-3/jul2021-jun2026/main-structure-and-greater-capital-city-statistical-areas>), developed by the Australian Bureau of Statistics. All data are presented by year of registration, which is not necessarily the same as the year the death occurred. For more information, see [Technical notes](#).

Direct estimates of suicide rates based on small numbers can be highly variable from year to year. As such, age standardised rates based on 20 or fewer deaths over the 5-year period have not been reported. Additionally, some areas with small numbers of deaths have had a random number of deaths assigned to them (instead of the true number) to protect the confidentiality of individuals. See [Technical notes](#) to ensure the data are interpreted appropriately.

### How to use these maps

Use the zoom and search functions to explore the map. Click on an area in the map to view additional information. The colour shading indicates different rates of deaths by suicide, with darker shades indicating a higher rate.

For the best experience, use Chrome, Edge or Firefox browsers. For more information on browser compatibility, see [Supported browsers](#). (<https://doc.arcgis.com/en/web-appbuilder/create-apps/supported-browsers.htm>)

### Suicide by local areas



(Map will open in a new window)

(<https://maps.arcgis.aihw.gov.au/portal/apps/experiencebuilder/experience/?id=47b5b317d1074ad09d93e861a6ebf55>)

Over the 5-year period 2018-2022, reportable age-standardised suicide rates in persons at the SA3 level, were:

- highest in the SA3 areas of Kimberley in Western Australia (32.9 deaths per 100,000 population), Burnett in Queensland (32.3) and Daly - Tiwi - West Arnhem in Northern Territory (31.4)
- lowest in the SA3 areas of Baulkham Hills in New South Wales (5.3 deaths per 100,000 population) Canterbury in New South Wales (5.5) and Tullamarine - Broadmeadows in Victoria (5.8).

Over the same period (2018-2022), reportable suicide rates in males, at the SA4 level, were:

- highest in the SA4 areas of Wide Bay, Queensland (37.6 deaths per 100,000 population), Northern Territory - Outback (35.6) and Western Australia - Outback (North) (35.4).
- lowest in the SA4 areas of Sydney - Inner South West (9.7 deaths per 100,000 population), Sydney - Baulkham Hills and Hawkesbury (9.8) and Sydney - Ryde (9.9).

For females, reportable suicide rates over the 5-year period 2018-2022, at the SA4 level, were:

- highest in the SA4 areas of Northern Territory - Outback (15.7 deaths per 100,000 population), Western Australia - Outback (North) (13.1), and Queensland - Outback (11.3)
- lowest in the SA4 areas of Sydney - South West (1.7 deaths per 100,000 population) and Sydney - Blacktown (3.4) and Sydney - Parramatta (3.7).

The AIHW is committed to continually improving the quality, ease-of-use, and timeliness of its products. In this product, we are using a new data visualisation tool to present results by geographical areas using maps. We welcome any feedback on this new presentation and hope that it will provide useful insights into the topic. As this tool is a relatively new addition to our website, we will be continuing to work to enhance its use and would welcome any feedback.



## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on [13 11 14](tel:131114), or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)

## Intentional self-harm hospitalisations by local areas

Hospitalisations data for patients with intentional self-harm injuries includes those with and without suicidal intent. For further information see the [Technical notes](https://www.aihw.gov.au/suicide-self-harm-monitoring/data/technical-notes) (<https://www.aihw.gov.au/suicide-self-harm-monitoring/data/technical-notes>).

The rates of hospitalisations for intentional self-harm in small geographic areas can provide insight into the incidence of intentional self-harm in local communities.

Statistical Areas Level 3 (SA3s) is a type of [geographical classification](#)

([https://www.abs.gov.au/websitedbs/D3310114.nsf/home/Australian+Statistical+Geography+Standard+\(ASGS\)](https://www.abs.gov.au/websitedbs/D3310114.nsf/home/Australian+Statistical+Geography+Standard+(ASGS))) defined by the Australian Bureau of Statistics (ABS) to provide a regional breakdown of Australia. There are 336 geographical areas which cover states and territories (excluding SA3s associated with overseas territories and other) with boundaries defined by the ABS. Each SA3 generally has a population of between 30,000 and 130,000 people. Allocation to an SA3 for hospitalisation data is based on the patient's usual place of residence, rather than where they received treatment.

Variations in hospitalisation rates between geographical areas may be due to a range of factors. Crude hospitalisation rates at SA3s should be interpreted with caution as areas with small populations are more sensitive to changes in the number of hospitalisations.

### How to use these maps

Use the zoom and search functions to explore the map. Click on an area in the map to view additional information. Change maps by selecting to 'open' or 'close' the eye icon. The colour shading indicates different rates of intentional self-harm hospitalisations, with darker shades indicating a higher rate.

For the best experience, use Chrome, Edge, Firefox or Safari browsers. For more information on browser compatibility, see [Supported browsers](#) (<https://doc.arcgis.com/en/web-appbuilder/create-apps/supported-browsers.htm>).

### Intentional self-harm hospitalisations by local areas



(Map will open in a new window)

(<https://maps.arcgis.aihw.gov.au/portal/apps/experiencebuilder/experience/?id=132ed362bfeb49f8b743e1168589e980>)

Note: Data behind these maps are available on the [Data downloads](https://www.aihw.gov.au/suicide-self-harm-monitoring/data/data-downloads) (<https://www.aihw.gov.au/suicide-self-harm-monitoring/data/data-downloads>) page: 2021-22 National Morbidity Database - Intentional self-harm hospitalisations.

### Variation across local areas

In 2021-22, rates of hospitalisations for intentional self-harm across SA3 geographies varied widely.

- Across Australia, rates ranged from 27 per 100,000 population for intentional self-harm hospitalisations in Mandurah (Western Australia) to 482 in Barkly (Northern Territory).
- For females, rates of hospitalisation ranged from 33 per 100,000 population in Mandurah (Western Australia) to 672 in Barkly (Northern Territory).
- For males, rates ranged from 14 hospitalisations per 100,000 population in Parramatta (New South Wales) to 536 in Caboolture Hinterland (Queensland).

Rates of intentional self-harm hospitalisations for different age groups also varied widely between SA3s.

- Rates of hospitalisations for intentional self-harm for those aged 24 and below ranged from 23 hospitalisations per 100,000 population in Bankstown (New South Wales) to 698 in Whitsunday (Queensland).
- For the 25-44 age group, rates ranged from 20 hospitalisations per 100,000 population in Parramatta (New South Wales) to 799 in Caboolture Hinterland (Queensland).
- For those aged 45 and over, rates ranged from 17 hospitalisations per 100,000 population in Ku-ring-gai (New South Wales) to 353 in Alice Springs (Northern Territory).



## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on [13 11 14](tel:131114), or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)

## International estimates of death by intentional self-harm

Global statistics on suicide provide a broad view of the issue across the world and provide a means of evaluation to allow governments, policy makers and researchers to learn from each other to improve suicide prevention planning and decision making. The intent in providing this information is to contribute to an informed, open debate about ways to prevent suicide in Australia—not to create comparisons ranking suicide rates around the world.

These data are estimates based on modelling assumptions from the most recent update to the Global Burden of Disease Study (GBD 2019) and are sourced from the Global Health Data Exchange (GHDx), a data catalogue created and supported by the Institute for Health Metrics and Evaluation (IHME). For further information see [Global Health Data Exchange \(http://ghdx.healthdata.org/\)](http://ghdx.healthdata.org/) and IHME [Global burden of disease \(http://www.healthdata.org/gbd/2019\)](http://www.healthdata.org/gbd/2019).


The interactive data visualisation below allows you to view the most recent data (rates of suicide and years of life lost) from Australia, Organisation for Economic Co-operation and Development (OECD) member countries, G20 nations (19 member nations plus the remaining 24 European Union nations individually represented) and World Health Organization regions. You can view data for any country or region using the 'multiple values' selector.

Deaths due to deliberate self-harm by region, 1990 to 2019.

This line graph shows the self-harm measures from 1990-2020, in OECD countries, G20 countries and WHO regions. Users can filter the graph in various ways, including viewing the age-standardised rate or Years of Life Lost (YLL) due to deliberate self-harm, viewing the latest year of data only and filtering by age groups and sex. Users can also compare Australia to OECD countries, G20 countries and WHO regions. Overall, Australia tracks slightly above the averages of OECD countries, G20 countries and WHO regions in 2020. The average age-standardised suicide rate has declined steadily in OECD countries, G20 countries and WHO regions.

## Deaths due to deliberate self-harm by region, 1990 to 2019

Rate (Age-standardised, per 100,000)

Show uncertainty  
HideMeasure  
DeathsYear  
1990 to 2019

■ Australia  
■ G20  
■ OECD Countries  
■ WHO regions

Chart type  
Time seriesAge  
Age-standardisedSex  
Both

Global Burden of Disease Collaborative Network.  
 Global Burden of Disease Study 2019 (GBD 2019) Results.  
 Seattle, United States: Institute for Health Metrics and Evaluation (IHME), 2020.  
 Supplementary Table: GBD S1  
 Available from <http://ghdx.healthdata.org/gbd-results-tool>

See notes ▶

Data are presented as deaths or years of life lost due to death by intentional self-harm. The terms self-harm and suicide are used interchangeably. It should be noted that this terminology is different to that used in other sections of the *Suicide & self-harm monitoring* website, where the term self-harm refers to non-fatal injury rather than death. The ICD-10 codes used here include: X60-X64.9, X66-X84.9, Y87.0 which are slightly different to those reported in other sections of *Suicide & self-harm monitoring*.

International rates of deaths due to self-harm should be interpreted with caution as the quality of mortality data can vary between countries and there is a lack of consistency in methods of death registration. Also, due to stigma associated with suicide—and the fact that it is illegal in some countries—some countries are likely to underestimate suicide rates and this may bring into question the reliability of suicide-related statistics (particularly in countries with low reported suicide rates).

Overall, there has been a reduction in suicide rates since 1990 driven mostly by declines in Europe and South East Asia. Across other regions, suicide rates have remained relatively stable.

### Suicide rates by country

Of OECD nations in 2019, age-standardised suicide rates ranged from 2.8 per 100,000 in Turkey to 23.9 per 100,000 in Lithuania. Australia's 2019 estimated suicide rate (10.4 per 100,000 population) was in the middle of OECD countries (18 of 36) and was similar to those reported in Canada, Czech Republic, New Zealand, and Sweden. The suicide rates in Austria and the United States were higher at 11.3 and 11.7 per 100,000 of the population respectively. Suicide rates have been rising in the United States prior to 2020 (see [Deaths of despair](#)).

Similarly, in comparison with G20 nations in 2019, Australia was 23 of 43 (19 members nations plus remaining 24 European Union nations individually represented).

### Suicide is more common in males than females in all countries

Suicide rates for males and females can be explored for any country or region on the interactive visualisations by selecting the drop down options for sex.

In 2019, in OECD countries, rates for males varied from 4.4 per 100,000 in Turkey to 42.2 in Lithuania, while female suicide rates ranged from 1.3 per 100,000 in Greece to 11.8 in the Republic of Korea. Again, Australia was in between with suicide rates of 16.2 per 100,000 for males and 4.8 per 100,000 for females.

### Suicide rates by age

Suicide is one of the leading causes of death in young people in Australia; however, this does not necessarily mean suicide is more likely to occur in young people than in older age groups—it is largely a reflection of the fact that older Australians also die from many other causes.





## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on [13 11 14](tel:131114), or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)

## Suicide & self-harm monitoring: Behaviours & risk factors

Risk factors are behaviours or aspects of lifestyle, environmental exposures or inherited characteristics that can interact to influence people's risk of suicidal behaviours. Therefore, looking at risk factors at a population level can help target assistance.

It is important to remember that the presence of one or more of these risk factors cannot predict or explain suicide or intentional self-harm as each person's experience is unique. Experiencing any of these risk factors does not necessarily mean a person has—or ever will—attempt suicide, but establishing whether a person has any of these risk factors can help determine whether they are at increased risk. Also, some people will have suicidal thoughts without having a history of any risk factors.

Risk factors and behaviours can be modifiable (change over time; for example, illicit drug use) or non-modifiable (permanent or constant; for example, a personal history of self-harm). They can also be background factors (such as a childhood history of abuse) or recent stressful life events. The presence of these factors and their influence is different from person to person over their lifetime and can vary by sex, culture and other characteristics.

Information on these risk factors in Australians has been obtained from a number of sources by making greater use of existing data sets or by integrating multiple data sets. This includes:

- the presence of psychosocial factors (for example, a past history of self-harm; relationship problems; legal issues; bereavement; unemployment; homelessness; and disability) in deaths by suicide obtained by manual review of reports and coronial findings held by the National Coronial Information System (NCIS) by the Australian Bureau of Statistics
- the effect of differences in educational attainment and labour force status in deaths by suicide obtained by integrating the ABS Causes of Death data set with that of the Census 2011
- risk factors associated with suicide and self-inflicted injuries included in the Australian Burden of Disease Study 2015 (to be updated with 2019 data as soon as possible as per the recent AIHW report [The health impact of suicide and self-inflicted injuries in Australia, 2019](#)).

## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on [13 11 14](tel:131114), or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)

## Behavioural risk factor burden for suicide and self-inflicted injuries

The National Suicide and Self-harm Monitoring Project provided funding for the AIHW to produce a report on [The health impact of suicide and self-inflicted injuries in Australia, 2019](#). The report estimates the combined impact of people dying prematurely from suicide and the direct health impacts on individuals living with injury due to self-harm. Note that the estimates do not take into consideration the potential mental health issues associated with self-harm or the effects suicide and self-harm can have on people's families, friends and communities. Through detailed data visualisations the report presents time series data for the Australian population by age, sex and key population groups. The contribution of various modifiable risk factors to disease burden is also estimated.

Data on risk factors associated with suicide and self-inflicted injuries from the Australian Burden of Disease Study 2022 are included below. The full report is available [here](https://www.aihw.gov.au/reports/burden-of-disease/australian-burden-of-disease-study-2022/contents/about). (<https://www.aihw.gov.au/reports/burden-of-disease/australian-burden-of-disease-study-2022/contents/about>)

According to the AIHW's Australian Burden of Disease Study 2022, suicide and self-inflicted injuries was the second leading cause of premature death from injury or disease, accounting for an estimated 6% of the total years of life lost in Australia (AIHW, 2022). Moreover, suicide and self-inflicted injuries is the leading cause of premature death in men aged 15-49 years. See [Burden of disease](#) for further information.

### What is burden of disease?

Burden of disease analysis measures the impact of living with illness and injury and dying prematurely. The method uses the summary measure 'disability-adjusted life years (or DALY) to measure the years of healthy life lost by combining premature death (years of life lost; YLL) with years lived with disability (YLD). For further information including a more comprehensive explanation of the methodology and data sources used, see [Australian Burden of Disease Study: methods and supplementary material 2022](#) (<https://www.aihw.gov.au/reports/burden-of-disease/australian-burden-of-disease-study-2022/contents/technical-notes>).


The burden of suicide and self-inflicted injuries due to behavioural risk factors, known as attributable burden, has also been estimated in the Australian Burden of Disease Study. These estimates reflect the amount of burden that could have been avoided if all people in Australia were not exposed to the risk factor.

In 2022, 'suicide and self-inflicted injuries' was the second leading cause of fatal burden among all people, with an estimated 159,200 total YLL. Approximately 121,200 YLL were lost to suicide and self-inflicted injuries among men and 38,000 YLL among women. In 2022, suicide and self-inflicted injuries were also the second leading cause of fatal burden among men and the ninth leading cause of fatal burden among women (down from eighth in 2018).

The interactive data visualisation shows the leading causes for years of life lost (YLL) based on leading causes of mortality in Australia. Sex (females, males, persons) and data year can be selected for viewing. Highlighted in purple indicates YLL due to suicide and self-inflicted injuries.

## Leading causes of years of life lost (YLL) by sex, Australia, 2018

Persons



Select sex:  
 Females  
 Males  
 Persons

Select data year:  
 2018

Notes: 2022 estimates are based on trend analyses from previous years. For more information please see notes.

Supplementary Table: S1.1

Source: AIHW Australian Burden of Disease Database, 2022

[See notes ►](#)


The visualisation directly below shows the average YLL per individual deceased person separately for each of the study years (2003, 2011 and 2018). The causes of death shown in this visualisation are the top 20 leading contributors to years of life lost initially identified according to total YLL. The causes of death included are not necessarily among the top 20 leading causes according to average YLL.

In 2018, an average of 42.2 years were lost to ‘suicide and self-inflicted injuries’ among males, and 41.6 years were lost to ‘suicide and self-inflicted injuries’ among females.

The interactive data visualisation shows average years of life lost (YLL). The causes of death shown in this visualisation are the top 20 leading contributors to years of life lost initially identified according to total YLL. Sex (females, males, persons) and data year can be selected for viewing. Highlighted in purple indicates YLL due to suicide and self-inflicted injuries.

## Average years of life lost (YLL) per death, for the top 20 causes of YLL by sex, Australia, 2018

Persons



Select sex:  
 Females  
 Males  
 Persons

Select data year:  
 2018

Note: Averaged from the top 20 years leading causes of life lost, by sex, in each year.

Supplementary Table: S1.1

Source: AIHW Australian Burden of Disease Database

[See notes ►](#)

'Child abuse and neglect' during childhood was:

- consistently the leading behavioural risk factor contributing to the years of healthy life lost due to suicide and self-inflicted injuries in both men and women since 2003 and has increased at each time point.
- associated with 32% of the years of healthy life lost due to 'suicide and self-inflicted injuries' in men (about 25,700 DALYs) and 43% of the years of healthy life lost due to 'suicide and self-inflicted injuries' in women (about 12,000 DALYs) in 2019 with the vast majority of these years of healthy life lost due to premature death.

Until 2018, among men, the second and third leading risk factors contributing to the years of healthy life lost due to suicide and self-inflicted injuries were 'alcohol use' and 'illicit drug use' across all years of the Australian Burden of Disease Study. Since 2018, 'illicit drug use' became the second leading risk factor contributing to the years of healthy life lost due to suicide and self-inflicted injuries among men followed by 'alcohol use'. In 2019 this trend continued:

- 'Illicit drug use' was responsible for 23% (about 18,600 DALYs) of the years of healthy life lost to 'suicide and self-inflicted injuries' among men.
- 'Alcohol use' was responsible for 22% of the years of healthy life lost due to 'suicide and self-inflicted' injuries in men (about 18,100 DALYs)


For women, the second greatest contributor to the years of healthy life lost due to 'suicide and self-inflicted injuries' was 'intimate partner violence' (estimated in women only) which was consistent over all study years. The third leading contributor of healthy life lost due to suicide and self-inflicted injuries among women has remained 'illicit drug use' since 2018. In 2019:

- 'Intimate partner violence' contributed 25% of the years of healthy life lost due to suicide and self-inflicted injuries in women (about 7,000 DALYs).
- 'Illicit drug use' contributed to 11% of the years of healthy life lost to suicide and self-inflicted injuries (about 3,100 DALYs) among women.

The interactive data visualisation shows the burden (based on frequency) of suicide and self-inflicted injuries attributable to selected risk factors and categorised by age (from 5 years old to over 85). Selection for sex (females and males), data year (2003, 2011, 2015, 2018 and 2019) and attributable DALY, YLD and YLL are available for viewing.

### Burden (YLL, YLD, DALY) of suicide and self-inflicted injuries attributable to selected risk factors, by age group and sex, Australia, 2019

#### Attributable DALY



Note: Intimate partner violence risk factor is estimated in females only as the evidence in the literature used to inform the linked diseases and relative risks was available only for females (Ayre et al. 2016).  
 Source: AIHW Australian Burden of Disease Database.  
 Supplementary Table: S1.2  
 Latest data: 2019

See notes ►

In 2019, 'child abuse and neglect' during childhood was the greatest contributor to the years of healthy life lost due to suicide and self-inflicted injuries in both men and women in all age groups. The exception to this are women aged 85 years and over where 'intimate partner violence' was the highest contributor. The majority of the 'child abuse and neglect' burden was experienced among people aged 15-44 years. In females, the number of DALYs was similar across these age groups (about 2,000-2,900 DALYs). The highest among men was between ages 25-34 years (7,000 DALYs).

Similarly, most of the years of healthy life lost due to suicide and self-inflicted injuries attributable to 'alcohol use' or 'illicit drug use' was experienced in ages 15-54 years. Both risk factors were highest among both men and women aged 15-34 years.

The years of healthy life lost due to suicide and self-inflicted injuries in women that were attributable to 'intimate partner violence' was highest among women aged 35-44 years.

## References

Australian Institute of Health and Welfare 2022. [Australian Burden of Disease Study 2022](https://www.aihw.gov.au/reports/burden-of-disease/australian-burden-of-disease-study-2022/contents/about) (https://www.aihw.gov.au/reports/burden-of-disease/australian-burden-of-disease-study-2022/contents/about). Cat. no. BOD 37. Canberra: AIHW.

---

© Australian Institute of Health and Welfare 2024



## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on [13 11 14](tel:131114), or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)

## Impact of suicide among Aboriginal and Torres Strait Islander Australians

Suicide rates among Aboriginal and Torres Strait Islander people are substantially higher than those of non-Indigenous Australians (see [Deaths by suicide amongst Indigenous Australians \(https://www.aihw.gov.au/suicide-self-harm-monitoring/data/populations-age-groups/suicide-indigenous-australians\)](https://www.aihw.gov.au/suicide-self-harm-monitoring/data/populations-age-groups/suicide-indigenous-australians)). Reducing deaths by suicide and suicidal behaviour among Indigenous Australians is a public health priority for all Australian governments (Cth of Australia, 2022). Therefore, providing data and information about suicide and self-harm among Indigenous Australians is a priority for the National Suicide and Self-harm Monitoring System.

For more information on suicide among Indigenous Australians and efforts to enhance and improve access to the evidence base, see the [Indigenous Mental Health and Suicide Prevention Clearinghouse \(https://www.indigenoumhspsc.gov.au/\)](https://www.indigenoumhspsc.gov.au/).

The following is an overview of the estimated impact of suicide among Indigenous Australians, from AIHW's Australian Burden of Disease study (see [Australian Burden of Disease Study: impact and causes of illness and death in Aboriginal and Torres Strait Islander people 2018](https://www.aihw.gov.au/reports/burden-of-disease/illness-death-indigenous-2018/summary)) (<https://www.aihw.gov.au/reports/burden-of-disease/illness-death-indigenous-2018/summary>). The visualisation directly below shows the estimated total number of years of life lost (YLL) among Indigenous Australians for the top 20 leading contributors to YLL. This information is displayed separately for each of the study years (2003, 2011 and 2018).

Suicide and self-inflicted injuries were the second highest cause of total YLL among Indigenous Australians, second to coronary heart disease across all three data years. In 2018, there was approximately 10,800 YLL for suicide and self-inflicted injuries, an increase of around 2,400 years since 2011 (8,400 YLL) and 4,100 years since 2003 (6,700 YLL).


Among Indigenous men, suicide and self-inflicted injuries were the second highest cause of YLL, after coronary heart disease. Approximately 8,000 years of life were lost to suicide and self-inflicted injuries in 2018, compared to around 5,900 YLL in 2011 and 5,400 YLL in 2003.

In 2018, approximately 2,800 years of life were lost due to suicide and self-inflicted injuries among Indigenous women. Though suicide and self-inflicted injuries were the second leading cause of YLL in both 2018 and 2011, they were the sixth leading cause of YLL among Indigenous women in 2003.

The interactive data visualisation shows the leading causes for years of life lost (YLL) among Aboriginal and Torres Strait Islander people, based on leading causes of mortality in Australia. Sex (females, males, persons) and data year (2003, 2011, 2015, 2018) can be selected for viewing. Highlighted in purple indicates YLL due to suicide and self-inflicted injuries.

## Leading causes of years of life lost (YLL) among Aboriginal and Torres Strait Islander people, by sex, 2018

Persons



Select sex:  
Persons

Select data year:  
2018

Supplementary Table: S2.1  
Source: AIHW Australian Burden of Disease Database  
Latest data: 2018

[See notes ►](#)

The visualisation directly below shows the average YLL per individual deceased among Indigenous Australians separately for each of the study years (2018, 2011, and 2003). The causes of death shown in this visualisation are the top 20 leading contributors to total YLL identified above. The causes of death included are not necessarily among the top 20 leading causes according to average YLL.


Among all Indigenous people who died by suicide and self-inflicted injuries, an average of around 55 years of life were lost in 2018, 2011, and 2003. The average YLL by suicide and self-inflicted injuries is higher for Indigenous people compared to non-Indigenous people. The average YLL by suicide and self-inflicted injuries for non-Indigenous people was 41 in 2018 and 2011, and 43 in 2003.

Among Indigenous men, the average YLL per death by suicide and self-inflicted injuries was 55 in 2018 and 2011 and 56 in 2003. Among Indigenous women, the average YLL per death by suicide and self-inflicted injuries was 57 in 2018 and 2011 and 58 in 2003.

The interactive data visualisation shows the leading causes for average years of life lost (YLL) based on leading causes of mortality in Australia. Sex (females, males, persons) and data year (2003, 2011, 2015, 2018) can be selected for viewing. Highlighted in purple indicates YLL due to suicide and self-inflicted injuries.

**Average years of life lost (YLL) per death for the top 20 causes of YLL among Aboriginal and Torres Strait Islander people, by sex, 2018**

Males



Select sex: Males  
 Select data year: 2018

Note: Averaged from the top 20 years leading causes of life lost by sex in each year  
 Supplementary Table: S2.1  
 Source: AIHW Australian Burden of Disease Database  
 Latest data: 2018

See notes ►

**References**

AIHW (Australian Institute of Health and Welfare) (2023) *Deaths by suicide amongst Indigenous Australians* (<https://www.aihw.gov.au/suicide-self-harm-monitoring/data/populations-age-groups/suicide-indigenous-australians>), accessed 31 Jan 2023.

The Commonwealth of Australia (Cth of Australia) (2022) *National Mental Health and Suicide Prevention Agreement* (<https://federalfinancialrelations.gov.au/agreements/mental-health-suicide-prevention-agreement>), The Federal Financial Relations website, accessed 3 March 2023.





## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on [13 11 14](tel:131114), or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)

## Psychosocial risk factors and deaths by suicide

Capturing information on risk factors relating to deaths by suicide can highlight areas of a person's life experience that may need additional attention to provide the most effective suicide prevention interventions. However, it is important to note that the presence of one or more of these risk factors in an individual's life does not necessarily mean they will experience suicidal behaviours. The vast majority of people who experience these risk factors will not experience suicidal behaviours.

As part of the National Suicide and Self-harm Monitoring Project the AIHW has funded the Australian Bureau of Statistics (ABS) to identify and code (using ICD-10) psychosocial risk factors for deaths referred to a coroner, including deaths by suicide.


From 2018 to 2022, around two-thirds of all deaths by suicide had at least one or more psychosocial risk factor identified (ABS, 2023). The types of psychosocial risk factors associated with deaths by suicide were age dependent and differed throughout the lifespan.

Most frequently occurring psychosocial risk factors in coroner-certified suicide deaths by age and sex, Australia, 2022.

The horizontal bar graph shows the proportion of coroner-certified deaths by suicide with psychosocial risk factors identified in males in Australia in 2022. The user can choose to view the data by sex, by age groups, and by the number of deaths by suicide with psychosocial risk factors identified.

## 2022 suicide deaths, by most frequent psychosocial risk factors for 2022 deaths by suicide

Per cent



Note: Data in this graph indicates the number and proportion of deaths with each specified risk factor recorded. Risk factors may not be mutually exclusive, and therefore people with multiple psychosocial factors recorded will be counted in more than one category.  
 Source: ABS Causes of Death, Australia 2023  
 Supplementary table: NMD S14  
 Latest data: 2022 (annual release)

[See notes ►](#)

In 2022, among those who died by suicide:

- ‘Personal history of self-harm (Z915)’ was the most commonly identified risk factor in males and females in all age groups (except males aged 45-54 years and males and females 65 years and over). The percentage of all suicide deaths with this risk factor was almost twice as high among females (34.0%) compared to males (17.4%). This code includes both intentional self-harm as well as suicide attempts (ABS, 2019).
- ‘Limitation of activities due to disability (Z736)’ was the most commonly identified risk factor in males and females aged 65 and over (27.1% and 35.0% respectively in 2022). This code included all types of disabilities as and health conditions which affected the individual’s abilities, such as chronic conditions in the elderly) and/or a perceived limitation of ability due to a newly diagnosed illness (ABS 2019).
- ‘Disruption of family by separation and divorce (Z635)’ was a common risk factor in males and females aged under 55. The percentage of all suicide deaths with this risk factor was higher among males compared to females across all age groups. This code includes relationship breakdowns or separation, divorce, and was also applied to children who died by suicide and were affected by separation of divorce of their guardians (ABS 2019).
- ‘Problems related to other legal circumstances (Z653)’ was a common risk factor in males aged 25-54 (associated with more than 10% of deaths by suicide). This code included domestic violence orders, child custody or support proceedings, litigation, restraining orders, potential or impeding legal circumstances or court appearances, charges which had been laid or the person was awaiting/anticipating commencement of legal proceedings and circumstances where death occurred in relation to illegal activities and not captured in a different code (ABS 2019).
- ‘Problems in relationship with spouse or partner (Z630)’ was also identified as a frequently occurring psychosocial risk factor in males and females across most age groups. This code includes intimate partner violence, relationship issues, acute events as well as ongoing/reoccurring, one off events which impacted the chain of events (leading to death) and domestic violence (ABS, 2019).
- ‘Other problems related to housing and economic circumstances’ codes Z599 and Z598 emerged as another common risk factor in males aged 55-64 and females aged 45-54. According to the ABS (2019) annex, code Z598 refers to ‘Other and unspecified problems related to economic circumstances’. This code included foreclosures on loans, problems with creditors, financial loss/difficulty/issues/troubles/concerns/problems/stress, bankruptcy, unemployment of a family member, unspecified financial issues or any other specific economic circumstances which do not fit in Z590-Z597 (ABS, 2019). Code Z599 refers to ‘Other and unspecified problems related to housing circumstances’. This included temporary housing, eviction from housing (owner or renter), concerns over security of keeping or obtaining housing, and other specified housing circumstances which did not fit within codes Z590-Z597 (ABS, 2019).

There is no national standard for the collection of data on psychosocial factors—each state and territory has its own legislation and processes relating to coroner-certified deaths meaning that the type of information collected and held by the NCIS database differs by jurisdiction. Also, due to the method used for the collection of data, protective factors are not included. For the full ABS annex and code

definitions, please click [here](https://www.abs.gov.au/statistics/research/psychosocial-risk-factors-they-relate-coroner-referred-deaths-australia#annex-listing-psychosocial-codes-inclusions-and-exclusions-) (<https://www.abs.gov.au/statistics/research/psychosocial-risk-factors-they-relate-coroner-referred-deaths-australia#annex-listing-psychosocial-codes-inclusions-and-exclusions->).

## COVID-19 psychosocial risk factors

Risk factors which were prevalent among cases of death by suicide in 2020-2021, namely those related to the COVID-19 pandemic, are rarely among the those most frequently occurring risk factors in 2022. Risk factors related to the pandemic are therefore low in prevalence when looking at the 2020 and 2021 years.

In 2020, the ABS added codes for the capture of the COVID-19 pandemic as a risk factor based on how it was described as part of the coronial investigation:

- F41.8 Pandemic-related anxiety and stress
- Z29.0 Isolation or quarantine (hotel or home), and
- Z29.9 Prophylactic measures put in place through health directives, including closure of business and stay at home measures.

In 2022, there were 84 people who died by suicide (2.6% of all suicides) who had the COVID-19 pandemic mentioned in either a police or pathology report, or a coronial finding. However, for people who died by suicide and had the COVID-19 pandemic mentioned as a risk factor, it did not appear as an isolated risk (they had, on average, 6.5 risk factors and 3.5 psychosocial risk factors). It is important to remember that circumstances relating to suicide are complex and multifaceted and a combination of multiple factors contribute to a person taking their own life rather than a single reason.

In 2022, of those who died by suicide with issues relating to the COVID-19 pandemic as a risk factor:

- 47.6% also had problems relating to employment or unemployment
- 46.4% also had mood disorders, including depression
- 17 people also had problems related to the social environment, including social isolation (this is down from 33 in 2020 but increase from 11 in 2021)
- the majority of people lived in New South Wales, Victoria, or Queensland (ABS 2023).

When COVID-19 was mentioned as a risk factor it manifested in different ways. For some people direct impacts from the pandemic, such as job loss, lack of financial security, family and relationship pressures, and not feeling comfortable with accessing health care were noted. For others, a general concern or anxiety about the pandemic and societal changes were stated or anxiety about contracting the virus itself. For further information about how the pandemic affected suicide rates, please visit the Suicide and Self-harm Monitoring COVID-19 page, [here](#).

The ABS reviewed and coded psychosocial risk factors (defined as social processes and social structures which can have an interaction with individual thought, behaviour and/or health outcomes) associated with deaths by suicide in 2017 through a review of police, toxicology and pathology reports and coronial findings held by the NCIS. The AIHW is working with the ABS to continue this work and embed psychosocial risk factors in future national mortality data sets.

## References

ABS (Australian Bureau of Statistics) 2019, *Psychosocial risk factors as they relate to coroner-referred deaths in Australia* (<https://www.abs.gov.au/statistics/research/psychosocial-risk-factors-they-relate-coroner-referred-deaths-australia#annex-listing-psychosocial-codes-inclusions-and-exclusions->), ABS Website, accessed 4 October 2023.

ABS 2023. *Causes of Death, Australia 2022*. ABS Catalogue number 3303.0.

## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on [13 11 14](tel:131114), or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)

## Social and economic factors and deaths by suicide

There is growing evidence that social factors, including education, employment status, income level and wealth, play an important role in determining the risk of suicide in high income countries (Blakely et al, 2003).

A combination of factors contribute to someone considering suicide. Although some social factors may be associated with an increased risk of suicide, they cannot be considered a direct cause.

Understanding how social factors affect the risk of suicide is important to better inform strategies to reduce suicide in Australia and may help in the planning of more effective evidence-based prevention and intervention programs.

Using linked data from the Multi-Agency Data Integration Project (MADIP), the AIHW has conducted two studies and a further study in collaboration with the Australian National University's Centre for Social Research and Methods to identify social and economic characteristics associated with greater risk of death by suicide. While these pieces of work are distinct, together they add to the growing understanding of population-level influences on suicide deaths in Australia.

The MADIP is a partnership among Australian Government agencies to link administrative and survey data. These studies used de-identified Australian Census of Population and Housing (2011) data linked with 7 years of Death Registrations (2011 to 2017). For more detailed information on the MADIP data asset, data linkage and analytical methods used, see [Technical notes](#).

Data linkage combines information from multiple sources, while preserving privacy. All linked data sets used for analysis at the AIHW comply with legislative and regulatory standards, are securely stored and accessed, and meet ethical standards and community expectations. Protocols are in place to prevent privacy breaches or the unauthorised identification of individuals, and to ensure data security and restricted access to information.

The initial analysis, [Educational attainment, employment and deaths by suicide](#), found that the cumulative risk of suicide in Australia is higher in those with fewer years of education and is lower among those who are employed. These results have been reported previously on [Suicide and self-harm monitoring](#).

Additional analysis, [Regression risk models for selected census variables](#), developed statistical regression models to examine the association between 10 identified predictive social and economic factors from the 2011 Census and deaths by suicide in Australia. The difference between this approach and the previous cumulative risk analysis, is that regression allows for adjustment for the various risk factors for suicide, which may make estimates more precise.

The multivariate (multiple variables) regression model showed that the strongest associations with deaths by suicide (relative to respective reference groups, and after adjusting for other variables in the model) included:

- being male (HR = 3.12; 95% CI 2.93 to 3.32)
- being widowed, divorced or separated (HR = 1.95; 95% CI 1.79 to 2.12)
- being in a lone person household (HR = 1.72; 95% CI 1.57 to 1.87)
- being unemployed (HR = 1.75; 95% CI 1.55 to 1.99) or not in the labour force (HR = 1.80; 95% CI 1.64 to 1.99)

Results for other variables are reported on [Regression risk models for selected census variables](#).

In further analysis, [Social and economic factors associated with suicide in Australia: a focus on individual income](#) reported here for the first time, a longitudinal approach was taken, which enabled the investigation of changes to individuals' income and employment status across time. It also examined the absolute risk, as well as relative odds of dying by suicide.

The longitudinal multivariate regression model confirmed findings from the Regression risk models for selected census variables study and produced additional insights into associations between deaths by suicide, income and income uncertainty including:

- those with higher income uncertainty had higher odds of suicide death relative to those with lower income uncertainty. Relative to those in the lowest income uncertainty quintile, the odds of dying by suicide increased by 1.91 (95% CI 0.29 to 0.44) for those in the highest income uncertainty quintile.
- people who experienced longer periods of unemployment had higher odds of suicide death. Relative to those with no periods of unemployment, the odds of dying by suicide increase by 1.57 (95% CI 1.21-2.05) for those unemployed for 2 years; 1.75 (95%CI 1.36-2.26) for those unemployed for 3 years; 2.03 (95% CI 1.61-2.57) for those unemployed for 4 years; and 1.96 (95% CI 1.61-2.57) for those unemployed for 5 years.

Additional results are reported on [Longitudinal analysis of income uncertainty](#) and the full report can be found on [Releases](#).

---



## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on [13 11 14](tel:131114), or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)

## Deaths by suicide, by socioeconomic areas

There is a strong association between socioeconomic status and deaths by suicide. Age-standardised rates and numbers of deaths by suicide tend to be higher for those living in lower socioeconomic areas (more disadvantaged areas). However, it is important to remember that suicide can affect all Australians and each person's experience is unique; not everyone who lives in these areas will experience suicidal behaviours.

Socioeconomic status classifies individuals according to the socioeconomic characteristics of the area in which they lived prior to their death by suicide. These areas are defined using the ABS Index of Relative Socio-Economic Disadvantage (IRSD), which estimate the level of socioeconomic disadvantage of the area, rather than individuals (see [Technical notes \(https://www.abs.gov.au/statistics/detailed-methodology-information/concepts-sources-methods/socio-economic-indexes-areas-seifa-technical-paper/2021\)](https://www.abs.gov.au/statistics/detailed-methodology-information/concepts-sources-methods/socio-economic-indexes-areas-seifa-technical-paper/2021) for more information). Variables used in calculating the IRSD index include household income, unemployment and levels of education.

Suicide deaths by socioeconomic area and mechanism, Australia, 2010 to 2022.

The series of line graphs show suicide deaths by socioeconomic areas (Quintiles 1 to 5) from 2001 to 2022. Users can choose to view age-standardised suicide rates or numbers of deaths by suicide. Users can choose to view suicide deaths by specified mechanisms (Firearms, gas, Hanging, Other mechanism, or Poisoning (except gas)). Users may also view the percentage of all suicide deaths that occurred by a specified mechanism.

---

### Content warning:

The data in this visualisation might be distressing to some readers as it contains data on the modality of suicide deaths and attempts. Please consider your need to read the following information. If this material raises concerns for you or if you need immediate assistance, please contact a [crisis support service](#), available free of charge, 24 hours a day, 7 days a week.

Please consider the [Mindframe guidelines](#) if reporting on these statistics.

[Proceed to visualisation](#)

## Highest rates of suicide occur in lowest socioeconomic areas

From 2001 to 2022, age-standardised suicide rates were highest for those who lived in the lowest socioeconomic areas (most disadvantaged areas), and generally decreased as the level of disadvantage lessened.

In 2022, the suicide rate for people living in the lowest socioeconomic (most disadvantaged) areas (18.4 deaths per 100,000 population; Quintile 1) was more than twice that of those living in the highest socioeconomic (least disadvantaged) areas (8.2 deaths per 100,000 population; Quintile 5). Similarly, the number of deaths by suicide generally declined as socioeconomic disadvantage decreased.

## Suicide rates increased over time in lowest socioeconomic areas

Overall, age-standardised suicide rates increased for those living in the lowest socioeconomic areas (Quintile 1); from 14.0 deaths per 100,000 population in 2001 to 18.4 deaths per 100,000 population in 2022. In contrast, smaller change was observed for those living in the higher socioeconomic areas (Quintiles 4 and 5).

Henley and Harrison (2019) found that over the period 2009-10 to 2015-16, suicide rates increased significantly for those living in the lowest socioeconomic areas (most disadvantaged) by an average 3.5% per year while little change was observed for those in the highest (least disadvantaged) socioeconomic areas (0.2% change per year).

## Methods of suicide vary by socioeconomic areas

Understanding the methods used for suicide can play an important role in suicide prevention. These data are provided to inform discussion around restriction of access to means as a policy intervention for the prevention of suicide.

Please consider your need to read the following information. If this material raises concerns for you or if you need immediate assistance, please contact a [crisis support service](#), available free of charge, 24 hours a day, 7 days a week.

Please consider the [Mindframe guidelines](https://mindframe.org.au/suicide/communicating-about-suicide/mindframe-guidelines) (<https://mindframe.org.au/suicide/communicating-about-suicide/mindframe-guidelines>) if reporting on these statistics.

The classification system used to code causes of deaths data, ICD-10, uses the term 'mechanism' to refer to the external cause of death. Throughout *Suicide & self-harm monitoring* website, 'mechanism' has been used in data visualisations, while the term 'method' has been used in the accompanying text.

Throughout 2001 to 2022, age-standardised suicide rates generally decreased with decreasing socioeconomic disadvantage for hanging (ICD-10 X70) and firearms (ICD-10 X72-X75). However, there was little difference in suicide rates between socioeconomic areas for poisoning excluding gas (ICD-10 X60-X66, X68-X69), poisoning by gas (ICD-10 X67), or other methods (ICD-10 X71, X76-X84, Y87.0).

In 2022 the rate of suicide by hanging for those living in the lowest socioeconomic areas (Quintile 1) was 2.5 times higher than that of those living in the highest socioeconomic areas (Quintile 5) (11.7 vs 4.6 deaths per 100,000 population).

Between 2001 and 2022, the proportion of all deaths by suicide that were completed by exposure to poisonous substances excluding gas or by other methods generally increased with decreasing socioeconomic disadvantage. Whereas the proportion of all deaths by suicide completed by hanging tended to decrease as socioeconomic disadvantage increased.

## Reference

AIHW: Henley G & Harrison JE 2019. Injury mortality and socioeconomic influence in Australia, 2015-16. Injury research and statistics series no. 128. Cat. no. INJCAT 208. Canberra: AIHW.

---

© Australian Institute of Health and Welfare 2024





## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on [13 11 14](tel:131114), or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)

## Intentional self-harm hospitalisations by socioeconomic areas

Hospitalisations data for patients with intentional self-harm injuries includes those with and without suicidal intent. For further information refer to the [Technical notes](#).

Socioeconomic status classifies individuals according to the socioeconomic characteristics of the area in which they live. These areas are defined using the ABS Index of Relative Socio-Economic Disadvantage (IRSD), which reflects the average level of socioeconomic disadvantage of the area (see [Technical notes](https://www.aihw.gov.au/suicide-self-harm-monitoring/data/technical-notes) (<https://www.aihw.gov.au/suicide-self-harm-monitoring/data/technical-notes>) for more information).

Intentional self-harm hospitalisations, by age, sex and socioeconomic areas, Australia, 2012-13 to 2020-22.

The line graph shows age-specific rates of intentional self-harm hospitalisations from 2012-13 to 2020-22 by socioeconomic areas from Quintile 1, the most disadvantaged, to Quintile 5, the least disadvantaged. Users can also choose to view age-specific rates, numbers and proportion of hospitalisations for intentional self-harm by socioeconomic areas by sex and specific age groups.

### Does socioeconomic status affect risk of intentional self-harm?

Rates of hospitalisations for intentional self-harm tend to be higher for those living in lower socioeconomic (more disadvantaged) areas.

In 2021-22:

- the rate for the most disadvantaged areas (Quintile 1) was 122 hospitalisations per 100,000 population, which is 1.5 times the rate for the least disadvantaged areas (Quintile 5; 82 per 100,000 population).

A similar pattern was seen in suicide rates in 2021, see [Suicide by socioeconomic areas](https://www.aihw.gov.au/suicide-self-harm-monitoring/data/behaviours-risk-factors/suicide-by-socioeconomic-areas) (<https://www.aihw.gov.au/suicide-self-harm-monitoring/data/behaviours-risk-factors/suicide-by-socioeconomic-areas>).

### How have rates of intentional self-harm hospitalisations changed for socioeconomic areas?

From 2012-13 to 2021-22:

- the highest proportion of intentional self-harm hospitalisations was for people living in the lowest socioeconomic (most disadvantaged) areas; this proportion has remained relatively stable over the period, averaging around 23%
- rates for males in the lowest socioeconomic areas, Quintile 1 and 2, increased from 115 and 98 hospitalisations per 100,000 population in 2012-13 to 129 and 110 in 2016-17, respectively, before decreasing to 87 and 70 hospitalisations per 100,000 population, respectively in 2021-22
- rates for females in the lowest (most disadvantaged) socioeconomic areas (Quintile 1) also increased from 179 in 2012-13 to 206 in 2016-17 and then decreased to 157 in 2021-22.

The highest age-specific rates of hospitalisations between 2012-13 and 2021-22 were recorded for those aged 25-44 for males and 0-24 for females, in the lowest socioeconomic areas (Quintile 1).

- Age-specific rates for intentional self-harm hospitalisations increased for all socioeconomic areas in females aged 0-24 from 2019-20 to 2020-21 before decreasing in 2021-22

- rates for females aged 25-44 in Quintile 1 increased from 243 per 100,000 population in 2012-13 to 272 in 2016-17 before falling to 174 in 2021-22
- rates for males aged 25-44 in Quintile 1 ranged from 197 in 2012-13 to 213 in 2016-17 then fell to 128 in 2021-22.

An increase in the rate of hospitalisations due to intentional self-harm for all socioeconomic areas was reported in 2016-17, which may be due to increases in hospitalisations in 3 states. Variation in hospital admission policy and practices between states and territories may have contributed to differences in the reporting of hospitalisation data. For further information, see the [data quality statement \(https://meteor.aihw.gov.au/content/724188\)](https://meteor.aihw.gov.au/content/724188).

---

## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on [13 11 14](tel:131114), or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)


## Deaths of despair

Since the late 1990s, there has been a marked increase in the overall mortality of middle-aged white non-Hispanic males and females in the United States (Case and Deaton 2015, 2017, 2020). This increase in mortality was largely attributed to increases in deaths by suicide, drug and alcohol poisonings (both accidental and undetermined intent) and deaths due to chronic liver diseases and cirrhosis - together termed 'deaths of despair' by Case and Deaton (2015, 2017, 2020). They linked this trend to a decline in economic security, a lack of universal health care and the widespread availability of opioids (Case and Deaton 2015, 2017, 2020). In 2017, Case and Deaton suggested that a similar increase in mortality from deaths of despair may be emerging in other countries (Case and Deaton 2017).

Selected causes of death, by sex, Australia, 1997 to 2022.

The line graph shows age-standardised rates of death by suicide, alcoholic liver disease and cirrhosis, accidental poisoning, and all of these causes combined from 1997 to 2022. Users can also choose to view age-standardised death rates and numbers of deaths for this period by sex and cause of death.

## Selected causes of death, by sex, Australia, 1997 to 2022



Note: Due to missing, not reported or overlapping data, numbers for demographic variables may not sum to the total.

Source: AIHW National Mortality Database and ABS Causes of Death, Australia 2023  
Supplementary table: NMD S13  
Latest data: 2022 (annual release)

[See notes ►](#)

An analysis of Australian mortality data using methods similar to those used by Case and Deaton shows that Australians are not increasingly dying due to these ‘deaths of despair’ over time. The rates of combined deaths by suicide, alcoholic liver disease and cirrhosis, and accidental poisoning (deaths of despair) over the period 1997 to 2022 show no clear trend. Since 2014 the rate has remained around 23 to 25 deaths per 100,000 population (from 2014 to 2022), similar to rates at the start of the period 1997 to 1999; between these dates rates remained lower (around 20.1 deaths per 100,000 population).

Males are more likely than females to die by these selected causes of death (suicide, alcoholic liver disease and cirrhosis, and accidental poisoning). At the start of the period, between 1997 and 1999, male rates of combined deaths by suicide, alcoholic liver disease and cirrhosis, and accidental poisoning ranged between 35.7 and 38.1 deaths per 100,000 population. Female rates, for the same period, ranged from 10.8 to 11.2 - 3.2 to 3.5 times lower than males. Since 2014, death rates for both males and females have shown little variation. Male rates ranged between 33.9 and 36.8 deaths per 100,000 population and female rates ranged from 12.3 to 13.8. Since 2014 these causes of death were, on average, 2.7 times more common in males than females.

### References

Case A & Deaton A 2020. Deaths of Despair and the Future of Capitalism. Princeton: Princeton University Press.

Case A & Deaton A 2017. Mortality and morbidity in the 21st century. Brookings Papers on Economic Activity 397.

Case A & Deaton A 2015. Rising morbidity and mortality in midlife among white non-Hispanic Americans in the 21st century. PNAS. 112(49):15078-15083.



## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on [13 11 14](tel:131114), or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)

## Technical notes

This section contains more detailed information about the data sources, codes and classifications, and analysis methods used in compiling data for *Suicide & self-harm monitoring*.

---

## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on [13 11 14](tel:131114), or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)

## Data sources

### National Mortality Database (NMD)

---

The AIHW National Mortality Database (NMD) contains records for deaths in Australia from 1964 to 2022. The database comprises information about causes of death and other characteristics of the person, such as sex, age at death, area of usual residence and Indigenous status.

The AIHW sources causes of deaths data from the Registries of Births, Deaths and Marriages in each state and territory and the National Coronial Information System (managed by the Victorian Department of Justice). The cause of death data are compiled and coded by the Australian Bureau of Statistics (ABS) to the International Statistical Classification of Diseases and Related Health Problems (ICD) and maintained at the AIHW in the NMD. Registration of deaths is the responsibility of the Registry of Births, Deaths and Marriages in each state and territory.

To improve the quality of data, the ABS annually revises the causes of death for coroner-referred deaths to reflect the latest available information. This process applies to deaths registered after 1 January 2006. Deaths registered between 2006 and 2019 are finalised. Deaths registered in 2020 are revised, deaths registered in 2021 are preliminary revised, and 2022 registered deaths are preliminary. Revised, preliminary revised, and preliminary data are subject to further revision by the ABS. For a more detailed description of the coverage and processing of deaths data, including deaths certified by the coroner, refer to the [ABS Causes of death, Australia methodology](#) (<https://www.abs.gov.au/methodologies/causes-death-australia-methodology/2022>), Australia (ABS Catalogue No. 3303.0).

In the NMD, the year the death occurred, the year the death was registered with the state and territory registry, and the in scope year the death was lodged with the ABS (ABS reference year) are provided. Year of registration has been used for the purposes of monitoring deaths by suicide. Deaths based on the year the death occurred have also been presented; however, as some deaths at the end of each calendar year may not be registered until the following year, year of death information for the latest available year (2022) is generally an underestimate of the actual number of deaths that occurred in that year. While not as significantly impacted, it should be noted that latest data by year of registration is also an underestimate and subject to revision.

In more recent years, there have been occasions where the ABS has received a large number of deaths, which were registered in Victoria in earlier years. For detailed information on this issue, see [Technical Note: Victorian additional registrations and time series adjustment](#) (<https://www.abs.gov.au/methodologies/causes-death-australia-methodology/2019#technical-note-victorian-additional-registrations-and-time-series-adjustment>) and [Technical Note: Victorian additional registrations \(2013-2016\)](#) (<https://www.abs.gov.au/methodologies/causes-death-australia-methodology/2021#technical-note-victorian-additional-registrations-2013-2016>).

Deaths (such as those from suicide) that are referred to a coroner can take time to be fully investigated, which can influence what information is available to assign a cause of death code during the ABS coding process. Each year, some coroner cases are coded by the ABS before the coronial proceedings are finalised. Coroner cases that have not been closed or had all information made available can impact on data quality as less specific ICD-10 codes often need to be applied. At the time of coding both 2021 and 2022 data there was a higher proportion of open coroner cases at preliminary coding than seen in previous years (67.2% in 2021 versus and 65.2% in 2022 versus a 5-year average for 2015-2019 of 56.2%). This is reflected in the 2021 and 2022 preliminary datasets by a higher rate of deaths due to 'other ill-defined and unspecified causes of mortality' (R99). In consideration of this, the ABS conducted an early revision of 2021 coroner certified deaths that had ill-defined causes. Therefore, data for 2021 are now considered preliminary revised. The ABS will apply the full revisions process for 2021 data and an early revision of 2022 data during the 2024 revisions cycle.

It is expected that deaths due to intentional self-harm will increase through the revisions process. For further information surrounding the revisions process, see Coding of suicide in the Deaths on the [ABS Causes of Death, Australia, 2022 methodology](https://www.abs.gov.au/methodologies/causes-death-australia-methodology/2022) (<https://www.abs.gov.au/methodologies/causes-death-australia-methodology/2022>) page.

The data quality statements underpinning the AIHW NMD can be found on the following ABS internet pages:

- ABS Quality declaration summary for [Causes of death, Australia methodology \(ABS cat. no. 3303.0\)](https://www.abs.gov.au/methodologies/causes-death-australia-methodology/2022#data-quality) (<https://www.abs.gov.au/methodologies/causes-death-australia-methodology/2022#data-quality>).
- ABS Quality declaration summary [Deaths, Australia](https://www.abs.gov.au/methodologies/deaths-australia-methodology/2021#data-quality) (<https://www.abs.gov.au/methodologies/deaths-australia-methodology/2021#data-quality>)

For more information on the AIHW NMD see [National Mortality Database](#) and [About National Mortality Database](#).

### Quality of Indigenous status data

The Indigenous status of a deceased person is captured through the death registration process; however, it is recognised that not all such deaths are captured through these processes, leading to under-identification. The Aboriginal and Torres Strait Islander (First Nations) origin of a deceased person is noted on the Death Registration Form (DRF) and the Medical Certificate of Cause of Death (MCCD).

For 2022 for New South Wales, information from the MCCD has been used for the first time. Using both sources (the DRF and MCCD) resulted in a greater proportion of deaths of First Nations origin, compared to 2021. This change has introduced a break in time series in First Nations death statistics in NSW and Australia. Therefore caution should be used when making comparisons with previous years. For more information on this change and the impacts refer to the Technical Note: [The impact of using two sources for deriving the Indigenous status of deaths in NSW in 2022](https://www.abs.gov.au/methodologies/causes-death-australia-methodology/2022#technical-note-the-impact-of-using-two-sources-for-deriving-the-indigenous-status-of-deaths-in-nsw-in-2022) (<https://www.abs.gov.au/methodologies/causes-death-australia-methodology/2022#technical-note-the-impact-of-using-two-sources-for-deriving-the-indigenous-status-of-deaths-in-nsw-in-2022>).

Data on deaths by suicide in Indigenous people have been compiled by jurisdiction of usual residence for New South Wales, Queensland, Western Australia, South Australia and the Northern Territory only. Data for Victoria, Tasmania and the Australian Capital Territory have been excluded in line with national reporting guidelines.

### National Hospital Morbidity Database (NHMD)

Data for patients who were hospitalised with intentional self-harm injuries are sourced from the AIHW's National Hospital Morbidity Database (NHMD). Most of the data used for the monitoring of hospitalisations for intentional self-harm are from 2008-09 to 2021-22. For each reference year, the NHMD includes all hospitalisations for patients who were discharged between 1 July and 30 June.

The NHMD is a compilation of episode-level records from admitted patient morbidity data collection systems in Australian hospitals. It is a comprehensive data set that has records for all episodes of admitted patient care from essentially all public and private hospitals in Australia.

The data supplied are based on the National Minimum Data Set (NMDS) for Admitted Patient Care and include administrative, demographic, clinical and length of stay data, as well as data on the diagnoses of the patients, the procedures they underwent in hospital and external causes of injury and poisoning.

The purpose of the NMDS for Admitted Patient Care is to collect information about care provided to admitted patients in Australian hospitals. The scope of the NMDS includes episodes of care for admitted patients in all public and private acute and psychiatric hospitals, free standing day hospital facilities, and alcohol and drug treatment centres in Australia. Hospitals operated by the Australian Defence Force, corrections authorities and in Australia's off-shore territories are not in scope but may be included. Hospitals specialising in dental, ophthalmic aids and other specialised acute medical or surgical care are included.

**episode of care:** The period of admitted patient care between a formal or statistical admission and a formal or statistical separation, characterised by only one care type (see care type and separation). METeOR identifier: 268956.

**separation:** The process by which an episode of care for an admitted patient ceases. A separation may be formal or statistical. METeOR identifier: 327268.

**formal separation:** The administrative process by which a hospital records the cessation of treatment and/or care and/or accommodation of a patient.

**statistical separation:** The administrative process by which a hospital records the cessation of an episode of care for a patient within the one hospital stay.

The criteria used to describe intentional self-harm hospitalisations reported in *Suicide & self-harm monitoring* is described in the [Codes and classifications](#) section.

### Data limitations

States and territories are primarily responsible for the quality of the data they provide. However, the AIHW undertakes extensive validations on receipt of data, checking for valid values, logical consistency and historical consistency. Where possible, data in individual data sets are checked with data from other data sets. Potential errors are queried with jurisdictions, and corrections and resubmissions may be made in response to these queries. Except as noted, the AIHW does not adjust data to account for possible data errors or missing or incorrect values.

The most recent [Data quality statement for Admitted Patient Care \(https://meteor.aihw.gov.au/content/index.phtml/itemId/724188\)](https://meteor.aihw.gov.au/content/index.phtml/itemId/724188) is available in METeOR. The Data Quality Statement contains information on other changes that may affect interpretation of the data for the relevant year.

### Quality of Indigenous status data

The Indigenous status data in the NHMD for all states and territories are considered to be of sufficient quality for statistical reporting. In 2011-12, an estimated 88% of Indigenous patients were correctly identified in public hospitals (AIHW 2013). The overall quality of the data provided for Indigenous status needs some improvement and varied between states and territories. It is unknown to what extent Indigenous Australians might be under-identified in private hospital admissions data.

### National Ambulance Surveillance System (NASS)

---

The National Ambulance Surveillance System (NASS) is a public health monitoring system, which aims to provide timely and comprehensive data on intentional self-harm (including suicidal behaviours with self-injurious intent), mental health, and alcohol and drug harms in the community. Data for the National Ambulance Surveillance System (NASS) are compiled by Turning Point in partnership with Monash University and are sourced from paramedic electronic patient care records provided by Australian state and territory-based ambulance services in New South Wales, Victoria, Queensland, Tasmania, and the Australian Capital Territory. As part of the National Suicide and Self-harm Monitoring Project, the AIHW has contracted Turning Point through Monash University to develop and maintain the National Ambulance Surveillance System (NASS) for self-harm related ambulance attendances. Self-harm (suicidal ideation, suicide attempt, self-injury) related modules from the NASS are reported here.

Information is obtained and coded through manual scrutiny of de-identified electronic patient care records (ePCRs), including paramedic clinical assessment, patient self-report, information from third parties and other evidence at the scene, such as written statements of intent (including social media, text messages and written notes), as recorded by paramedics. Intent of self-harm behaviours derived from the ePCR may be from either stated or physical evidence, or where there is evidence but the patient may have denied the behavioural intent (Lubman et al. 2020).

Self-harm related ambulance attendances are included if self-harm occurred in the preceding (past 24 hours) or during the ambulance attendance, with 4 categories of self-harm related ambulance attendances defined and coded as:

- self-injury (non-fatal intentional injury without suicidal intent)
- suicidal ideation (thinking about killing oneself without acting on the thoughts)
- suicide attempt (non-fatal intentional injury with suicidal intent, regardless of likelihood of lethality)
- suicide (fatal intentional injury with suicidal intent).

Suicide, suicide attempt and suicidal ideation are considered mutually exclusive; however, self-injury could be simultaneously coded with any other self-harm case category.

The number of attendances related to suicide is under-represented as ambulances do not attend all deaths. Furthermore, when they do attend there may be insufficient information to determine suicidal intent at the scene.

A modified ICD-10 coding framework is used to code the methods of suicide, suicide attempt or suicidal ideation, methods of self-injury and categories of suicidal ideation preparation (planned, unplanned and unknown if planned).

For more information see [Lubman et al. 2020 \(https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0236344\)](https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0236344).

### Data limitations

Data are collected for operational rather than monitoring or research purposes with paramedics only recording information that they either observe or is provided to them by the patient or bystanders, and which they deem clinically relevant to patient care. It is possible that relevant information with respect to self-harm or mental health variables is not recorded, or similar events may not be recorded consistently by different paramedics over time.

### Multi-Agency Data Integration Project (MADIP)

---

The Multi-Agency Data Integration Project (MADIP) is a partnership among Australian Government agencies to develop a secure and enduring approach for combining information on healthcare, education, government payments, personal income tax, and demographics (including the Census) to create a comprehensive picture of Australian populations over time (ABS 2018). The key MADIP datasets used in analysis published on the Suicide and Self-Harm Monitoring site were:

- Person Linkage Spine (Australian Bureau of Statistics)
- 2011 Census of Housing and Population (Australian Bureau of Statistics)
- Causes of Death (Australian Bureau of Statistics)
- Personal Income Tax (Australian Taxation Office)
- Social Security and Related Information (Department of Social Services)
- Synthetic income data developed by the Australian National University using personal income tax data, social security payment information and Census (for more information see [Biddle & Marasinghe 2021 \(https://taxpolicy.crawford.anu.edu.au/publication/ttppi-working-papers/18706/using-census-social-security-and-tax-data-multi-agency-data\)](https://taxpolicy.crawford.anu.edu.au/publication/ttppi-working-papers/18706/using-census-social-security-and-tax-data-multi-agency-data)).



## Linkage approach

In order to identify socioeconomic factors associated with deaths by suicide in Australia, 2011 Census and 2011 to 2017 Causes of Death data were linked to the ABS Person Linkage Spine (Spine). The Spine is comprised of all persons in the Medicare Enrolments Database, Personal Income Tax or Social Security and Related Information data sets at any point between 2006 and 2016 (ABS 2019). As the baseline population, 2011 Census was considered a closed population and several assumptions were made about this population. These include:

- everyone in the 2011 Census who did not die over the period were still in the population up to the end of 2017, that is, no migration occurred
- person information in the 2011 Census were held constant over the analysis period. However, in the modelling analysis conducted, time varying age and income of the year before suicide were calculated and applied.

Table 1 shows the linkage coverage of Census 2011 and deaths by suicide from the ABS Causes of Death. The Estimated Residential Population of Australia at 30 September 2011 was 22.43 million people (ABS 2021). Of these, 20,739,159 were accounted for in the Census 2011, noting that the Census 2011 started in August 2011. In total, the linked Census 2011 population was 16,700,062 (74.4% of the total Australian population of September 2011). According to the National Deaths Index, there are 17,306 deaths by suicide from September 2011 to December 2017, of which 11,580 (67%) deaths by suicide were linked to the linkable Census 2011 data. Suicide was defined by ICD-10 external cause codes X60-X84 and Y87.0

Table 1: Linkage coverage of 2011 Census population and deaths by suicide in ABS MADIP

|                                  | Total (n)  | Linked (n) | Linked (%) |
|----------------------------------|------------|------------|------------|
| ERP <sup>(a)</sup> at Sept 2011  | 22,432,771 | 16,700,062 | 74         |
| Deaths by suicide <sup>(b)</sup> | 17,306     | 11,580     | 67         |

a. Estimated resident population. Linked records are from 2011 Census population.

b. Linked deaths by suicide weighted to all deaths by suicide from September 2011 to December 2017.

## Estimated suicide risk by educational attainment and employment method

### Imputing weights for unlinked suicide deaths and 2011 Census

To address the issue of unlinked deaths by suicide and 2011 Census records, an imputation weighting technique was used. This section describes the method used to develop these weights, which involved a three-staged approach.

First stage: imputing weights to scale up the Census population. The ABS historical ERP for 31 December 2011 by states, sex and 5-year age groups were used to derive weights by these demographic characteristics, based on the assumption that there were no significant differences in the age distribution of the population. The derived weight was applied at the person level for each record of Census that has ABS Person Linkage Spine (Spine) information to enable analysts to weight the analyses to the 31 December 2011 total ERP.

Unlike the original ABS research paper (ABS 2016) describing the creation of a linked data set between 2011 Census and deaths registered in the following 13 months, the imputation method did not calculate weights by Indigenous and non-Indigenous populations. Also, note that Diplomatic personnel resident in Australia have not been excluded from total ERP.

Second stage: suicide weights were calculated by using all deaths by suicide from 2011 to 2017 by states and territories, sex and 5-year age groups. Suicide weights were then applied at person level to only those linked Census records with suicide information. This made it possible to weight the analyses to all deaths by suicide (18,848) from 2011-2017.

An issue with applying suicide weights is that suicide weights are slightly higher when compared with population weights applied in the first stage. As such, the combined weights of the linked records with both 2011 Census and suicide information when aggregated, the weighted ERP will be slightly higher than that of 31 December 2011. Hence the need for a scale down adjustment factor.

Third stage: Finally, a scale down adjustment factor, derived based on total ERP, linked deaths by suicide and all deaths by suicide, was applied at the person level to only Census records without linked death by suicide information. Hence the weights of the Census population with or without linked death by suicide information, aggregated to the 31 December 2011 ERP (22,340,025).

### Cumulative suicide incidence

Australian residents in the 2011 Census, weighted to 31 December 2011 estimated resident population (ERP) and linked to ABS Causes of Death data from 2011 to 2017 created a binary outcome of either died by suicide (ICD 10 external cause codes X60-X84, Y87.0) or not. Note that deaths by suicide used in this analysis are based on year of occurrence. These may differ from deaths by suicide data used in other AIHW publications which are based on year of registration. In addition to the closed population assumptions noted above, due to data quality issues the age in this analysis is at the time of the 2011 Census except for those who have died by suicide.

Over the period 2011 to 2017, Australia recorded more than 18,800 deaths by suicide of people who were in the 2011 Census. This resulted in a cumulative incidence of about 84 per 100,000 people during the 7-year period. The cumulative number and incidence of deaths by suicide that occurred over the 7 years varies considerably by sex, educational attainment and labour force status.

### Uncertainty in the estimates

All data are subject to some level of uncertainty. For the data presented in this analysis the sources of uncertainty include:

Linkage error: Uncertainty is introduced when there is error in linking data sets. The data used in this report carries some risk of linkage error. An attempt has been made to reduce this error through imputation weighting process but some uncertainty remains.

Timeliness of data: Some of the data used in this analysis is Census data collected in August 2011. A person's education status and employment status can change over time, particularly for certain population groups. The use of out-of-date information introduces a source of error to the analysis.

Randomness in the number of deaths by suicide that occur in a given time period, 2011-2017: The number of deaths by suicide that occur in a given time period fluctuate, even if the underlying population risk remains the same. The exact distribution of the counts is unknown. With deaths by suicide being a rare event it is often assumed that the counts follow a Poisson distribution. If this is the case then the relative level of uncertainty due to randomness decreases as the number of deaths by suicide increase.

## Regression risk models for selected census variables

The MADIP datasets used in this modelling are outlined in the Data section of these Technical notes. In this analysis, only people aged 25 to 64 years in the linked 2011 Census have been included, representing, over 9 million people in the 2011 Census and 7,000 deaths by suicide from 2011 to 2017. This age group was chosen because most deaths by suicide occur between these ages and because of the relative stability of socioeconomic factors over time (such as level of education) among this age group. While suicide is the leading cause of death among people aged 15 to 24 years, people in this age group were excluded from the modelling because of their lack of socioeconomic stability.

Missing values have been excluded from this analysis. Educational attainment has the highest proportion of missing values (5.5%). Unlike with the cumulative suicide risk estimations, the data used in the regression modelling has not been weighted.

To identify modelling predictors and explore their association with suicide deaths, an extensive literature review of social factors was carried out. This included earlier analyses published by AIHW, which showed deaths by suicide varied by factors such as employment and educational attainment.

Socioeconomic factors identified from the 2011 Census were used as predictors and deaths by suicide as the outcome variable. A total of 10 factors were included:

- Age (10-year age groups)
- Sex
- Indigenous status
- Registered marital status
- Family household composition
- Highest level of educational attainment
- Labour force participation
- Occupation
- Synthetic total income (quartiles, see Biddle & Marasinghe 2021)
- Need for assistance with core activities of daily living.

## Method

Two modelling approaches were tested: Poisson regression and competing-risks regression (as described by Fine & Gray 1999). For Poisson regression, counts of the outcome variable with the value 1 for deaths by suicide and 0 for those who did not die by suicide were created and data aggregated by socioeconomic factor.

For the competing-risks regression, the influence of other causes of death is considered. This is because people who died from any other causes (such as cancer and coronary heart disease) are no longer at risk of dying by suicide.

Sex-stratified and Indigenous-stratified multivariate models were also fitted to investigate the associations within males and females, and within Indigenous and non-Indigenous people. Due to data quality issues including small sample sizes, Indigenous-stratified models have not been published. Univariate and multivariate models (including quasi-Poisson to deal with slight overdispersion) were also refitted. The coefficients obtained were back transformed so they could be interpreted as rate ratios (for Poisson models) and subhazard ratios (for competing-risks models). Analysis was conducted using R (glm package) and Stata (version 16) software.

Of the models tested, competing-risks regression, a method that accounts for people being censored from the risk set because of a competing cause, was used to estimate the risk of death by suicide and the selected socioeconomic factors. Univariate, multivariate and sex-stratified competing-risks models were developed. Generally, competing-risks regression models can be regarded as an extension of the Cox proportional hazards model, where subjects who experience competing events (deaths from other causes) are adequately counted as not having any chance of dying by suicide.

Estimated coefficients of competing-risks models can be interpreted in a similar way as coefficients estimated from a Cox model, except that they estimate the effect of certain covariates in the presence of competing events. Note that the transformed coefficients are known as subhazard ratios, similar to hazard ratios estimated in Cox regression. The subhazard ratio can be interpreted as a rate ratio (Henan 2010), but here we are considering the relative change in rates of the event in those subjects who are either currently event-free or who have previously experienced a competing event (Austin & Fine 2017). For simplicity and ease of understanding, coefficients in this report are referred to as hazard ratios.

## Social and economic factors associated with suicide in Australia: a focus on individual income

Researchers from the Australian National University’s Centre for Social Research and Methods (CSRМ), in close collaboration with the AIHW, have extended the analysis [Regression risk models for selected census variables](#). An extract from the Multiagency Data Integration Project (MADIP) was used. For this analysis, the following MADIP datasets were utilized. 2011 Census information, 2011-2016 Personal income tax (PIT) records, 2011-2016 Social security and related information (SSRI) and 2011-2016 cause of death data. The data linkage process was identical to the linkage process described in the Data section of these Technical notes.

The following set of explanatory variables were used to capture the social and economic factors. More specifically, the CSRМ used the time-invariant 2011 Census data to capture social factors and time-variant PIT and SSRI to capture economic factors of suicide.

Table 2 - Explanatory variables and sources

| Variable                                    | Source           | Type     |
|---|------------------|----------|
| 1. Highest level of education               | 2011 Census      | Social   |
| 2. Age                                      | 2011 Census      | Social   |
| 3. Sex                                      | 2011 Census      | Social   |
| 4. Indigenous status                        | 2011 Census      | Social   |
| 5. Need for assistance with core activities | 2011 Census      | Social   |
| 6. Household composition                    | 2011 Census      | Social   |
| 7. Total income                             | 2011 - 2016 PIT  | Economic |
| 8. SSRI                                     | 2011 - 2016 SSRI | Economic |

In addition to the variables presented in Table 1, the following variables were derived using Census, PIT and SSRI data.

1. A synthetic measure of income - A measure of income that was derived using a machine learning algorithm. This income variable provides a representative measure of income of the entire Australian population (Biddle and Marasinghe 2021). This measure was then divided into quintiles to obtain a relative measure of income.
2. Coefficient variation of income (relative standard deviation) - Coefficient of variation of income was used to capture the income uncertainty of each individual. The measure was defined such that it captured income uncertainty between the current year and the previous year (i.e. variation between t and t-1). This measure was then divided into quintiles. An income uncertainty of quintile 1 implied an individual had low-income variation relative those in higher income uncertainty quintiles. Furthermore, since this measure uses data from t-1, the scope of the study was limited to 2012 January to 2016 December.
3. Proxy for unemployment - Unemployment status of an individual was captured using SSRI information. More specifically, if an individual received Newstart allowance and Youth allowance, the individual was then classified as being unemployed that period.

## Methods

### Longitudinal (panel) regression

The MADIP dataset is a longitudinal dataset therefore, utilizing longitudinal regression methods allows us to control for unobserved individual heterogeneity across the time period. For example, longitudinal regression methods would allow us to control for unobserved measures such as behavioural differences and cultural differences across individuals. Given that suicide is a complex individual decision and may not be entirely explained by observed variables, controlling for these unobserved heterogeneity would allow us to obtain unbiased estimates.

In this analysis, two competing longitudinal regression methods were tested - a random effects logistic model and a population-averaged logistic model. Both models were estimated with robust standard errors to account for heteroskedasticity.

Given that the random effects logistic model is dependent on the strong assumption that the underlying variation have no serial correlation, the population-averaged logistic model was selected as the primary regression method due to its robustness to serial correlation (Hill et al., 2010). Furthermore, a likelihood-ratio test was also undertaken to decide between the two models. The result suggested that the population-averaged model was more appropriate than the Random effects model. Equation (1) and (2) outline the population-averaged model.


$$y_{i,t}^* = \alpha + \beta x_i + \delta z_{i,t} + \varepsilon_{i,t} \quad (1)$$

$$y_{i,t} = \begin{cases} 1, & y_{i,t}^* > 0 \\ 0, & \text{otherwise} \end{cases} \quad (2)$$

Where  $y_{i,t}$  is the dependent variable which takes the value 1 if individual i has completed suicide at time t and 0 otherwise.  $x_i$  is a vector of time-invariant explanatory variables,  $z_{i,t}$  is a vector of time-varying explanatory variables and  $\varepsilon$  is the error term which is assumed to be independent and identically distributed with  $\varepsilon \sim (0, \sigma^2)$ .

Setting up the dataset as a panel allowed us to account for individual heterogeneity. However, given that suicide is a rare event, explanatory variables with large number of categories (for example - occupation) were excluded from the longitudinal analysis. This was primarily due the low number of suicides each year, which in turn made the models more sensitive to variables with a large number of categories. Given this drawback of panel data, a cross-sectional analysis was also conducted as a part of the sensitivity analysis.

Figure 1 - Annual suicide counts<sup>1</sup>



<sup>1</sup> Year 2011 was not included in the analysis.

## Reference

Hill, R.C., Griffiths, W.E. and Lim, G.C., 2010. Principles of econometrics. pp 537-560. John Wiley & Sons.

## Australian Defence Force (ADF) Suicide Data Sources

In addition to the NMD, the Australian Defence Force (ADF) suicide monitoring analysis used the following data sources:

### National Death Index (NDI)

The NDI is managed by the AIHW and contains person-level records of all deaths in Australia since 1980 obtained from the Registrars of Births, Deaths and Marriage in each state and territory. Its use is confined to data linkage studies approved by the AIHW Ethics Committee for health and medical research. NDI records are supplemented with cause of death information from the NMD. In this study, the NDI is linked with Defence payroll data to create the linked Defence payroll-NDI data set used in analysis of suicide in the ADF population.

### Department of Defence personnel system data

The Department of Defence compiled a file of current and historical Defence personnel systems covering ADF members who have served since 1 January 1985. This combines PMKeyS, Core HR system, D1, CENRESPAY (for reservists), ADFPAY (for permanent members) and other historical payment systems. The Department of Defence and AIHW assessed the resulting file for completeness and duplicates. Comparisons were made with records from Department of Defence annual reports and other sources to validate the list. Data from the National Archives was also investigated for its suitability in validation, however as the majority of records are electronic files based on photos of paper records, this was not usable.

For further information see [Technical notes of Serving and ex-serving Australian Defence Force members who have served since 1985: suicide monitoring 1997 to 2021](#).

## Australian Burden of Disease Study (ABDS)

Estimates of fatal (years of life lost, YLL) and non-fatal burden (years lived with disability, YLD) were sourced from the Australian Burden of

Disease Study (ABDS) 2015. The ABDS 2015 used burden of disease analysis to measure the impact of 216 diseases and injuries on the health of the Australian population. The study provides a detailed picture of the burden of disease and injury in the Australian population in 2003, 2011 and 2015. It also includes estimates of the contribution made by selected risk factors on the disease and injury burden in Australia, and by socioeconomic areas for some risk factors.

The ABDS 2015 uses and adapts the methods of global studies to produce estimates that are more relevant to the Australian health policy context. The chosen reference period (2015) reflects the data availability from key data sources (such as the National Health Survey, deaths data, hospital admissions data and various disease registers) at the time of analysis.

Results from the study provide an important resource for health policy formulation, health service planning and population health monitoring. The results provide a foundation for further assessments.

Full details on the various methods, data sources and standard inputs used in the ABDS 2015 are available in [Australian Burden of Disease Study 2015: methods and supplementary material](#).

## Data from suicide registers

---

### New South Wales Suicide Monitoring System

The New South Wales (NSW) Suicide Monitoring System contains data on all suspected and confirmed suicide deaths from 2019 to present. Established in October 2020, the system is a collaboration between the NSW Ministry of Health, Department of Communities and Justice, the State Coroner and NSW Police.

The NSW Suicide Monitoring System contains initial police information of suspected suicide deaths from the JusticeLink information system, which is managed by NSW Department of Communities and Justice. The information is manually checked against other data sources, including coronial determination.

NSW Health publishes monthly reports on suspected deaths by suicide in NSW on their website [NSW Suicide Monitoring System](#) (<https://www.health.nsw.gov.au/towardszerosuicides/Pages/suicide-monitoring-system.aspx>).

### Victorian Suicide Register

The Victorian Suicide Register (VSR) contains data on all suicide deaths reported to the Coroners Court of Victoria (CCOV) from 2000 to present. The Coroners Prevention Unit, a specialist investigative service for Victorian Coroners, has managed the VSR since its implementation in 2012. The VSR operates with funding from the Victorian Department of Health.

VSR data are used to inform investigations into suspected suicides and support coronial recommendations to prevent similar deaths. VSR data are regularly shared with the Victorian Department of Health, AIHW, Victoria Police and other organisations involved in suicide prevention.

Data on the VSR are obtained and coded from materials gathered throughout the course of coronial investigation, including police notification of death, forensic reports (autopsy and toxicology), witness statements and medical records.

CCOV releases a range of suicide data reports from the VSR on their [website](#) ([https://www.coronerscourt.vic.gov.au/forms-resources/publications?combine=&field\\_audience\\_target\\_id=All&field\\_publication\\_type\\_target\\_id=All&year=&page=5](https://www.coronerscourt.vic.gov.au/forms-resources/publications?combine=&field_audience_target_id=All&field_publication_type_target_id=All&year=&page=5)), including information on suspected deaths by suicide on a monthly basis, and overviews of First Nations suicides in Victoria. CCOV has also published a report on suicide among LGBTQ+ people in Victoria.

### Queensland Suicide Register and interim Queensland Suicide Register

In Queensland, there are two systems that are used to monitor suicide deaths: the Queensland Suicide Register (QSR), which includes suicide data since 1990 and is used to monitor longer-term trends, and the interim Queensland Suicide Register (iQSR). The iQSR was established in 2011 to provide real-time information on suicide deaths.

Data on this website are from the iQSR. The iQSR contains interim data on suspected suicides in Queensland, recorded shortly after the death occurs. The data are based on initial police reports and other information that is available to police at the time when they refer the death to the coroner.

The QSR contains information on suicide deaths for which coronial investigations have been finalised. The QSR is based on more information than the iQSR, including toxicology reports, post-mortem examination and the finding from the coroner, including details on the context and circumstances of the death.

The QSR and iQSR are currently managed by the Queensland Mental Health Commission (QMHC) on behalf of the Queensland Government, with support from the Coroners Court of Queensland (CCQ) and Queensland Police Service (QPS). Prior to September 2023, the iQSR was managed by the Australian Institute for Suicide Research and Prevention (AISRAP) at Griffith University. The QMHC publishes [monthly reports](#) (<https://info.qmhc.qld.gov.au/suicide-data#:~:text=on%20suicide%20deaths.,The%20Queensland%20Suicide%20Register,and%20entered%20into%20the%20QSR.>) based on data from the iQSR, around eight weeks from the last day of the reporting month.

## National Integrated Health Services Information (NIHSI)

---

### End of life service use for those who died from suicide in Australia

## Data sources

Data are from the National Integrated Health Services Information Analysis Asset (NIHSI AA) version 0.5. This data asset includes mortality data together with information from hospital admissions, Medicare Benefits Schedule (MBS), Pharmaceutical Benefits Scheme (PBS) and residential aged care data.

The analysis population was those who had died between 1 July 2010 and 31 December 2017 in the linked National Deaths Index (NDI). Suicide was defined as the principal external cause of death in X60-X84 and Y87.0 based on the International Classification of Diseases, Tenth Revision (ICD-10) codes. Patient demographic information was taken from the NDI and is therefore accurate at the time of death not time of service. Only people whose age at death was between 15 to 64 years were included in the analysis. This was due to people in this age range making up the majority of those who die from suicide and to allow for better comparisons with deaths from other causes, which mostly occur in people older than 65 (AIHW 2022a). People without a primary cause of death and with sex not stated were also removed from the analysis due to small cell sizes.

## Method

The analysis included MBS, PBS emergency department presentation and outpatient services in addition to hospital admissions datasets. For more information on MBS item classification and PBS item classification, drawn from the Anatomical Therapeutic Chemical (ATC) codes (AIHW 2022b), visit [Mental health services in Australia - Data source and key concepts](#).

Hospitalisation data was taken from two sources: admitted patients and emergency department (ED) presentations. The method for counting hospital admissions in this analysis based on the method in a similar study by [Clapperton et al. \(2021\)](#). (<https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0252682>)

Within the NIHSI AA v0.5, hospital data pertains to only New South Wales (NSW), Victoria (Vic) (excluding Albury-Wodonga), South Australia (SA) and Tasmania (Tas) public hospitals. Admitted patient information also contains information from private hospitals in Victoria. To ensure accurate comparisons with hospitals data, only deaths registered in NSW, Vic, SA and Tas are included in the analysis.

Admitted patient data refers to only acute admitted and mental health separations (Admitted Patient Care National Minimum Data Set care types of 1, 7.1, 7.2, and 11). In scope separations where the patient was transferred from another hospital or had a change of care type in the same hospital are not counted in the total to avoid duplication.

Any hospital episode (ED presentation, hospital admission) that ended in “death” was excluded as it was considered to be a result of the fatal (suicide) incident. The only exception to this were episodes where the intentional self-harm was coded as occurring in a health service area—these episodes were retained as they were most likely inpatient suicides ([Clapperton et al. 2021](#)).

For admitted patient data, the definitions for mental health and self-harm behaviours include:

- ‘Any mental health’ hospitalisations are defined as any diagnosis (principal, secondary, etc.) of a mental disorder (ICD-10-AM codes F00-F99).
- Intentional self-harm hospitalisations are defined based on the ICD-10-AM principal diagnosis in the range S00-T75 or T79 and has a principal external cause code in the range X60-X84 or Y87.0.
- Suicidal ideation hospitalisations (ICD-10-AM code R45.81) are grouped with ‘any mental health’ and intentional self-harm hospitalisations, as this code is usually coded in the absence of a mental health condition.

For ED presentation data, the definitions ‘mental health-related ED presentations’ refers to presentations that have a principal diagnosis that falls within the *Mental and behavioural disorders* chapter (Chapter 5) of ICD-10-AM (codes F00-F99). It should be noted that this definition does not encompass all mental health-related presentations to ED. See [Mental health services in Australia](#) (<https://www.aihw.gov.au/reports/mental-health-services/mental-health-services-in-australia>) for further information.

Note that diagnosis codes for intentional self-harm sit outside the *Mental and behavioural disorders* chapter (X60-X84). Additionally, an ED presentation for self-harm may have a principal diagnosis relating to the injury. These presentations cannot be identified as mental health-related presentations and are not included in this analysis (AIHW 2022).

Presentations to hospital emergency departments relating to suicide attempts or intentional self-harm cannot be easily identified in the current national emergency department data collection. Furthermore, ICD-10-AM diagnosis codes for intentional self-harm do not specify if there was suicidal intent or not— and therefore includes both suicide attempts and non-suicidal self-harming behaviours (AIHW 2022b). See [Suicide & self-harm monitoring: Intentional self-harm hospitalisations](#). (<https://www.aihw.gov.au/suicide-self-harm-monitoring/data/intentional-self-harm-hospitalisations>) for further information.

‘Any hospitalisation’ refers to any acute admitted/mental health care separation or ED presentation.

Limitations of this analysis includes:

- Mental health items could be miscoded or reported, for example, GP mental health services are typically billed under general GP consultations.
- Service use captured in the NIHSI is influenced by severity of condition, a person’s ability and desire to access a service, and the availability of alternative services not captured in the data (e.g. private community mental health services).

## References

Australian Institute of Health and Welfare (AIHW) (2022a) Deaths in Australia (<https://www.aihw.gov.au/reports/life-expectancy-death/deaths-in-australia>), AIHW, Australian Government, accessed 11 October 2022.

Australian Institute of Health and AIHW Welfare (2022b) Mental health services in Australia, AIHW, Australian Government, accessed 09 September 2022

AIHW (2022c) Suicide and self-harm monitoring: Intentional self-harm hospitalisations (<https://www.aihw.gov.au/suicide-self-harm-monitoring/data/intentional-self-harm-hospitalisations/intentional-self-harm-hospitalisations-by-age-sex>), AIHW, Australian Government, accessed 14 October 2022.

Clapperton A, Dwyer J, Millar C, Tolhurst P and Berecki-Gisolf J (2021) 'Sociodemographic characteristics associated with hospital contact in the year prior to suicide: A data linkage cohort study in Victoria, Australia', PLoS ONE, 16(6): e0252682, doi:10.1371/journal.pone.0252682.

## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on [13 11 14](tel:131114), or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)

## Codes and classifications

### International Statistical Classification of Diseases (ICD) and Related Health Problems

The ICD, which was developed by the World Health Organization (WHO), is the international standard for coding morbidity and mortality statistics. It was designed to promote international comparability in collecting, processing, classifying and presenting these statistics. The ICD is periodically reviewed to reflect changes in clinical and research settings.

For *Suicide & self-harm monitoring*, deaths since 1964 (included in the NMD) classified as 'intentional self-harm' according to the relevant revisions of the ICD classification were included:

Table 1: Versions of ICD and years applicable in Australia

| ICD version   | Years applicable | Intentional self-harm codes |
|---------------|------------------|-----------------------------|
| 7th revision  | 1958-1967        | E970-E979 and E963          |
| 8th revision  | 1968-1978        | E950-E959                   |
| 9th revision  | 1979-1996        | E950-E959                   |
| 10th revision | 1997 to date     | X60-X84 and Y87.0           |

For deaths prior to 1964, please see [General Record of Incidence of Mortality \(GRIM\) books](#) GRIM 2017 Intentional self-harm (suicide) X60-X84, Y87.0 for ICD versions and codes used.

### ICD-10-AM

Diagnosis, intervention and external cause data are reported to the NHMD by all states and territories using the International Statistical Classification of Diseases and Related Health Problems, 10th revision, Australian Modification (ICD-10-AM) and the Australian Classification of Health Interventions (ACHI). The Australian Coding Standards (ACS) are designed to be used in conjunction with the ICD-10-AM and ACHI to support sound coding convention.

The hospital separations reported were coded according to the applicable ICD-10-AM edition for the following years:

- 2008-09 to 2010-11: ICD-10-AM 6th edition
- 2010-11 to 2012-13: ICD-10-AM 7th edition
- 2013-14 to 2014-15: ICD-10-AM 8th edition
- 2015-16 to 2016-17: ICD-10-AM 9th edition
- 2017-18 to 2020-21: ICD-10-AM 10th edition.

Records that satisfied the following criteria were included:

- a principal diagnosis in the ICD-10-AM range S00-T75, T79 (Injury, poisoning and certain other consequences of external causes)
- the first reported external cause code in the record in the ICD-10-AM range X60-X84, Y87.0 (external causes of morbidity).

Excluded from the criteria are:



- separations for which the care type was reported as Newborn (without qualified days), and records for Hospital boarders or Posthumous organ procurement
- separations with a mode of admission of 'transfer from another hospital'
- separations with reported ICD-10-AM code Z50 (Care involving the use of rehabilitation procedures) in additional diagnosis.

Changes to the Australian Coding Standard for Rehabilitation in 1 July 2015 ICD-10-AM (9th Edition), means that the 'reason' for rehabilitation (codes S00-T98 Injury, poisoning and certain other consequences of external causes) will be assigned the principal diagnosis and the rehabilitation code (Z50) will be sequenced as the additional diagnosis. This change results in an increase in the number of separations in principal diagnoses with codes from S00-T98 from 1 July 2015 onwards. In order to reflect the number of injury separations where the primary clinical intent is acute care and not rehabilitation, records with Z50 (Care involving the use of rehabilitation procedures) in principal diagnosis or additional diagnosis for all years are excluded in the data set before and after the coding change.

Intentional self-harm hospitalisations reported in *Suicide & self-harm monitoring* may differ from other publications. The differences are small and may reflect differences in the inclusion criteria (e.g. Y87.0 included here) and/or exclusion criteria. Data may also be subject to periodic updates occurring after the original publication date.



## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on [13 11 14](tel:131114), or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)

## Methods

### Crude rates

A crude rate provides information on the number of events relative to the population 'at risk' (for example, the entire population) in a specified period based on the Australian estimated resident population for the relevant analysis year. No age adjustments are made when calculating such a rate. Crude rates are used throughout this publication and expressed per 100,000 population.

### Age-specific rates

Age-specific rates are calculated by dividing the number of events (for example, deaths) in each specified age group, by the total population at risk of the event in the same age group. Where age-specific rates are reported they are expressed per 100,000 population.

### Age-standardised rates

Age-standardised rates are incidence rates that enable comparisons between populations that have different age structures and over time as the age structure of the population of interest may change. This effectively removes the influence of the age structure on the summary rate—it is the overall death rate that would have prevailed in the standard population if it had experienced at each age the death rates of the population under study.

Direct standardisation was used in this report. To calculate age-standardised rates, age-specific rates (grouped in 5-year intervals) were multiplied against a standard population. Directly age-standardised rates were adjusted using the current Australian standard population (that is, the non-recast Australian estimated resident population (ERP) as at 30 June 2001).

Rates are expressed as per 100,000 per population years.

### Standardised mortality ratio

Standard mortality ratio (SMR) is a widely recognised measure used to account for differences in age structures when comparing death rates between populations. This method of standardisation can be used when analysing relatively rare events (i.e. where number of deaths is less than 25 for the analysed time period) (Curtin and Klein, 1995). The SMR has been used in the analysis of Australian Defence Force (ADF) deaths by suicide. It is used to control for the fact that the 3 ADF service status groups have a younger age profile than the Australian population, and rates of suicide vary by age in both the study populations and the Australian population. The SMRs control for these differences, enabling comparisons of suicide counts between the 3 service status groups and Australia without the confounding effect of differences in age. The SMR is calculated as the observed number of events (deaths by suicide) in the study population divided by the number of events that would be expected if the study population had the same age and sex specific rates as the comparison population.

### Geography

Geographic location data are based on the area of usual residence of the deceased in the NMD or admitted patient in the NHMD. These data are specified using Statistical Area Level 2 (SA2) of the Australian Bureau of Statistics (ABS) Australian Statistical Geography Standard (ASGS) Edition 2016 for all states and territories. From 2016-17, the area of usual residence in the NHMD was voluntarily provided by some jurisdictions in the form of a Statistical Area level 1 (SA1).

### Remoteness areas

Data for remoteness areas are based on a person's usual residence, rather than where they died (NMD) or received treatment (NHMD). Data by remoteness are aligned to the 2016 Australian Statistical Geography Standard (ASGS) Remoteness Area Structure. Correspondence files are sourced from Australian Statistical Geography Standard (ASGS): Volume 1 - Main Structure and Greater Capital City Statistical Areas (ABS cat. no. 1270.0.55.001). The 2016 ASGS Remoteness Structure categorises geographic areas in Australia into 5 classes of remoteness areas based on their relative access to services using the Accessibility/Remoteness Index of Australia which is, in turn, derived by measuring the road distance of a location from the nearest urban centre. The 5 classes are: *Major cities*, *Inner regional*, *Outer regional*, *Remote*, and *Very remote*. See the [Australian Statistical Geography Standard \(ASGS\): Remoteness Structure, 2016](https://www.abs.gov.au/ausstats/abs@.nsf/mf/1270.0.55.005) (<https://www.abs.gov.au/ausstats/abs@.nsf/mf/1270.0.55.005>) for further information on Remoteness areas including details of the nature of the changes between the ASGS 2011 and ASGS 2016.

## Socioeconomic status

The Socio-Economic Indexes for Areas (SEIFA) is a suite of 4 summary measures, developed by the ABS based on Census data that ranks geographic areas across Australia in terms of their relative socioeconomic advantage and disadvantage. The SEIFA index used is the 2016 SEIFA Index of Relative Socioeconomic Disadvantage (IRSD) for use at Statistical Area Level 2 except for NHMD 2012-13 to 2016-17 data which uses the 2011 SEIFA IRSD.

The IRSD includes only measures of relative disadvantage. A low score indicates greater disadvantage in general (for example, an area has many households with low income, many people with no qualifications and many people working in low skill occupations). A high score indicates a relative lack of disadvantage in general (for example, an area has few households with low incomes, few people with no qualifications and few people working in low skilled occupations). It is important to understand that a high score reflects a relative lack of disadvantage rather than advantage and that the IRSD relates to the average disadvantage of all people living in a geographic area and does not reflect the socioeconomic status of all individuals living within the area.

Population-based Australian cut-offs for SEIFA quintiles have been used in this report. Population-based quintiles are calculated by dividing SEIFA areas into 5 equal groups in such a way that the population in each group is approximately equal. As SEIFA measures the characteristics of an area rather than individuals, the population in the most disadvantaged population-based quintile ('1–Lowest') is the 20% of the national population residing in the most disadvantaged areas, rather than the most disadvantaged 20% of the population.

See the [Census of Population and Housing: Socio-Economic Indexes for Areas \(SEIFA\) Australia, 2016](https://www.abs.gov.au/ausstats/abs@.nsf/mf/2033.0.55.001) (<https://www.abs.gov.au/ausstats/abs@.nsf/mf/2033.0.55.001>) for further information on SEIFA.

## Primary Health Network

Primary Health Networks (PHNs) were established in 2015 by the Department of Health to commission medical services and improve the coordination of care for patients across specific geographic areas (PHN areas). There are 31 PHN areas that cover the whole of Australia.

Statistics for PHN areas are derived by aligning deaths or hospitalisations area of usual residence data at Statistical Area Level 2 (SA2) to the 2017 PHN structure using ABS correspondence files, sourced from [Australian Statistical Geography Standard \(ASGS\): Volume 3 - Non ABS Structures, July 2018](https://www.abs.gov.au/ausstats/abs@.nsf/mf/1270.0.55.003) (ABS cat. no. 1270.0.55.003) (<https://www.abs.gov.au/ausstats/abs@.nsf/mf/1270.0.55.003>).

## Statistical Areas

Statistical Areas are a geographic classification defined by the Australian Bureau of Statistics. They encompass 4 levels, with increasing size and population: Statistical Areas Level 1 (SA1s); Statistical Areas Level 2 (SA2s); Statistical Areas Level 3 (SA3s); and Statistical Areas Level 4 (SA4).

Deaths by suicide and hospitalisations for intentional self-harm data at Statistical Area Level 2 (SA2) were aligned to Statistical Area Level 3 (SA3) and 4 (SA4) geographies based on the 2016 Australian Statistical Geography Standard (ASGS) structure. Correspondence files are sourced from [Australian Statistical Geography Standard \(ASGS\): Volume 1 - Main Structure and Greater Capital City Statistical Areas](https://www.abs.gov.au/ausstats/abs@.nsf/mf/1270.0.55.001) (ABS cat. no. 1270.0.55.001) (<https://www.abs.gov.au/ausstats/abs@.nsf/mf/1270.0.55.001>).

## Using confidence intervals to test for statistical significance

Statistical significance is a measure that indicates how likely it is that an observed difference, or a larger one, would occur under the conditions of the null hypothesis.

In the analysis of deaths by suicide in Australian Defence Force personnel, 95% confidence intervals (CIs) are provided for each standardised mortality ratio to indicate the level of uncertainty around these estimates due to random fluctuations in the number of deaths by suicide over time. Estimates produced using low numbers can be sensitive to small changes in numbers of deaths over time and will therefore have wide CIs. 95% CIs are provided within this report as they may account for the variation in absolute numbers of deaths by suicide over time (related to the small sample size). It is important to note that there are other sources of uncertainty, such as linkage error, that are not captured by the provided CIs.

Use of CIs is the simplest way to test for significant differences between service groups and Australian comparison groups. For the purpose of this monitoring site, differences are deemed to be statistically significant if CIs do not overlap with 1.0 in the case of an SMR. The CIs in this report cannot be used to determine the significance of differences over time between overlapping 3-year time periods.

## References

ABS (Australian Bureau of Statistics) 2019. Microdata: Multi-Agency Data Integration Project, Australia, March 2019. Cat. 1700.0.

ABS 2018. Multi-Agency Data Integration Project (MADIP) Research Projects

([https://www.abs.gov.au/ausstats/abs@.nsf/Lookup/by%20Subject/1700.0-Australia-Main%20Features-MADIP%20Basic%20Longitudinal%20Extract,%202011-2016%20\(2011-2016%20Cohorts\)-10000](https://www.abs.gov.au/ausstats/abs@.nsf/Lookup/by%20Subject/1700.0-Australia-Main%20Features-MADIP%20Basic%20Longitudinal%20Extract,%202011-2016%20(2011-2016%20Cohorts)-10000)).

ABS 2016. Research Paper: Death Registrations to Census Linkage Project - A Linked Dataset for Analysis, Mar 2016. Cat. 1351.0.55.058.

Curtin, LR, & Klein, R J 1995. *Direct standardization (age-adjusted death rates)* (No. 6). Hyattsville, MD: US Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Center for Health Statistics.

Lubman DI, Heilbronn C, Ogeil RP, Killian JJ, Matthews S, Smith K, et al. 2020. National Ambulance Surveillance System: A novel method using coded Australian ambulance clinical records to monitor self-harm and mental health-related morbidity

(<https://doi.org/10.1371/journal.pone.0236344>). PLoS ONE 15(7): e0236344.

---

© Australian Institute of Health and Welfare 2024





## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on [13 11 14](tel:131114), or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)

## Data downloads

---

## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on [13 11 14](tel:131114), or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)

## Notes

### Latest data updates

For information on future planned updates to the publication see [Data update schedule](#).

#### 18 April 2024

- Data from suicide registers

#### 14 March 2024

- Data from suicide registers

#### 18 January 2024

- Data from suicide registers

#### 12 December 2023

- LGBTIQ+ Australians: suicidal thoughts and behaviours and self-harm
- Data from suicide registers

#### 22 November 2023

- Suicide among refugee and humanitarian entrants and other permanent migrants
- LGBTIQ+ Australians: suicidal thoughts and behaviours and self-harm
- Australian Defence Force suicide monitoring
- Fact sheets and key messages

#### 27 October 2023

- Deaths by suicide in Australia
- Topic Summary
- Data from suicide registers

#### 6 October 2023

- Data from suicide registers

#### 28 September 2023

- Youth Self-Harm Atlas
- Ambulance attendances: suicidal and self-harm behaviours
- Intentional self-harm hospitalisations 2021-22
- Featured report - Feasibility Study for Identifying Suicide Clusters Using Real-time Coronial Data

#### 5 September 2023

- Data from suicide registers

#### 13 July 2023

- Data from suicide registers

#### 7 July 2023

- Suicidal and self-harming thoughts and behaviours among LGBTIQ+ Australians
- Ambulance attendances: suicidal and self-harm behaviours

#### 15 June 2023

- Data from suicide registers

#### 16 May 2023

- Data from suicide registers

#### 6 April 2023

- Impact of suicide among Aboriginal and Torres Strait Islander Australians
- Ambulance attendances: suicidal ideation and self-harm behaviours
- Deaths by suicide among people who used disability service
- Burden of disease studies - Suicide & self-inflicted injuries
- Data from suicide registers

#### 17 February 2023

- Data from suicide registers

#### 17 January 2023

- Data from suicide registers

#### 12 December 2022

- ANU paper: Spatiotemporal Analysis of Suicide Deaths 2001 - 2020

#### 6 December 2022

- Australian Defence Force suicide monitoring

#### 18 November 2022

- Data: Deaths by Suicide in Australia

#### 4 November 2022

- Data from suicide registers

#### 5 October 2022

- Data from suicide registers

#### 8 September 2022

- Data from suicide registers

#### 5 August 2022

- Data from suicide registers
- Australian prevalence estimates of suicidal behaviours

#### 26 July 2022

- Research & information | Releases | Featured reports - Evaluation of the National Suicide and Self-harm Monitoring Project and System| Final Report
- Behaviours & risk factors - Longitudinal analysis of income uncertainty & suicide (MADIP data asset)
- Intentional self-harm hospitalisations 2020-21
- Ambulance attendances - suicidal and self-harm behaviours
- Research & information | Releases | Consultations - Data Requirements for the Portal

#### 8 July 2022

- Data from suicide registers
- COVID-19 - The use of mental health services, psychological distress, loneliness, suicide, ambulance attendances and COVID-19
- Australia's health 2022: Suicide & intentional self-harm

## 8 June 2022

- Data from suicide registers

## 27 April 2022

- Data from suicide registers
- COVID-19 - The use of mental health services, psychological distress, loneliness, suicide, ambulance attendances and COVID-19

## 3 February 2022

- Data from suicide registers

## 9 December 2021

- Data from suicide registers
- COVID-19 - The use of mental health services, psychological distress, loneliness, suicide, ambulance attendances and COVID-19
- Research & information | Releases | Featured Reports - A scoping review of analytic methods used within the peer reviewed literature
- Research & information | Releases | Featured Reports - Addendum | Suicide mortality in Australia: Estimating and projecting monthly variation and trends from 2007 to 2018 and beyond
- Research & information | Releases | Consultations - Consultation with young people - Suicide & self-harm monitoring website

## 8 December 2021

- Ambulance attendances: suicidal and self-harm behaviours
- Research & information | Releases | Featured Reports - Patterns of suicide in the context of COVID-19: Evidence from three Australian states

## 4 November 2021

- Data from suicide registers
- The health impact of suicide and self-inflicted injuries in Australia, 2019

## 15 October 2021

- Ambulance attendances: suicidal and self-harm behaviours
- Behaviours & risk factors - Social factors & suicide (MADIP data asset)
- COVID-19 - The use of mental health services, psychological distress, loneliness, suicide, ambulance attendances and COVID-19

## 30 September 2021

- Deaths by suicide in Australia, Populations & age groups, Behaviours & Risk Factors [ABS Causes of Death 3303.0]
- Populations & age groups - Australian Defence Force suicide monitoring
- Data from suicide registers
- Geography - Intentional self-harm hospitalisations by local areas

## 1 September 2021

- Data from suicide registers

## 20 July 2021

- Deaths by suicide in Australia - Deaths by suicide over time
- Data from suicide registers
- COVID-19 - The use of mental health services, psychological distress, loneliness, suicide, ambulance attendances and COVID-19
- Ambulance attendances: suicidal and self-harm behaviours
- Populations & age groups - Suicide & Indigenous Australians
- Intentional self-harm hospitalisations (all pages excluding Intentional self-harm hospitalisations by local areas)
- Geography - International estimates of death by self-harm

## 30 March 2021

- Victoria & New South Wales Suicide Register data
- The use of mental health services, psychological distress, loneliness, suicide, ambulance attendances and COVID-19

## 18 November 2020

- Mortality data; Geography - Suicide by PHN areas
- Victoria and New South Wales Suicide Register data; COVID-19 - Data from suicide registers

## 9 November 2020

- Mortality data; Death by suicide in Australia; Populations & age groups; Geography; Behaviours & risk factors



9 October 2020

- Populations & age groups - Australian Defence Force suicide monitoring
- 


## Amendments

28 July 2022

- Ambulance attendances - Ambulance attendances: suicidal and self-harm behaviours

9 November 2020

- Populations & age groups - Suicide among young people.
- 

© Australian Institute of Health and Welfare 2024 

## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on [13 11 14](tel:131114), or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)

## Glossary

**Aboriginal or Torres Strait Islander:** A person of Aboriginal and/or Torres Strait Islander descent who identifies as an Aboriginal and/or Torres Strait Islander. See also [Indigenous](#).

**additional diagnosis:** The diagnosis of a condition or recording of a complaint—either coexisting with the principal diagnosis or arising during an episode of admitted patient care (hospitalisation)—that requires the provision of care. Multiple diagnoses may be recorded.

**ADF personnel:** Serving, reserve and ex-serving members of the Australian Defence Force; civilian personnel employed by the Department of Defence are excluded.

**admission:** An admission to hospital. The term hospitalisation is used to describe an episode of hospital care that starts with the formal admission process and ends with the formal separation process.

**administrative data collection:** A data set that results from the information collected for the purposes of delivering a service or paying the provider of the service. This type of collection is usually complete (all in-scope events are collected), but it may have limitations for population-level analysis because the data are collected primarily for an administrative purpose.

**age structure:** The relative number of people in each age group in a population.

**age-specific rate:** The number of events for a specified age group over a specified period (e.g. calendar or financial year), divided by the total population in that age group. Reported as number per 100,000. The numerator and denominator relate to the same age group.

**age-standardised rates:** are incidence rates that enable comparisons to be made between populations that have different age structures. The age structures of the different populations are converted to the same 'standard' structure, and then the rates that would have occurred with that structure are calculated and compared. Rates are expressed as per 100,000 per population years.

**associated cause(s) of death:** All causes of death listed on the death certificate, other than the [underlying cause of death](#). They include the immediate cause, any intervening causes, and conditions which contributed to the death but were not related to the disease or condition causing the death.

**attributable burden:** The disease burden attributed to a particular risk factor. It is the amount of burden that could be avoided if the risk factor were removed or reduced to the lowest possible exposure.

**Australian Statistical Geography Standard (ASGS):** Common framework defined by the Australian Bureau of Statistics (ABS) for collecting and disseminating geographically classified statistics. It replaced the Australian Standard Geographical Classification (ASGC) in July 2011.

**burden of disease:** The quantified impact of a disease, injury or risk factor on a population, using the [disability-adjusted life year \(DALY\)](#) measure. One DALY is one year of 'healthy life' lost due to illness and/or death. The more DALY associated with a disease or injury, the greater the burden. The DALY is produced by combining the non-fatal and fatal burden together. People generally experience more burden as they age.

**cause(s) of death:** All diseases, morbid conditions or injuries that either resulted in or contributed to death—and the circumstances that produced any such injuries—that are entered on the death certificate. The coding of causes of death produces an [underlying cause of death](#) and, for many deaths, one or more [associated cause\(s\) of death](#). See also [multiple causes of death](#).

**child:** A person aged 0-14 years.

**comorbidity:** The occurrence of 2 or more health conditions in a person at one time. While the coexistence of these multiple conditions may be unrelated, in many instances there is some association between them.

**confidence interval:** A statistical term describing a range (interval) of values within which we can be 'confident' that the true value lies, usually because it has a 95% or higher chance of doing so.

**contemporary ex-serving (Australian Defence Force):** Australian Defence Force members who have had at least 1 day of full-time or reserve service on or after 1 January 2001, and have since been discharged from the Australian Defence Force.

**current serving (Australian Defence Force):** Australian Defence Force members who have had at least 1 day of full-time service on or after 1 January 2001, and are still serving in the Australian Defence Force.

**crude rate:** The crude rate is the number of events recorded during a specified time period (e.g. calendar year) per 100,000 estimated resident population.

**DALY:** See **disability-adjusted life year**.

**data linkage:** The process of combining (linking) information from two or more different data sources that are believed to relate to the same entity (for example, the same individual or the same institution). This linkage can yield more information about the entity and, in certain cases, provide a time sequence—helping to 'tell a story', show 'pathways' and perhaps unravel cause and effect. The term is used synonymously with 'record matching' and 'data integration'.

**death:** Any death which occurs in, or en route to Australia and is registered with a State or Territory Registry of Births, Deaths and Marriages.

**determinant:** Any factor that influences how likely a population or individual will stay healthy or become ill or injured. Factors that increase the chances of ill health are known as risk factors, while those that promote good health are protective factors. Services or other programs that aim to improve health are usually not included in this definition.

**disability-adjusted life year (DALY):** A measure of healthy life lost, either through premature death or living with disability due to illness or injury. It is the basic unit used in burden of disease and injury estimates.

**episode of care:** The period of admitted patient care between a formal or statistical admission and a formal or statistical separation, characterised by only one care type (see care type and **separation**).

**estimated resident population (ERP):** The official ABS estimate of the Australian population. The ERP is derived from the 5-yearly Census counts and is updated quarterly between each Census. It is based on the usual residence of the person. Rates are calculated per 1,000 or 100,000 mid-year (30 June) ERP.

**external cause:** The environmental event, circumstance, or condition that is regarded as the cause of injury, poisoning and other adverse effect.

**fatal burden:** The quantified impact on a population of dying prematurely due to disease or injury, measured by years of life lost (YLL).

**hospitalisation:** An episode of admitted patient care, which can be a total hospital stay (from admission to discharge, transfer or death) or a portion of a hospital stay beginning or ending in a change of type of care (e.g. from acute care to rehabilitation).

**incidence:** Incidence is a measure of the number of new cases of a characteristic that develop in a population in a specified time period; whereas prevalence is the proportion of a population who have a specific characteristic in a given time period, regardless of when they first developed the characteristic.

**incidence rates:** incidence rates for death by suicide refers to the number of suicides during a specified period over the population within the same period. Rates are expressed as per 100,000 per population years.

**Index of Relative Socioeconomic Disadvantage (IRSD):** One of the set of **Socio-Economic Indexes for Areas (SEIFA)** for ranking the average socioeconomic conditions of a population in a geographic area. The IRSD was developed by the ABS for use at Statistical Area Level 2 and summarises attributes of the population that indicate disadvantage, such as low income, low educational attainment, high unemployment and jobs in relatively unskilled occupations.

**Indigenous:** A person of Aboriginal and/or Torres Strait Islander descent who identifies as an Aboriginal and/or Torres Strait Islander. See also **Aboriginal or Torres Strait Islander**.

**intentional self-harm:** Includes attempts to suicide, as well as cases where people have intentionally hurt themselves, but not necessarily with the intention of suicide (e.g. acts of self-mutilation).

**International Statistical Classification of Diseases and Related Health Problems (ICD):** The World Health Organization's internationally accepted classification of death and disease. The 10th Revision (ICD-10) is currently in use. The ICD-10-AM is the Australian Modification of the ICD-10; it is used for diagnoses and procedures recorded for patients admitted to hospitals.

**monitoring (of public health):** A process of keeping a regular and close watch over important aspects of the public's health and health services through various measurements, and then regularly reporting on the situation, so that the health system and society more generally can plan and respond accordingly. The term is often used interchangeably with surveillance, although surveillance may imply more urgent watching and reporting, such as the surveillance of infectious diseases and their epidemics.

**morbidity:** The ill health of an individual and levels of ill health in a population or group.

**mortality:** Number or rate of deaths in a population during a given time period.

**multiple causes of death:** All causes listed on the death certificate. This includes the underlying cause of death and all associated causes of death. This information is useful for describing the role of all diseases involved in deaths, where there is more than one cause contributing to the death. For deaths where the underlying cause was identified as an external cause multiple causes include circumstances of injury, the nature of injury as well as any other conditions reported on the death certificate.

**non-fatal burden:** The quantified impact on a population of ill health due to disease or injury, measured as years lived with disability (YLD).

**non-Indigenous:** People who have declared that they are not of Aboriginal or Torres Strait Islander descent.

**prevalence:** The number or proportion (of cases, instances, and so forth) in a population at a given time.

**prevention (of suicide):** Action to reduce or eliminate the onset, causes, complications or recurrence of suicide.

**Primary Health Networks (PHNs):** Primary Health Networks were established on 1 July 2015 by the Australian Government Department of Health. They are independent primary health care organisations that commission services and are operated by not-for-profit companies, informed by clinical councils and community advisory committees.

**Primary Health Network (PHN) areas:** PHNs connect health services across a specific geographic area (a PHN area), with the boundaries defined by the Australian Government Department of Health. There are 31 PHN areas that cover the whole of Australia.

**principal diagnosis:** The diagnosis established after study to be chiefly responsible for occasioning an episode of admitted patient care (hospitalisation). Diagnoses are recorded using the relevant edition of the International statistical classification of diseases and related health problems, 10th revision, Australian modification (ICD-10-AM).

**protective factors:** Factors that enhance the likelihood of positive outcomes and reduce the chance of negative consequences from exposure to risk.

**psychological distress:** Psychological distress is commonly measured using the Kessler Psychological Distress Scale—10 items (K10). The K10 questionnaire was developed to yield a global measure of psychosocial distress, based on questions about people's level of nervousness, agitation, psychological fatigue and depression in the past four weeks. The Kessler 6 Scale is an abbreviated version of K10.

**psychosocial factors:** Social processes and social structures which can have an interaction with individual thought, behaviour and/or health outcomes.

**public health:** Activities aimed at benefiting a population, with an emphasis on prevention, protection and health promotion as distinct from treatment tailored to individuals.

**quintile:** A group derived by ranking the population or area according to specified criteria and dividing it into five equal parts. Commonly used to describe socioeconomic areas.

**rate:** A rate is one number (the numerator) divided by another number (the denominator). The numerator is commonly the number of events in a specified time. The denominator is the population 'at risk' of the event. Rates (crude, age-specific and age-standardised) are generally multiplied by a number such as 100,000 to create whole numbers.

**remoteness area:** A classification of the remoteness of a location using the Australian Statistical Geography Standard Remoteness Area Structure (2016) which divides Australia into 5 classes of remoteness based on their relative access to services using the Accessibility and Remoteness Index of Australia which is, in turn, derived by measuring the road distance of a location from the nearest urban centre. The 5 Remoteness Areas are Major cities, Inner regional, Outer regional, Remote and Very remote.

**reserve (Australian Defence Force):** Australian Defence Force members who have had at least 1 day of reserve service on or after 1 January 2001.

**risk factor:** Any attributes, characteristics or exposures that increase the likelihood of a person developing a health condition or experiencing an event.

**separation (from hospital):** An episode of care for an admitted patient, which can be a total hospital stay (from admission to discharge, transfer or death) or a portion of a hospital stay beginning or ending in a change of type of care (for example, from acute care to rehabilitation). Separation also means the process by which an admitted patient completes an episode of care either by being discharged, dying, transferring to another hospital or changing type of care.

**social determinants of health:** The circumstances in which people are born, grow up, live, work and age, and the systems put in place to deal with illness. These circumstances are in turn shaped by a wider set of forces: economics, social policies and politics.

**socioeconomic status:** The social and economic position of an individual or group within the larger society. In this monitoring site, socioeconomic status is reported using the Socio-Economic Indexes for Areas, typically for 5 groups, from the most disadvantaged (lowest socioeconomic status areas) to the least disadvantaged (highest socioeconomic status areas).

**Socio-Economic Indexes for Areas (SEIFA):** A set of indexes, created from Census data, that represent the socioeconomic status of geographical areas in Australia according to their relative socioeconomic advantage and disadvantage. The SEIFA index used in this report is the **Index of Relative Socioeconomic Disadvantage (IRSD)**. It is important to understand that the index value reflects the overall or average level of disadvantage of the population of an area; it does not reflect the socioeconomic status of individuals living within the area.

**Socio-Economic Indexes for Areas (SEIFA) quintiles:** Population-based quintiles are calculated by dividing SEIFA areas into 5 equal groups in such a way that the population in each group is approximately equal. As SEIFA measures the characteristics of an area rather than individuals, the population in the most disadvantaged population-based quintile ('1–Lowest') is the 20% of the national population residing in the most disadvantaged areas, rather than the most disadvantaged 20% of the population.

**statistical areas:** A geographical classification defined by the ABS. They encompass four levels, with increasing size and population: Statistical Areas Level 1 (SA1s); Statistical Areas Level 2 (SA2s); Statistical Areas Level 3 (SA3s); and Statistical Areas Level 4 (SA4s).

**statistical significance:** A statistical measure indicating how likely the observed difference or association is due to chance alone. Rate differences are deemed to be statistically significant when their confidence intervals do not overlap, since their difference is greater than what could be explained by chance.

**suicidal ideation:** Serious thoughts about ending one's own life.

**suicidal behaviours:** The collective term for suicidal ideation, suicide plans and suicide attempts.

**suicide:** An action intended to deliberately end one's own life.

**total burden:** The sum of fatal burden (YLL) and non-fatal burden (YLD).

**underlying cause of death:** The disease or injury that initiated the train of events leading directly to a person's death, or the circumstances of the accident or violence that produced the fatal injury. See also **cause(s) of death** and **associated cause(s) of death**.

**usual residence:** The area of the address at which the deceased lived or intended to live, for 6 months or more prior to death.

**years lived with disability (YLD):** The number of years of what could have been a healthy life that were instead spent in states of less than full health. YLD represent non-fatal burden.

**years of life lost (YLL):** The number of years of life lost due to premature death, defined as dying before the ideal life span. YLL represent fatal burden.

## Viewing the monitoring data

Caution: Some people may find parts of this content confronting or distressing.

Please carefully consider your needs when reading the following information about suicide and self-harm. If this material raises concerns for you contact Lifeline on [13 11 14](tel:131114), or [see other ways you can seek help](#).

The information included here places an emphasis on data, and as such, can appear to depersonalise the pain and loss behind the statistics. The AIHW acknowledges the individuals, families and communities affected by suicide each year in Australia.

Aboriginal and Torres Strait Islander readers are advised that information relating to Indigenous suicide and self-harm is included.

The AIHW supports the use of the [Mindframe guidelines](#) on responsible, accurate and safe suicide and self-harm reporting. Please consider these guidelines when reporting on statistics on the monitoring of suicide and self-harm.

## Suicide & self-harm monitoring

### Need help now?

Lifeline 13 11 14

More (</suicide-self-harm-monitoring/research-information/crisis-support>)

## Archived content

For the latest data, please see [Data downloads](#).

The below data tables contain previously published data that have now been superseded.

### Notes for archived data downloads

- **National Hospital Morbidity Database—Intentional self-harm hospitalisations**

The estimated resident populations used in rates calculations throughout this data table have been revised in more recent updates.

- **National Mortality Database—Suicide (ICD-10 X60-X84, Y87.0)**

The estimated resident populations used in rates calculations throughout this data table have been revised in more recent updates.

---