

Indicator 5.3: Incidence by location

Table 94: New cases of cervical cancer by age and location, 1995–1998 and 1999–2002

Age group	Major cities		Regional		Remote		Australia	
	1995–1998	1999–2002	1995–1998	1999–2002	1995–1998	1999–2002	1995–1998	1999–2002
0–4	0	0	0	0	0	0	0	0
5–9	0	0	0	0	0	0	0	0
10–14	0	0	0	0	0	0	0	0
15–19	4	3	2	1	0	0	6	4
20–24	23	25	14	10	2	1	40	36
25–29	123	120	58	54	8	5	190	179
30–34	212	192	120	85	11	11	344	290
35–39	294	209	142	114	20	14	458	336
40–44	305	207	123	111	10	11	439	328
45–49	275	215	108	102	9	14	392	332
50–54	190	198	83	69	9	5	283	272
55–59	161	122	63	73	11	9	235	204
60–64	155	135	76	70	8	6	240	212
65–69	174	118	76	69	6	4	256	191
70–74	168	127	62	50	4	2	235	179
75–79	128	117	59	47	5	1	191	166
80–84	97	96	43	45	3	1	144	142
85+	87	69	27	33	2	2	115	104
All ages	2,395	1,952	1,056	933	108	86	3,568	2,975
Target age								
20–69 years	1,912	1,541	863	756	94	79	2,877	2,380

Notes

1. The numbers are presented as 4-year rolling blocks of data.
2. In the periods 1995–1998 and 1999–2002 there were 9 and 4 cases respectively that were excluded from these data because the respective postcodes were not able to be matched to the coding used for this analysis.

Source: National Cancer Statistics Clearing House (AIHW).

Table 95: Age-specific and age-standardised incidence rates for cervical cancer by age and location, 1995–1998 and 1999–2002

Age group	Major cities		Regional		Remote		Australia	
	1995–1998	1999–2002	1995–1998	1999–2002	1995–1998	1999–2002	1995–1998	1999–2002
0–4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
5–9	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
10–14	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
15–19	0.2	0.2	0.3	0.1	0.0	0.0	0.2	0.2
20–24	1.2	1.3	2.0	1.6	2.8	1.6	1.5	1.4
25–29	6.1	5.9	7.7	7.4	9.6	5.9	6.7	6.3
30–34	10.9	9.4	14.2	10.4	13.1	13.9	11.9	9.9
35–39	15.2	10.3	15.5	12.5	25.8	17.4	15.6	11.2
40–44	16.7	10.6	14.4	11.9	14.5	15.1	16.0	11.1
45–49	15.9	11.9	14.0	12.0	16.6	23.4	15.4	12.2
50–54	13.6	11.7	12.9	8.9	19.6	9.0	13.6	10.8
55–59	14.9	9.6	11.7	11.4	30.5	21.5	14.2	10.4
60–64	16.5	13.1	15.8	12.7	30.0	20.2	16.6	13.2
65–69	18.9	13.2	16.3	14.6	27.8	16.7	18.2	13.7
70–74	19.1	14.5	14.9	11.3	23.7	12.6	17.9	13.4
75–79	19.0	15.2	18.6	13.0	36.1	7.5	19.1	14.4
80–84	20.3	18.1	19.6	18.0	37.4	15.8	20.3	18.1
85+	21.9	14.2	15.1	14.9	19.2	24.3	19.8	14.5
All ages								
AS rate (A)	9.7	7.4	9.3	7.6	13.9	10.1	9.7	7.5
95% CI	9.4–10.1	7.0–7.7	8.7–9.9	7.1–8.1	11.3–16.9	8.0–12.5	9.4–10.0	7.2–7.8
AS rate (W)	8.2	6.2	8.0	6.5	11.7	8.7	8.2	6.4
95% CI	7.9–8.5	5.9–6.5	7.5–8.5	6.1–7.0	9.6–14.2	6.9–10.7	7.9–8.5	6.1–6.6
Target age 20–69 years								
AS rate (A)	12.5	9.4	12.2	10.1	17.8	14.0	12.5	9.7
95% CI	12.0–13.1	8.9–9.9	11.4–13.1	9.3–10.8	14.2–21.7	11.0–17.5	12.1–13.0	9.3–10.1
AS rate (W)	11.9	9.0	11.7	9.7	16.9	13.3	11.9	9.3
95% CI	11.3–12.4	8.5–9.4	11.0–12.6	9.0–10.4	13.6–20.7	10.5–16.6	11.5–12.4	8.9–9.7

Notes

1. The numbers are presented as 4-year rolling blocks of data.
2. Rates are expressed per 100,000 women and age-standardised to the Australian 2001 population (A) and the WHO World Standard Population (W).

Source: National Cancer Statistics Clearing House (AIHW).

Indicator 6.1: Mortality

Table 96: Deaths from cervical cancer by age, Australia, 1984–2004

Age group	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	00	01	02	03	04	
0–4	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
5–9	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
10–14	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
15–19	0	0	0	0	0	1	1	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1
20–24	0	0	2	2	0	1	1	3	0	0	0	0	1	0	3	1	1	0	2	0	0	0
25–29	10	6	6	5	3	3	10	5	5	2	6	3	1	2	6	2	4	1	2	5	4	4
30–34	13	20	12	15	12	21	14	13	15	11	11	7	13	8	5	6	10	11	6	13	7	7
35–39	19	17	16	20	15	18	30	25	19	25	11	16	23	18	19	7	12	12	9	12	9	9
40–44	20	18	26	20	24	24	36	19	27	32	28	21	20	16	19	18	14	19	13	12	13	13
45–49	26	21	24	19	27	31	36	29	26	23	35	32	30	28	16	25	27	23	15	22	17	17
50–54	25	25	25	24	19	27	17	21	13	29	37	26	13	21	24	15	19	21	32	17	15	15
55–59	21	31	41	32	41	20	25	25	23	20	26	34	22	24	15	14	19	20	15	19	21	21
60–64	41	41	41	28	41	33	34	33	31	25	24	30	21	22	28	15	24	25	19	21	15	15
65–69	43	52	50	46	41	54	43	35	25	30	37	37	29	30	19	21	26	20	18	20	17	17
70–74	33	43	32	55	34	48	25	37	45	38	33	43	41	36	28	30	37	28	18	23	17	17
75–79	29	29	23	29	35	29	32	30	32	28	30	30	38	32	26	26	25	30	26	29	16	16
80–84	26	26	23	20	34	24	8	22	35	24	26	27	22	27	26	19	23	28	26	21	23	23
85+	21	29	24	16	17	22	25	32	23	24	24	20	24	30	31	21	26	24	26	24	37	37
All ages	327	359	343	329	343	355	337	329	319	311	329	328	296	294	265	220	267	262	227	238	212	212
Target age 20–69 years	218	230	242	210	222	231	246	208	184	197	216	207	172	169	154	124	156	152	131	141	118	118

Notes

- Deaths were derived by year of registration.
- A comparability factor of 0.98 was applied to mortality data for years prior to 1997 because in processing deaths registered from 1 January 1997, Australia adopted the use of the Automated Coding System (ACS) and introduced ICD10 codes. The comparability factor provides a link between the two data series (i.e. pre 1997 and 1997 to 2004). Comparability factors close to 1.0 indicate there were no significant coding differences between automated ICD10 and manual ICD9 coding.

Source: AIHW Mortality Database.

Table 97: Age-specific and age-standardised death rates for cervical cancer by age, Australia, 1984–2004

Age group	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	00	01	02	03	04
0–4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
5–9	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
10–14	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
15–19	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.0	0.0	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1
20–24	0.0	0.0	0.3	0.3	0.0	0.1	0.1	0.4	0.0	0.0	0.0	0.0	0.1	0.0	0.5	0.2	0.2	0.0	0.3	0.0	0.0
25–29	1.5	0.9	0.9	0.7	0.4	0.4	1.4	0.7	0.7	0.3	0.9	0.4	0.1	0.3	0.8	0.3	0.6	0.1	0.3	0.7	0.6
30–34	2.1	3.1	1.9	2.3	1.8	3.0	2.0	1.8	2.0	1.5	1.5	0.9	1.8	1.1	0.7	0.8	1.4	1.5	0.8	1.7	0.9
35–39	3.2	2.8	2.5	3.1	2.3	2.7	4.6	3.7	2.7	3.6	1.5	2.2	3.1	2.4	2.5	0.9	1.6	1.6	1.2	1.6	1.2
40–44	4.3	3.7	5.4	3.7	4.1	3.9	5.9	2.9	4.3	5.0	4.3	3.1	2.9	2.3	2.7	2.5	1.9	2.6	1.7	1.6	1.7
45–49	6.9	5.2	5.7	4.4	6.3	6.9	7.6	5.8	4.9	3.9	5.9	5.2	4.7	4.4	2.4	3.8	4.0	3.4	2.2	3.1	2.4
50–54	7.0	7.1	6.8	6.4	4.9	7.1	4.2	5.0	3.0	6.8	8.2	5.6	2.6	3.9	4.2	2.5	3.0	3.2	4.9	2.6	2.3
55–59	5.5	8.4	11.1	8.8	11.3	5.4	6.8	7.1	6.2	5.2	6.9	8.7	5.3	5.7	3.5	3.1	4.0	4.0	2.8	3.3	3.5
60–64	11.5	11.3	11.2	7.7	11.1	9.0	9.3	9.0	8.6	7.1	6.6	8.5	5.8	6.0	7.5	3.9	6.0	6.1	4.5	4.9	3.3
65–69	14.9	17.8	16.4	14.6	12.5	15.7	12.4	10.0	7.2	8.5	10.5	10.5	8.3	8.5	5.4	6.1	7.5	5.8	5.1	5.5	4.5
70–74	13.2	16.6	12.3	20.5	12.8	18.1	9.4	13.2	15.4	12.6	10.5	13.4	12.6	11.0	8.5	9.0	11.1	8.4	5.4	7.0	5.2
75–79	16.7	16.0	11.8	14.8	17.1	13.7	14.7	13.5	14.1	12.4	13.3	13.0	15.7	12.5	9.7	9.3	8.7	10.3	8.8	9.7	5.3
80–84	23.4	22.9	19.0	15.8	26.6	17.6	5.6	14.8	23.3	14.9	15.8	15.9	12.2	15.0	14.3	10.4	12.1	13.9	12.3	9.5	10.0
85+	24.7	33.1	24.9	16.1	16.7	20.9	23.2	29.4	19.5	19.3	18.4	14.6	16.6	20.1	19.8	12.6	14.8	13.1	13.6	12.2	18.3
All ages																					
AS rate (A)	4.7	5.0	4.7	4.4	4.5	4.5	4.2	4.0	3.8	3.6	3.8	3.7	3.2	3.1	2.7	2.2	2.6	2.5	2.1	2.2	1.9
AS rate (W)	3.6	3.8	3.7	3.4	3.4	3.5	3.3	3.1	2.8	2.8	2.9	2.8	2.4	2.3	2.1	1.7	2.0	1.9	1.6	1.7	1.4
Target age 20–69 years																					
AS rate (A)	4.9	5.0	5.2	4.4	4.6	4.6	4.8	4.0	3.5	3.8	4.1	3.8	3.0	3.0	2.7	2.1	2.6	2.5	2.1	2.2	1.8
AS rate (W)	4.5	4.6	4.8	4.1	4.2	4.3	4.5	3.8	3.3	3.4	3.7	3.5	2.8	2.7	2.5	1.9	2.4	2.3	1.9	2.0	1.7

Notes: Rates are expressed per 100,000 women and age-standardised to the Australian 2001 population (A) and the WHO World Standard Population (W).

Source: AIHW Mortality Database.

Table 98: Deaths from cervical cancer by age, states and territories, 1997–2000

Age group	NSW	Vic	Qld	WA	SA	Tas	ACT	NT	Australia
0–4	0	0	0	0	0	0	0	0	0
5–9	0	0	0	0	0	0	0	0	0
10–14	0	0	0	0	0	0	0	0	0
15–19	0	0	0	0	0	0	0	0	0
20–24	2	2	1	0	0	0	0	0	5
25–29	1	4	5	2	1	0	1	0	14
30–34	7	5	5	8	3	1	0	0	29
35–39	21	12	10	6	2	3	0	2	56
40–44	26	14	14	7	0	1	3	2	67
45–49	33	22	17	9	10	0	3	2	96
50–54	34	12	18	5	4	3	2	1	79
55–59	26	12	18	9	3	2	1	1	72
60–64	25	19	23	8	5	8	0	1	89
65–69	37	14	16	12	9	4	4	0	96
70–74	45	36	21	14	8	4	1	2	131
75–79	39	28	18	7	9	4	2	2	109
80–84	34	27	12	12	7	2	0	1	95
85+	42	24	16	17	5	3	1	0	108
All ages	372	231	194	116	66	35	18	14	1,046
Target age 20–69 years	224	124	118	68	35	26	16	11	603

Notes

1. Numbers were averaged over 4 years to smooth annual variations that may occur in the smaller states and territories.
2. Deaths were derived by year and state of registration.

Source: AIHW Mortality Database.

Table 99: Age-specific and age-standardised death rates for cervical cancer by age, states and territories, 1997–2000

Age group	NSW	Vic	Qld	WA	SA	Tas	ACT	NT	Australia
0–4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
5–9	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
10–14	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
15–19	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
20–24	0.2	0.3	0.2	0.0	0.0	0.0	0.0	0.0	0.2
25–29	0.1	0.5	0.9	0.7	0.5	0.0	1.9	0.0	0.5
30–34	0.7	0.7	1.0	2.9	1.4	1.5	0.0	0.0	1.0
35–39	2.1	1.6	1.8	2.0	0.9	4.0	0.0	6.2	1.9
40–44	2.7	2.0	2.7	2.4	0.0	1.4	6.0	7.1	2.4
45–49	3.8	3.4	3.5	3.4	4.7	0.0	6.1	8.2	3.6
50–54	4.3	2.1	4.2	2.3	2.1	5.1	4.8	5.4	3.4
55–59	4.3	2.7	5.5	5.4	2.0	4.3	3.7	8.7	4.0
60–64	4.7	4.9	8.7	5.8	3.9	19.7	0.0	14.2	5.9
65–69	7.5	3.9	6.7	9.9	7.3	10.7	25.0	0.0	6.9
70–74	9.5	10.4	9.5	12.8	6.5	11.4	6.8	61.4	9.9
75–75	10.0	9.9	9.8	7.9	8.6	13.3	16.9	92.6	10.0
80–84	12.9	14.3	9.7	20.1	10.0	9.6	0.0	75.6	12.9
85+	18.5	14.0	15.1	30.8	7.9	17.5	17.9	0.0	16.7
All ages									
AS rate (A)	2.7	2.3	2.9	3.3	1.9	3.4	3.3	8.8	2.7
95% CI	2.5–3.0	2.0–2.6	2.5–3.3	2.7–3.9	1.5–2.4	2.4–4.8	1.9–5.3	4.0–15.9	2.5–2.8
AS rate (W)	2.1	1.7	2.2	2.4	1.5	2.7	2.6	6.1	2.0
95% CI	1.8–2.3	1.5–1.9	1.9–2.6	2.0–2.9	1.1–1.9	1.8–3.7	1.5–4.1	3.0–10.6	1.9–2.2
Target age 20–69 years									
AS rate (A)	2.7	2.0	3.0	3.0	1.9	3.6	3.8	4.7	2.6
95% CI	2.3–3.0	1.6–2.4	2.5–3.6	2.3–3.8	1.3–2.6	2.3–5.4	2.0–6.4	2.0–9.1	2.4–2.8
AS rate (W)	2.4	1.8	2.8	2.8	1.7	3.3	3.4	4.4	2.4
95% CI	2.1–2.8	1.5–2.2	2.3–3.3	2.1–3.5	1.2–2.4	2.1–5.0	1.8–5.8	1.8–8.4	2.2–2.6

Notes

1. The age-standardised rates were averaged over 4 years to smooth annual variations that may occur in the smaller states and territories.
2. Deaths were derived by year and state of registration.
3. Rates are expressed per 100,000 women and age-standardised to the Australian 2001 population (A) and the WHO World Standard Population (W).

Source: AIHW Mortality Database.

Table 100: Deaths from cervical cancer by age, states and territories, 2001–2004

Age group	NSW	Vic	Qld	WA	SA	Tas	ACT	NT	Australia
0–4	0	0	0	0	0	0	0	0	0
5–9	0	0	0	0	0	0	0	0	0
10–14	0	0	0	0	0	0	0	0	0
15–19	0	0	0	1	0	0	0	0	1
20–24	2	0	0	0	0	0	0	0	2
25–29	1	1	3	3	1	2	0	1	12
30–34	11	4	11	8	1	0	2	0	37
35–39	14	9	9	2	2	3	2	1	42
40–44	21	8	17	3	5	2	0	1	57
45–49	23	15	13	8	12	4	1	1	77
50–54	27	16	12	12	9	5	0	4	85
55–59	30	15	13	6	6	4	1	0	75
60–64	33	18	14	6	7	1	1	0	80
65–69	25	18	12	9	8	1	1	1	75
70–74	31	19	21	10	3	2	0	0	86
75–79	28	29	20	9	9	4	2	0	101
80–84	33	26	18	14	3	3	0	1	98
85+	35	28	14	15	15	4	0	0	111
All ages	314	206	177	106	81	35	10	10	939
Target age 20–69 years	187	104	104	57	51	22	8	9	542

Notes

1. Numbers were averaged over 4 years to smooth annual variations that may occur in the smaller states and territories.
2. Deaths were derived by year and state of registration.

Source: AIHW Mortality Database.

Table 101: Age-specific and age-standardised death rates for cervical cancer by age, states and territories, 2001–2004

Age group	NSW	Vic	Qld	WA	SA	Tas	ACT	NT	Australia
0–4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
5–9	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
10–14	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
15–19	0.0	0.0	0.0	0.4	0.0	0.0	0.0	0.0	0.0
20–24	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1
25–29	0.1	0.1	0.6	1.1	0.5	3.6	0.0	2.9	0.4
30–34	1.1	0.5	1.9	2.7	0.5	0.0	3.8	0.0	1.2
35–39	1.4	1.2	1.6	0.7	0.9	4.4	4.0	3.1	1.4
40–44	2.1	1.1	2.9	1.0	2.1	2.7	0.0	3.3	1.9
45–49	2.5	2.2	2.5	2.8	5.4	5.7	2.0	3.8	2.7
50–54	3.1	2.5	2.4	4.6	4.2	7.6	0.0	17.6	3.2
55–59	4.1	2.8	3.1	2.9	3.3	7.0	2.8	0.0	3.4
60–64	5.7	4.2	4.4	3.7	5.0	2.2	4.2	0.0	4.7
65–69	5.0	4.9	4.7	6.8	6.5	2.6	5.5	17.8	5.2
70–74	6.6	5.6	9.1	8.7	2.6	5.7	0.0	0.0	6.5
75–75	6.7	9.4	9.9	9.2	8.1	12.9	14.7	0.0	8.5
80–84	10.7	11.7	12.2	19.7	3.6	12.8	0.0	57.0	11.3
85+	12.9	13.9	10.9	22.7	20.2	19.4	0.0	0.0	14.4
All ages									
AS rate (A)	2.1	1.9	2.3	2.7	2.2	3.3	1.6	3.7	2.2
95% CI	1.9–2.4	1.6–2.1	2.0–2.7	2.2–3.2	1.8–2.8	2.3–4.6	0.8–3.0	1.5–7.4	2.0–2.3
AS rate (W)	1.6	1.4	1.8	2.0	1.7	2.6	1.4	2.9	1.7
95% CI	1.4–1.8	1.2–1.6	1.5–2.0	1.6–2.4	1.3–2.2	1.7–3.6	0.7–2.5	1.3–5.5	1.5–1.8
Target age 20–69 years									
AS rate (A)	2.2	1.6	2.2	2.3	2.5	3.6	2.0	4.4	2.1
95% CI	1.9–2.5	1.3–2.0	1.8–2.6	1.8–3.0	1.9–3.3	2.2–5.4	0.8–3.9	1.8–8.6	1.9–2.3
AS rate (W)	2.0	1.5	2.0	2.2	2.3	3.3	1.9	4.0	2.0
95% CI	1.7–2.3	1.2–1.8	1.7–2.5	1.6–2.8	1.7–3.0	2.1–5.1	0.8–3.8	1.7–7.7	1.8–2.1

Notes

1. The age-standardised rates were averaged over 4 years to smooth annual variations that may occur in the smaller states and territories.
2. Deaths were derived by year and state of registration.
3. Rates are expressed per 100,000 women and age-standardised to the Australian 2001 population (A) and the WHO World Standard Population (W).

Source: AIHW Mortality Database.

Indicator 6.2: Mortality by location

Table 102: Deaths from cervical cancer by age and location, 1997–2000 and 2001–2004

Age group	Major cities		Regional		Remote		Australia	
	1997–2000	2001–2004	1997–2000	2001–2004	1997–2000	2001–2004	1997–2000	2001–2004
0–4	0	0	0	0	0	0	0	0
5–9	0	0	0	0	0	0	0	0
10–14	0	0	0	0	0	0	0	0
15–19	0	1	0	0	0	0	0	1
20–24	3	0	2	2	0	0	5	2
25–29	7	6	6	5	0	1	14	12
30–34	19	26	6	9	3	2	29	37
35–39	32	25	22	16	1	1	56	42
40–44	43	30	21	23	4	4	67	57
45–49	71	46	20	30	3	0	96	77
50–54	51	53	26	29	1	2	79	85
55–59	44	47	24	28	4	0	72	75
60–64	53	44	34	35	3	1	89	80
65–69	64	46	28	27	4	2	96	75
70–74	78	57	48	27	3	2	131	86
75–79	71	61	35	35	3	3	109	101
80–84	65	63	25	34	4	1	95	98
85+	71	83	36	25	1	3	108	111
All ages	672	589	332	324	35	23	1,046	939
Target age								
20–69 years	387	324	189	203	23	13	603	542

Notes

1. Deaths were derived from place of usual residence and by year of registration.
2. The number of deaths is presented as 4-year rolling blocks of data.
3. In 1997–2000 and 2001–2004 there were 6 and 3 deaths respectively that were excluded from these data because the respective postcodes were not able to be matched to the coding used for this analysis or postcodes were not provided.
4. Totals may not add up due to rounding.

Source: AIHW Mortality Database.

Table 103: Age-specific and age-standardised death rates for cervical cancer by age and location, 1997–2000 and 2001–2004

Age group	Major cities		Regional		Remote		Australia	
	1997–2000	2001–2004	1997–2000	2001–2004	1997–2000	2001–2004	1997–2000	2001–2004
0–4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
5–9	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
10–14	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
15–19	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0
20–24	0.2	0.0	0.3	0.3	0.0	0.0	0.2	0.1
25–29	0.3	0.3	0.7	0.7	0.5	1.2	0.5	0.4
30–34	1.0	1.3	0.8	1.1	4.1	2.1	1.0	1.2
35–39	1.6	1.3	2.3	1.8	1.3	1.5	1.9	1.4
40–44	2.3	1.5	2.3	2.4	5.1	6.0	2.4	1.9
45–49	4.0	2.5	2.5	3.4	5.2	0.7	3.6	2.7
50–54	3.3	3.0	3.7	3.5	2.6	4.0	3.4	3.2
55–59	3.8	3.3	4.0	3.9	11.9	0.0	4.0	3.4
60–64	5.4	4.0	6.6	5.9	9.0	1.9	5.9	4.7
65–69	7.1	5.0	6.1	5.4	15.7	8.2	6.9	5.2
70–74	8.9	6.6	11.1	6.1	18.3	9.0	9.9	6.5
75–79	9.6	7.8	10.2	9.3	23.4	21.4	10.0	8.5
80–84	13.2	10.8	10.6	12.3	46.2	14.5	12.9	11.3
85+	16.0	15.9	18.0	10.4	18.5	32.2	16.7	14.4
All ages								
AS rate (A)	2.6	2.1	2.7	2.4	5.2	3.1	2.7	2.2
95% CI	2.4–2.8	1.9–2.2	2.4–3.0	2.1–2.7	3.5–7.2	1.9–4.6	2.5–2.8	2.0–2.3
AS rate (W)	1.9	1.5	2.0	1.9	3.9	2.2	2.0	1.7
95% CI	1.8–2.1	1.4–1.7	1.8–2.3	1.6–2.1	2.6–5.3	1.4–3.3	1.9–2.2	1.5–1.8
Target age 20–69 years								
AS rate (A)	2.5	1.9	2.5	2.5	4.6	2.4	2.6	2.1
95% CI	2.2–2.8	1.7–2.1	2.2–2.9	2.2–2.9	2.9–6.9	1.2–4.0	2.4–2.8	1.9–2.3
AS rate (W)	2.3	1.8	2.4	2.3	4.3	2.2	2.4	2.0
95% CI	2.1–2.5	1.6–2.0	2.0–2.7	2.0–2.7	2.7–6.4	1.1–3.7	2.2–2.6	1.8–2.1

Notes

1. The age-standardised rates are presented as 4-year rolling blocks of data.
2. Deaths were derived from place of usual residence and by year of registration.
3. Rates are expressed per 100,000 women and age-standardised to the Australian 2001 population (A) and the WHO World Standard Population (W).

Source: AIHW Mortality Database.

Indicator 6.3: Mortality by Indigenous status

Table 104: Number of deaths and age-specific and age-standardised death rates for cervical cancer by age and Indigenous status, Queensland, South Australia, Western Australia and Northern Territory, 2001–2004

Age group	Indigenous Australians		Other Australians	
	Number	Rate	Number	Rate
0–4	0	0.0	0	0.0
5–9	0	0.0	0	0.0
10–14	0	0.0	0	0.0
15–19	0	0.0	1	0.0
20–24	0	0.0	0	0.1
25–29	0	0.0	8	0.0
30–34	3	6.4	17	0.8
35–39	1	2.5	13	1.6
40–44	7	21.0	19	1.2
45–49	3	11.6	31	1.7
50–54	3	14.8	34	3.0
55–59	1	7.3	24	3.5
60–64	2	19.3	25	3.0
65–69	2	28.4	28	4.0
70–74	2	41.7	32	5.5
75+	3	48.9	115	6.9
All ages	27		347	
AS rate (A)		10.5		2.3
95% CI		6.3–16.0		2.0–2.5
AS rate (W)		7.9		1.7
95% CI		5.0–11.7		1.5–1.9
Target age 20–69 years	22		199	
AS rate (A)		9.9		2.1
95% CI		6.0–15.3		1.9–2.5
AS rate (W)		9.1		2.0
95% CI		5.5–13.9		1.7–2.3

Notes

1. Deaths were derived by state and year of registration.
2. The number of deaths is presented as a 4-year rolling block of data.
3. Only Queensland, South Australia, Western Australia and the Northern Territory have Indigenous death registration data considered to be of a publishable standard.

Source: AIHW Mortality Database.