

Tables

Indicator 1: Participation rate for cervical cancer screening

Table A1a: Number of women participating in the National Cervical Screening Program by age, State or Territory, 1996–1997

Age group	NSW ^{(a)(b)}	Vic	WA	SA ^(b)	Tas	ACT ^(c)	NT ^(d)	Australia
20–24	100,579	84,743	38,551	28,157	10,458	7,354	4,958	274,800
25–29	142,392	123,651	48,584	36,478	11,786	8,716	6,184	377,791
30–34	148,690	127,168	50,173	37,889	12,293	8,571	5,511	390,296
35–39	144,884	125,992	48,770	37,991	12,242	8,217	4,722	382,818
40–44	121,267	107,973	42,240	32,341	10,358	7,409	3,802	325,390
45–49	102,807	93,757	34,555	28,267	8,757	6,841	3,094	278,078
50–54	77,168	72,536	23,546	20,925	6,201	4,621	1,827	206,824
55–59	52,031	50,217	16,292	14,740	4,441	2,744	1,028	141,493
60–64	37,832	37,706	11,973	11,506	3,336	1,830	518	104,701
65–69	28,365	30,597	8,693	9,298	2,498	1,182	284	80,917
70–74	19,105	14,576	3,416	7,022	929	477	116	45,640
75–79	n.a.	5,292	1,153	n.a.	332	155	49	6,981
80–84	n.a.	1,869	414	n.a.	122	51	22	2,478
85+	n.a.	997	15	n.a.	39	9	6	1,066
Not stated	9,750	n.a.	n.a.	1,029	10	77	96	10,962
Total	984,870	877,074	328,375	265,643	83,802	58,254	32,217	2,630,235
20–69 years	956,015	854,340	323,377	257,592	82,370	57,485	31,929	2,563,108

(a) The New South Wales Pap Test register commenced in July 1996, therefore data has been estimated for the period January to July 1996.

(b) New South Wales and South Australia have grouped all women aged 70 years or more, and for the purposes of this table they appear in the 70–74 age group.

(c) The ACT register only registers women with an ACT address.

(d) The Northern Territory Pap Smear register commenced in March 1996, therefore data has been estimated for the period January to March 1996.

Note: The Queensland Health Pap Smear Register began operations in February 1999; therefore no data are available for this report.

Table A1b: Number of women participating in the National Cervical Screening Program by age, State or Territory, 1997–1998

Age group	NSW	Vic	WA	SA ^(a)	Tas	ACT ^(b)	NT	Australia
20–24	106,771	82,031	36,422	27,668	9,989	7,042	4,880	274,803
25–29	152,260	122,571	47,752	36,982	11,899	8,772	6,228	386,464
30–34	154,446	126,446	48,703	38,060	11,989	8,552	5,441	393,637
35–39	154,060	127,261	49,431	39,175	12,698	8,407	4,846	395,878
40–44	130,315	110,367	42,817	34,069	10,682	7,587	3,799	339,636
45–49	109,283	95,905	35,005	29,617	9,051	7,018	3,091	288,970
50–54	84,773	76,368	25,537	23,309	6,896	5,257	1,898	224,038
55–59	56,631	52,128	16,903	15,820	4,701	3,101	1,069	150,353
60–64	41,976	39,569	12,735	12,639	3,582	2,001	535	113,037
65–69	31,116	31,903	9,323	9,972	2,706	1,363	305	86,688
70–74	13,371	15,051	3,590	7,547	864	535	117	41,075
75–79	5,154	5,357	1,274	n.a.	319	195	52	12,351
80–84	1,562	1,721	389	n.a.	106	51	15	3,844
85+	564	660	38	n.a.	35	15	8	1,320
Not stated	9,080	n.a.	n.a.	389	4	40	43	9,556
Total	1,051,362	887,338	329,919	275,247	85,521	59,936	32,327	2,721,650
20–69 years	1,021,631	864,549	324,628	267,311	84,193	59,100	32,092	2,653,504

(a) South Australia has grouped all women aged 70 years or more, and for the purposes of this table they appear in the 70–74 age group.

(b) The ACT register only registers women with an ACT address

Note: The Queensland Health Pap Smear Register began operations in February 1999; therefore no data are available for this report.

Table A2a: Participation rates in the National Cervical Screening Program by age, State or Territory, 1996–1997

Age group	NSW ^(a)	Vic	WA	SA ^(a)	Tas	ACT ^(b)	NT	Australia
20–24	44.6	49.8	58.1	55.5	66.3	52.1	60.1	49.9
25–29	59.8	67.8	70.9	68.7	71.7	66.0	67.1	65.0
30–34	62.4	71.5	73.0	69.5	71.4	68.3	67.1	67.6
35–39	63.4	75.0	73.4	70.8	69.7	68.6	67.0	69.2
40–44	62.0	74.1	71.8	68.8	68.6	69.2	66.6	68.0
45–49	60.9	74.2	70.0	67.8	67.5	70.3	69.8	67.3
50–54	64.1	82.0	71.8	71.1	67.6	75.7	67.2	71.5
55–59	53.2	69.8	62.1	62.0	58.0	66.1	62.5	60.7
60–64	44.2	59.7	55.0	54.5	50.3	60.1	48.4	51.7
65–69	32.8	48.8	42.1	42.5	38.3	43.9	37.4	40.1
70–74	24.4	25.5	19.4	33.9	15.6	19.9	22.7	25.0
75–79	n.a.	12.3	8.7	n.a.	7.1	9.2	14.9	5.0
80–84	n.a.	6.6	4.6	n.a.	3.9	5.0	12.3	2.7
85+ ^(c)	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.
Total	50.9	61.1	61.7	56.9	57.8	62.5	67.3	56.3
20–69 years	56.7	68.0	67.4	64.9	65.9	65.1	65.1	62.4

(a) New South Wales and South Australia have grouped all women aged 70 years or more, and for the purposes of this table they appear in the 70–74 age group.

(b) The ACT register only registers women with an ACT address.

(c) Rates cannot be calculated for women in the 85+ age group because hysterectomy fractions are not available for this age group.

Note: The Queensland Health Pap Smear Register began operations in February 1999; therefore no data are available for this report.

Table A2b: Participation rates in the National Cervical Screening Program by age, State or Territory, 1997–1998

Age group	NSW	Vic	WA	SA ^(a)	Tas	ACT ^(b)	NT	Australia
20–24	48.2	48.8	54.5	55.8	65.3	52.5	60.6	50.6
25–29	62.6	66.4	68.1	69.4	72.4	66.5	66.0	65.5
30–34	65.7	71.5	71.3	71.6	72.3	70.2	65.8	69.0
35–39	66.6	74.9	73.2	72.5	72.7	70.8	67.1	70.7
40–44	65.4	74.6	71.4	72.0	70.2	71.7	64.3	69.8
45–49	64.3	75.6	69.6	71.0	69.8	72.8	68.1	69.4
50–54	66.1	80.7	72.0	74.0	70.9	79.4	64.7	72.5
55–59	56.6	70.8	62.3	65.0	60.4	71.3	60.7	62.9
60–64	48.3	61.6	57.0	59.0	53.4	63.2	46.8	54.9
65–69	36.4	51.3	45.1	46.6	41.5	50.2	39.4	43.4
70–74	17.0	26.2	20.2	12.8	14.7	22.1	22.0	22.4
75–79	8.2	11.9	9.1	n.a.	6.6	10.6	15.5	8.5
80–84	4.0	6.0	4.3	n.a.	3.3	4.9	7.8	4.2
85+ ^(c)	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.
Total	53.7	61.0	60.7	60.8	58.8	64.2	66.0	57.7
20–69 years	60.1	68.1	66.4	67.2	67.5	67.4	64.1	63.9

(a) South Australia have grouped all women aged 70 years or more, and for the purposes of this table they appear in the 70–74 age group.

(b) The ACT register only registers women with an ACT address.

(c) Rates cannot be calculated for women in the 85+ age group because hysterectomy fractions are not available for this age group.

Note: The Queensland Health Pap Smear Register began operations in February 1999; therefore no data are available for this report.

Indicator 2: Early rescreening

Table A3: Number of women with repeat screenings in the 24 months following a negative Pap smear in February 1997 by State and Territory, and Australia 1996-1997 and 1997-1998

No. of tests	NSW	Vic	WA	SA	Tas	ACT ^(a)	NT	Australia 1997-1998	Australia ^(b) 1996-1997
Number of women									
0	29,603	24,855	9,402	7,980	2,520	1,358	842	76,560	44,125
1	20,827	18,803	6,546	4,556	1,628	686	410	53,456	31,407
2	4,050	4,225	1,197	961	311	86	92	10,922	7,295
3	657	887	202	241	57	8	28	2,080	1,567
4	136	266	33	53	11	2	7	508	470
5 or more	30	136	6	19	2	0	3	196	195

(a) The ACT register only registers women with an ACT address.

(b) The NSW Pap Test Register began operations on 29 August 1996, therefore no data were available for this indicator for the period 1996-1997.

Note: The Queensland Health Pap Smear Register began operations in February 1999; therefore no data are available for this report.

Table A4: Percentage of women with repeat screenings in the 24 months following a negative Pap smear in February 1997 by State and Territory, and Australia 1996-1997 and 1997-1998

No. of tests	NSW	Vic	WA	SA	Tas	ACT ^(a)	NT	Australia 1997-1998	Australia ^(b) 1996-1997
Per cent of women									
0	53.5	50.5	54.1	57.8	55.6	63.5	60.9	53.3	51.9
1	37.7	38.2	37.7	33.0	36.0	32.1	29.7	37.2	36.9
2	7.3	8.6	6.9	7.0	6.9	4.0	6.7	7.6	8.6
3	1.2	1.8	1.2	1.7	1.3	0.4	2.0	1.4	1.9
4	0.2	0.5	0.2	0.4	0.2	0.1	0.5	0.4	0.6
5 or more	0.1	0.3	0.0	0.1	0.0	0.0	0.2	0.1	0.2

(a) The ACT register only registers women with an ACT address.

(b) The NSW Pap Test Register began operations on 29 August 1996, therefore no data were available for this indicator for the period 1996-1997.

Note: The Queensland Health Pap Smear Register began operations in February 1999; therefore no data are available for this report.

Indicator 3: Low-grade abnormality detection

Table A5a: Number of low- and high-grade abnormalities on histology for women aged 20–69 years, by State and Territory, 1997

Abnormalities	NSW	Vic	WA	SA	Tas	NT	Australia
Low-grade	6,447	3,419	2,209	2,370	543	326	15,314
High-grade	3,601	3,388	1,432	1,310	430	231	10,392
Ratio	1.79	1.01	1.54	1.81	1.26	1.41	1.47

Notes: (1) The Queensland Health Pap Smear Register began operations in February 1999; therefore no data are available for this report.

(2) The ACT did not collect histology details during this period; therefore no data are available for this indicator.

Table A5b: Number of low- and high-grade abnormalities on histology for women aged 20–69 years, by State and Territory, 1998

Abnormalities	NSW	Vic	WA	SA	Tas	NT	Australia
Low-grade	5,799	3,329	2,090	2,179	756	258	14,411
High-grade	3,960	2,994	1,414	1,505	534	298	10,701
Ratio	1.46	1.11	1.48	1.45	1.42	0.87	1.35

Notes: (1) The Queensland Health Pap Smear Register began operations in February 1999; therefore no data are available for this report.

(2) The ACT did not collect histology details during this period; therefore no data are available for this indicator.

Indicator 4: High-grade abnormality detection

Table A6a: Number of histologically confirmed high-grade abnormalities by age, by State and Territory, 1997

Age group	NSW	Vic	WA	SA ^(a)	Tas	NT	Australia
20–24	788	670	320	184	115	46	2,123
25–29	1,022	1,015	370	335	109	62	2,913
30–34	753	696	286	265	74	41	2,115
35–39	453	429	193	216	61	32	1,384
40–44	267	277	110	90	32	19	795
45–49	154	139	82	89	17	15	496
50–54	64	80	28	50	9	9	240
55–59	34	35	16	27	5	5	122
60–64	31	23	15	33	2	2	106
65–69	35	24	12	21	6	0	98
70–74	20	16	1	35	n.a.	n.a.	72
75–79	7	8	2	n.a.	n.a.	n.a.	17
80–84	2	3	4	n.a.	n.a.	n.a.	9
85+	3	n.a.	n.a.	n.a.	n.a.	n.a.	3
Age not stated	5	n.a.	n.a.	1	n.a.	n.a.	6
Total	3,638	3,415	1,439	1,346	430	231	10,499
20–69 years	3,601	3,388	1,432	1,310	430	231	10,392

(a) South Australia has grouped all women aged 70 years or more, and for the purposes of this table they appear in the 70–74 age group.

Note: The Queensland Health Pap Smear Register began operations in February 1999; therefore no data are available for this report.

Table A6b: Number of histologically confirmed high-grade abnormalities by age, by State and Territory, 1998

Age group	NSW	Vic	WA	SA ^(a)	Tas	NT	Australia
20–24	847	566	309	305	127	66	2,220
25–29	1,180	895	408	411	137	95	3,126
30–34	748	591	276	259	109	58	2,041
35–39	531	437	184	192	85	39	1,468
40–44	284	233	124	134	39	19	833
45–49	163	116	52	87	21	8	447
50–54	101	65	25	51	7	8	257
55–59	51	38	19	29	4	4	145
60–64	37	38	11	23	5	1	115
65–69	17	15	6	14	0	0	52
70–74	19	8	6	29	0	0	62
75–79	7	7	7	n.a.	0	0	21
80–84	6	3	1	n.a.	1	0	11
85+	2	2	0	n.a.	0	0	4
Age not stated	3	n.a.	n.a.	1	0	0	4
Total	3,996	3,014	1,428	1,535	535	298	10,806
20–69 years	3,960	2,994	1,414	1,505	534	298	10,704

(a) South Australia has grouped all women aged 70 years or more, and for the purposes of this table they appear in the 70–74 age group.

Note: The Queensland Health Pap Smear Register began operations in February 1999; therefore no data are available for this report.

Table A7a: Number of women screened by age, by State and Territory, 1997

Age group	NSW	Vic	WA	SA ^(a)	Tas	NT	Australia
20–24	58,842	45,403	21,222	15,231	5,705	2,800	149,203
25–29	85,421	70,504	27,798	20,617	7,121	3,497	214,958
30–34	89,016	72,728	28,423	21,266	7,173	3,055	221,661
35–39	87,766	72,287	28,506	21,413	7,377	2,612	219,961
40–44	73,682	62,536	24,720	18,245	6,277	2,073	187,533
45–49	62,596	54,917	20,078	16,097	5,352	1,748	160,788
50–54	48,045	43,848	14,270	12,215	3,997	1,052	123,427
55–59	32,031	29,632	9,554	8,494	2,706	579	82,996
60–64	23,277	21,639	6,902	6,681	2,057	285	60,841
65–69	17,239	16,715	4,897	5,211	1,551	168	45,781
70–74	11,439	7,989	n.a.	3,887	488	59	23,862
75–79	n.a.	2,957	n.a.	n.a.	161	29	3,147
80–84	n.a.	996	n.a.	n.a.	64	9	1,069
85+	n.a.	n.a.	n.a.	n.a.	19	3	22
Age not stated	0	0	0	333	4	22	359
Total	589,354	502,151	186,370	149,690	50,052	17,991	1,495,608
20–69 years	577,915	490,209	186,370	145,470	49,316	17,869	1,467,149

(a) South Australia has grouped all women aged 70 years or more, and for the purposes of this table they appear in the 70–74 age group.

Note: The Queensland Health Pap Smear Register began operations in February 1999; therefore no data are available for this report.

Table A7b: Number of women screened by age, by State and Territory, 1998

Age group	NSW	Vic	WA	SA ^(a)	Tas	NT	Australia
20–24	60,919	47,286	22,267	16,122	5,532	2,678	1,54,804
25–29	90,720	73,379	29,342	21,785	6,883	3,584	2,25,693
30–34	92,053	76,937	29,693	22,260	6,911	3,170	2,31,024
35–39	93,293	77,383	30,288	23,111	7,387	2,896	2,34,358
40–44	79,402	67,374	26,259	20,351	6,177	2,249	2,01,812
45–49	66,812	58,146	21,676	17,464	5,207	1,783	1,71,088
50–54	52,675	46,514	15,753	13,857	4,073	1,092	1,33,964
55–59	34,574	31,584	10,123	9,087	2,723	615	88,706
60–64	25,243	23,974	7,609	7,138	2,000	308	66,272
65–69	17,970	19,257	5,453	5,526	1,476	153	49,835
70–74	7,685	9,235	n.a.	4,161	503	73	21,657
75–79	2,841	3,170	n.a.	n.a.	183	32	6,226
80–84	805	979	n.a.	n.a.	51	8	1,843
85+	293	518	n.a.	n.a.	16	6	833
Age not stated	4,335	n.a.	n.a.	132	3	22	4,492
Total	629,620	535,736	198,463	160,994	49,125	18,669	1,592,607
20–69 years	613,661	521,834	198,463	156,701	48,369	18,528	1,557,556

(a) South Australia has grouped all women aged 70 years or more, and for the purposes of this table they appear in the 70–74 age group.

Note: The Queensland Health Pap Smear Register began operations in February 1999; therefore no data are available for this report.

Table A8a: Rate of histologically confirmed high-grade abnormalities per 1,000 women screened, by State and Territory, 1997

Age group	NSW	Vic	WA	SA ^(a)	Tas	NT	Australia
20–24	13.4	14.8	15.1	12.1	20.2	16.4	14.2
25–29	12.0	14.4	13.3	16.3	15.3	17.7	13.6
30–34	8.5	9.6	10.1	12.5	10.3	13.4	9.5
35–39	5.2	5.9	6.8	10.1	8.3	12.3	6.3
40–44	3.6	4.4	4.5	4.9	5.1	9.2	4.2
45–49	2.5	2.5	4.1	5.5	3.2	8.6	3.1
50–54	1.3	1.8	1.9	4.1	2.3	8.6	1.9
55–59	1.1	1.2	1.7	3.2	1.8	8.6	1.5
60–64	1.3	1.1	2.2	4.9	1.0	7.0	1.7
65–69	2.0	1.4	2.5	4.0	3.9	0.0	2.1
70–74	1.8	2.0	n.a.	9.0	n.a.	0.0	3.0
75–79	n.a.	2.7	n.a.	n.a.	n.a.	0.0	5.4
80–84	n.a.	3.0	n.a.	n.a.	n.a.	0.0	8.4
85+	n.a.	n.a.	n.a.	n.a.	n.a.	0.0	136.4
Total	6.2	6.8	7.7	9.0	8.6	12.8	7.0
20–69 years	6.2	6.9	7.7	9.0	8.7	12.9	7.1

(a) South Australia has grouped all women aged 70 years or more, and for the purposes of this table they appear in the 70–74 age group.

Notes: (1) The Queensland Health Pap Smear Register began operations in February 1999; therefore no data are available for this report.

(2) The ACT did not collect histology details during this period; therefore no data are available for this indicator.

Table A8b: Rate of histologically confirmed high-grade abnormalities per 1,000 women screened, by State and Territory, 1998

Age group	NSW	Vic	WA	SA ^(a)	Tas	NT	Australia
20–24	13.9	11.9	13.9	18.9	22.9	24.6	14.3
25–29	13.0	12.2	13.9	18.9	19.9	26.5	13.9
30–34	8.1	7.7	9.3	11.6	15.8	18.3	8.8
35–39	5.7	5.6	6.1	8.3	11.5	13.5	6.3
40–44	3.6	3.5	4.7	6.6	6.3	8.4	4.1
45–49	2.4	1.9	2.4	4.9	4.0	4.5	2.6
50–54	1.9	1.4	1.6	3.7	1.7	7.3	1.9
55–59	1.5	1.2	1.9	3.2	1.5	6.5	1.6
60–64	1.5	1.6	1.5	3.2	2.5	3.2	1.7
65–69	0.9	0.8	1.1	2.5	0.0	0.0	1.0
70–74	2.5	0.8	n.a.	6.9	0.0	0.0	2.9
75–79	2.5	2.2	n.a.	n.a.	0.0	0.0	3.4
80–84	7.5	3.1	n.a.	n.a.	19.6	0.0	6.0
85+	6.8	3.9	n.a.	n.a.	0.0	0.0	4.8
Total	6.4	5.6	7.2	9.5	10.9	16.0	6.8
20–69 years	6.4	5.7	7.1	9.6	11.0	16.1	6.9

(a) South Australia has grouped all women aged 70 years or more, and for the purposes of this table they appear in the 70–74 age group.

Notes: (1) The Queensland Health Pap Smear Register began operations in February 1999; therefore no data are available for this report.

(2) The ACT did not collect histology details during this period; therefore no data are available for this indicator.

Table A9a: Age-standardised high-grade abnormality rate on histology per 1,000 women screened aged 20-69 years, by State and Territory, 1997

	NSW	Vic	WA	SA	Tas	NT	Australia
Standardised rate	6.1	6.9	7.3	8.8	8.5	11.3	6.9
95% CI	5.9–6.3	6.7–7.1	6.9–7.7	8.3–9.2	7.6–9.2	9.8–12.7	6.8–7.0

Notes: (1) The Queensland Health Pap Smear Register began operations in February 1999; therefore no data are available for this report.

(2) The ACT did not collect histology details during this period; therefore no data are available for this indicator.

(3) Standardised to the 1991 Australian total population.

Table A9b: Age-standardised high-grade abnormality rate on histology per 1,000 women screened aged 20-69 years, by State and Territory, 1998

	NSW	Vic	WA	SA	Tas	NT	Australia
Standardised rate	6.3	5.8	6.7	9.5	10.5	13.3	6.7
95% CI	6.1–6.5	5.6–6.0	6.4–7.1	9.1–10.0	9.6–11.4	11.8–14.8	6.6–6.9

Notes: (1) The Queensland Health Pap Smear Register began operations in February 1999; therefore no data are available for this report.

(2) The ACT did not collect histology details during this period; therefore no data are available for this indicator.

(3) Standardised to the 1991 Australian total population.

Indicator 5: Incidence of micro-invasive cervical cancer

Table A10: New cases of micro-invasive cervical cancer by age, Australia, 1985–1996

Age group	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996
0–4	0	0	0	0	0	0	0	0	0	0	0	0
5–9	0	0	0	0	0	0	0	0	0	0	0	0
10–14	0	0	0	0	0	0	0	0	0	0	0	0
15–19	1	0	0	0	0	0	1	0	0	0	0	0
20–24	8	1	3	4	1	4	0	5	1	7	1	6
25–29	11	9	8	13	12	15	14	13	7	17	16	17
30–34	23	17	18	20	27	30	31	32	31	32	42	18
35–39	19	17	25	12	10	24	38	23	26	31	29	36
40–44	13	6	14	11	17	22	32	24	16	26	29	23
45–49	11	7	10	9	5	17	10	12	15	27	22	11
50–54	5	4	3	6	4	4	11	12	17	9	12	10
55–59	2	3	4	5	5	8	6	12	5	5	8	7
60–64	8	1	3	2	6	8	7	7	6	10	9	6
65–69	3	2	1	2	2	6	7	9	10	6	7	10
70–74	2	3	0	0	0	2	4	2	4	6	5	3
75–79	0	1	1	1	1	3	3	2	1	3	5	2
80–84	0	0	0	0	0	0	2	0	0	0	1	1
85+	0	0	0	0	0	0	0	0	1	2	1	1
Total	106	71	90	85	90	143	166	153	140	181	187	151

Source: National Cancer Statistics Clearing House (AIHW).

Table A11: Age-specific and age-standardised rates of micro-invasive cervical cancer by age, Australia, 1985–1996

Age group	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996
0–4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
5–9	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
10–14	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
15–19	0.2	0.0	0.0	0.0	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.0
20–24	1.2	0.2	0.5	0.6	0.2	0.6	0.0	0.7	0.1	1.0	0.1	0.9
25–29	1.7	1.3	1.2	1.9	1.7	2.1	2.0	1.9	1.0	2.5	2.3	2.4
30–34	3.7	2.7	2.8	3.0	4.0	4.3	4.4	4.4	4.2	4.4	5.7	2.5
35–39	3.2	2.7	4.0	1.9	1.5	3.7	5.7	3.4	3.8	4.4	4.1	4.9
40–44	2.8	1.2	2.6	1.9	2.9	3.6	5.0	3.7	2.5	4.0	4.3	3.4
45–49	2.8	1.7	2.4	2.1	1.1	3.6	2.0	2.2	2.6	4.5	3.6	1.7
50–54	1.4	1.1	0.8	1.6	1.0	1.0	2.7	2.8	3.9	2.0	2.5	2.0
55–59	0.5	0.8	1.1	1.4	1.4	2.2	1.7	3.3	1.3	1.3	2.0	1.7
60–64	2.2	0.3	0.8	0.5	1.6	2.2	1.9	1.9	1.7	2.8	2.5	1.7
65–69	1.0	0.7	0.3	0.6	0.6	1.7	2.0	2.5	2.8	1.7	2.0	2.8
70–74	0.8	1.1	0.0	0.0	0.0	0.7	1.4	0.7	1.3	1.9	1.5	0.9
75–79	0.0	0.5	0.5	0.5	0.5	1.4	1.3	0.9	0.4	1.3	2.1	0.8
80–84	0.0	0.0	0.0	0.0	0.0	0.0	1.4	0.0	0.0	0.0	0.6	0.6
85+	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.8	1.6	0.7	0.7
All ages												
AS Rate (A)	1.4	0.9	1.2	1.1	1.1	1.7	1.9	1.7	1.6	2.0	2.0	1.6
AS Rate (W)	1.2	0.8	1.0	0.9	0.9	1.4	1.6	1.5	1.3	1.7	1.7	1.4
Ages 20–69												
AS Rate (A)	2.2	1.4	1.8	1.7	1.7	2.6	2.9	2.7	2.4	3.0	3.1	2.5
AS Rate (W)	2.1	1.3	1.7	1.6	1.6	2.5	2.7	2.6	2.3	2.9	2.9	2.4

Note: Rates are expressed per 100,000 women and age-standardised to the Australian 1991 population (A) and the World Standard Population (W).

Source: National Cancer Statistics Clearing House (AIHW).

Indicator 6: Incidence of invasive squamous, adenocarcinoma, adeno-squamous and other cervical cancer

Table A12: New cases of cervical cancer by age, Australia, 1985–1996

Age group	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996
0–4	0	0	0	1	0	0	0	0	0	0	0	0
5–9	0	0	0	0	0	0	0	0	0	1	0	0
10–14	1	0	0	0	0	0	0	0	0	0	0	0
15–19	1	3	3	4	1	1	1	0	1	1	2	1
20–24	23	12	16	18	17	13	11	9	10	16	5	14
25–29	70	61	66	76	66	61	48	51	36	47	49	43
30–34	109	120	137	130	128	110	119	110	104	117	113	70
35–39	131	125	149	136	121	154	140	128	127	131	117	137
40–44	108	114	113	125	128	136	153	130	130	130	115	118
45–49	83	114	95	92	90	122	102	100	100	133	100	102
50–54	87	75	65	63	81	66	91	77	90	87	58	78
55–59	83	78	60	65	85	82	57	78	80	73	66	62
60–64	97	83	106	91	83	77	84	77	74	89	69	63
65–69	82	81	88	104	97	74	90	88	93	93	77	64
70–74	69	64	78	56	66	67	79	70	65	79	73	57
75–79	43	42	56	50	49	49	48	52	49	64	51	48
80–84	26	27	31	34	26	29	40	34	36	39	30	41
85+	20	20	25	22	20	24	35	21	22	25	34	25
Total	1,033	1,019	1,088	1,067	1,058	1,065	1,098	1,025	1,017	1,125	959	923

Note: The above table includes the incidence of micro-invasive and invasive cervical cancers.

Source: National Cancer Statistics Clearing House (AIHW).

Table A13: Age-specific and age-standardised incidence rates of cervical cancer by age, Australia, 1985–1996

Age group	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996
0–4	0.0	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
5–9	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.0	0.0
10–14	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
15–19	0.2	0.5	0.4	0.6	0.1	0.1	0.2	0.0	0.2	0.2	0.3	0.2
20–24	3.5	1.8	2.5	2.8	2.6	1.9	1.6	1.3	1.4	2.3	0.7	2.0
25–29	10.7	9.1	9.7	10.9	9.3	8.6	6.9	7.4	5.3	6.9	7.1	6.1
30–34	17.4	18.9	21.2	19.7	18.9	15.8	16.7	15.2	14.2	15.9	15.5	9.7
35–39	21.7	20.0	23.9	21.4	18.7	23.5	21.1	18.9	18.5	18.8	16.4	18.8
40–44	22.9	23.1	21.1	21.9	21.5	22.0	23.9	20.3	20.1	19.8	17.2	17.4
45–49	20.8	27.9	22.5	21.1	19.7	25.5	20.3	18.6	17.5	22.3	16.2	15.9
50–54	24.3	20.8	17.7	16.7	20.8	16.5	22.0	18.2	20.7	19.2	12.2	15.7
55–59	22.2	21.0	16.3	17.9	23.5	22.8	15.9	21.3	21.3	18.9	16.7	15.2
60–64	26.7	22.6	28.8	24.6	22.4	20.8	22.7	21.1	20.6	24.9	19.3	17.7
65–69	28.0	26.6	27.8	31.6	28.3	21.2	25.6	24.9	26.2	26.2	21.7	18.0
70–74	26.6	24.3	29.2	20.9	24.8	24.8	28.0	23.9	21.4	24.9	22.6	17.4
75–79	23.4	21.9	28.2	24.3	22.8	22.2	21.3	22.7	21.3	28.1	21.9	19.7
80–84	22.5	22.7	25.0	26.3	19.4	20.8	27.5	22.5	22.7	23.3	17.4	23.2
85+	22.5	21.1	25.7	22.0	19.4	22.7	31.8	18.2	18.1	19.6	25.3	17.7
All ages												
AS Rate (A)	13.4	13.0	13.4	12.9	12.5	12.3	12.4	11.4	11.1	12.0	10.0	9.4
AS Rate (W)	11.1	10.8	11.0	10.6	10.3	10.2	10.1	9.3	9.1	9.9	8.2	7.8
Ages 20–69												
AS Rate (A)	18.4	18.0	18.1	17.7	17.2	16.9	16.6	15.4	15.1	16.1	13.4	12.8
AS Rate (W)	18.1	17.6	17.5	17.2	16.8	16.5	16.1	15.0	14.7	15.8	12.9	12.5

Note: Rates are expressed per 100,000 women and age standardised to the Australian 1991 population (A) and the World Standard Population (W).
Source: National Cancer Statistics Clearing House (AIHW).

Table A14: Number of new cases of cervical cancer by age, by State and Territory, 1993–1996

Age group	NSW	Vic	Qld	WA	SA	Tas	ACT	NT	Australia
0–4	0	0	0	0	0	0	0	0	0
5–9	0	0	0	0	1	0	0	0	1
10–14	0	0	0	0	0	0	0	0	0
15–19	3	0	2	0	0	0	0	0	5
20–24	11	15	13	5	0	1	0	0	45
25–29	47	40	38	22	15	6	5	2	175
30–34	135	105	72	44	27	15	4	2	404
35–39	183	124	110	44	27	10	6	8	512
40–44	146	135	99	54	33	9	10	7	493
45–49	151	104	91	36	26	14	6	7	435
50–54	113	75	65	25	18	5	6	6	313
55–59	100	63	53	27	13	13	7	5	281
60–64	102	83	45	33	16	11	3	2	295
65–69	111	86	59	34	19	11	3	4	327
70–74	108	56	51	21	23	10	3	2	274
75–79	76	63	42	12	12	7	0	0	212
80–84	47	44	24	11	11	5	3	1	146
85+	36	30	17	12	9	2	0	0	106
Total	1,369	1,023	781	380	250	119	56	46	4,024

Source: National Cancer Statistics Clearing House (AIHW).

Table A15: Age-specific and age-standardised incidence rates for cervical cancer, by State and Territory, 1993–1996

Age group	NSW	Vic	Qld	WA	SA	Tas	ACT	NT	Australia
0–4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
5–9	0.0	0.0	0.0	0.0	0.5	0.0	0.0	0.0	0.0
10–14	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
15–19	0.4	0.0	0.4	0.0	0.0	0.0	0.0	0.0	0.2
20–24	1.2	2.1	2.5	1.9	0.0	1.5	0.0	0.0	1.6
25–29	5.1	5.6	7.8	8.3	7.0	9.0	9.8	5.7	6.4
30–34	13.6	14.2	13.9	15.6	11.6	20.0	7.7	6.0	13.8
35–39	19.2	17.5	22.0	15.8	11.8	13.4	11.7	27.1	18.1
40–44	16.5	20.3	21.0	20.5	15.2	13.0	19.8	27.5	18.6
45–49	18.5	17.1	20.8	15.6	12.9	22.2	13.0	34.0	17.9
50–54	17.7	16.0	19.5	14.5	11.6	10.2	19.4	44.2	16.8
55–59	18.4	15.7	19.7	18.9	9.7	30.6	31.7	59.0	18.0
60–64	20.2	22.3	18.8	26.4	12.7	28.1	17.6	34.4	20.6
65–69	21.9	23.3	25.1	28.9	14.5	28.7	19.2	97.9	23.0
70–74	23.6	16.9	24.5	20.7	19.0	28.2	22.1	72.2	21.6
75–79	22.5	26.2	27.0	16.2	13.4	26.0	0.0	0.0	22.7
80–84	19.5	24.8	22.0	19.6	16.9	26.1	50.9	110.9	21.6
85+	19.5	21.1	20.2	27.1	17.6	14.3	0.0	0.0	20.2
All ages									
AS Rate (A)	10.5	10.6	11.9	11.0	7.7	11.8	9.7	20.6	10.6
95% CI	10.0–11.0	10.0–11.0	11.0–13.0	10.0–12.0	7.0–9.0	10.0–14.0	7.0–12.0	15.0–27.0	10.0–11.0
AS Rate (W)	8.6	8.6	9.8	9.1	6.3	9.7	8.1	17.0	8.7
95% CI	8.1–9.0	8.1–9.1	9.1–10.4	8.2–10.0	5.5–7.1	8.0–11.4	6.0–10.3	12.1–21.9	8.4–9.0
Ages 20–69									
AS Rate (A)	14.0	14.3	16.0	15.2	10.2	15.8	13.3	27.1	14.3
95% CI	13.2–14.8	13.3–15.3	14.8–17.2	13.5–16.8	8.8–11.6	12.6–19.0	9.6–17.0	19.0–35.2	13.8–14.8
AS Rate (W)	13.7	13.9	15.6	14.8	9.9	15.5	13.3	26.8	14.0
95% CI	13.6–13.8	13.8–14.0	15.5–15.8	14.6–14.9	9.8–10.0	15.4–15.6	13.2–13.4	26.6–27.0	13.9–14.1

Note: Rates are expressed per 100,000 women and age standardised to the Australian 1991 population (A) and the World Standard Population (W).

Source: National Cancer Statistics Clearing House (AIHW).

Table A16: Number of new cases of cervical cancer by histological type for women aged 20–69 years, Australia, 1985–1996

Histological type	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996
Squamous	665	637	660	640	671	620	635	604	588	621	532	515
Adenocarcinoma	91	114	124	151	103	146	137	133	134	189	133	140
Adenosquamous	36	42	41	40	48	50	45	51	46	39	35	39
Other	81	70	71	69	74	79	78	60	76	68	69	57
Micro-invasive	103	67	89	84	89	138	156	149	134	170	175	144
Total	976	930	985	984	985	1,033	1,051	997	978	1,087	944	895

Source: National Cancer Statistics Clearing House (AIHW).

Table A17: Age-standardised incidence rates for cervical cancer by histological type for women aged 20–69 years, Australia, 1985–1996

Histological type	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996
Squamous	14.0	13.2	13.3	12.6	12.9	11.7	11.7	11.0	10.5	10.9	9.2	8.8
Adenocarcinoma	1.9	2.4	2.5	3.0	2.0	2.8	2.5	2.4	2.4	3.3	2.3	2.4
Adenosquamous	0.8	0.9	0.8	0.8	0.9	0.9	0.8	0.9	0.8	0.7	0.6	0.7
Other	1.7	1.4	1.4	1.3	1.4	1.5	1.4	1.1	1.4	1.2	1.2	1.0
Micro-invasive	2.2	1.4	1.8	1.7	1.7	2.6	2.9	2.7	2.4	3.0	3.1	2.5

Note: Rates are expressed per 100,000 women and age-standardised to the Australian 1991 population (A).

Source: National Cancer Statistics Clearing House (AIHW).

Indicator 8: Incidence by location

Table A18: Number of new cases of cervical cancer by age and location, 1993–95 to 1995–96

Age group	Metropolitan			Rural			Remote		
	1993–95	1994–96	1995–96	1993–95	1994–96	1995–96	1993–95	1994–96	1995–96
0–4	0	0	0	0	0	0	0	0	0
5–9	1	1	0	0	0	0	0	0	0
10–14	0	0	0	0	0	0	0	0	0
15–19	3	2	2	1	2	1	0	0	0
20–24	17	19	12	4	8	6	2	1	0
25–29	86	89	63	26	32	25	6	4	2
30–34	204	180	118	81	75	52	13	9	8
35–39	241	245	185	80	85	54	10	11	11
40–44	240	235	170	85	81	55	11	7	3
45–49	213	220	155	67	63	36	17	14	5
50–54	145	137	99	55	50	30	7	10	6
55–59	135	118	90	55	51	29	7	9	6
60–64	146	129	89	50	53	38	5	7	3
65–69	160	141	91	64	54	39	12	11	8
70–74	155	149	98	41	40	30	5	5	2
75–79	95	97	68	44	42	29	4	3	1
80–84	65	69	49	27	29	19	3	1	1
85+	52	55	45	16	16	10	2	2	2
Total	1,957	1,886	1,334	696	681	453	104	94	58

Notes: The numbers are presented as 3-year rolling blocks of data for the periods 1993–1995 and 1994–1996, and 2-yearly for the period 1995–1996.

Years refer to calendar years.

Source: National Cancer Statistics Clearing House (AIHW).

Table A19: Age-specific and age-standardised incidence rates for cervical cancer by age and location, 1993–95 to 1995–96

Age group	Metropolitan			Rural			Remote		
	1993–95	1994–96	1995–96	1993–95	1994–96	1995–96	1993–95	1994–96	1995–96
0–4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
5–9	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0
10–14	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
15–19	0.2	0.1	0.2	0.2	0.4	0.3	0.0	0.0	0.0
20–24	1.0	1.2	1.1	0.9	1.9	2.1	3.0	1.6	0.0
25–29	5.6	5.7	6.0	5.8	7.1	8.3	8.4	5.6	4.2
30–34	12.9	11.4	11.2	15.0	14.1	14.8	17.6	12.4	16.7
35–39	16.1	16.1	18.0	14.9	15.5	14.6	15.7	17.0	25.3
40–44	16.9	16.3	17.6	17.2	16.0	16.1	20.7	13.0	8.3
45–49	16.4	16.3	16.9	15.1	13.7	11.6	37.8	30.4	16.1
50–54	14.9	13.4	14.2	15.5	13.5	11.9	20.6	28.5	25.2
55–59	16.7	14.2	16.0	17.1	15.4	13.0	26.8	33.5	33.1
60–64	19.6	17.4	18.0	16.3	17.3	18.6	24.3	33.8	21.7
65–69	21.4	18.9	18.3	21.4	18.0	19.4	71.5	64.6	70.0
70–74	23.1	21.6	21.2	15.8	15.1	16.8	39.1	38.4	22.9
75–79	19.2	19.3	19.9	23.4	21.8	22.3	45.4	33.6	16.6
80–84	18.2	18.6	19.6	20.1	20.8	20.2	48.9	15.7	23.3
85+	18.7	18.9	22.7	15.9	15.0	13.6	37.7	34.4	49.2
All ages									
AS Rate (A)	9.7	9.2	9.6	9.5	9.2	9.2	16.5	14.8	13.3
	9.3–10.2	8.7–9.6	9.1–10.1	8.8–10.2	8.5–9.9	8.3–10.0	13.2–19.8	11.7–18.0	9.8–16.9
AS Rate (W)	8.0	7.5	7.9	7.8	7.5	7.5	13.3	12.4	10.9
	7.6–8.3	7.1–7.8	7.4–8.3	7.2–8.4	6.9–8.1	6.7–8.2	10.7–15.9	9.8–14.9	8.0–13.8
Ages 20–69									
AS Rate (A)	13.0	12.2	12.8	12.9	12.5	12.4	21.2	19.8	18.3
	12.4–13.7	11.6–12.8	12.1–13.6	11.9–14.0	11.4–13.5	11.1–13.6	16.7–25.8	15.4–24.2	13.2–23.5
AS Rate (W)	12.7	11.9	12.5	12.5	12.1	12	20.8	19.6	17.6
	12.4–13.6	11.6–12.8	12.1–13.6	11.9–14.0	11.5–13.5	11.1–13.6	16.8–25.7	15.5–24.2	13.4–23.3

Notes: Rates are expressed per 100,000 women and age-standardised to the Australian 1991 population (A) and the World Standard Population (W).

The age-standardised rates are presented as 3-year rolling blocks of data for the periods 1993–1995 and 1994–1996, and 2-yearly for the period 1995–1996.

Years refer to calendar years.

Source: National Cancer Statistics Clearing House.

Indicator 7: Mortality

Table A20: Number of deaths from cervical cancer by age, Australia, 1986–1998

Age group	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998
0–4	0	0	0	0	0	0	0	0	0	0	0	0	0
5–9	0	0	0	0	0	0	0	0	0	0	0	0	0
10–14	0	0	0	0	0	0	0	0	0	0	0	0	0
15–19	0	0	0	2	0	0	0	0	0	1	0	0	0
20–24	2	2	0	1	1	3	0	0	0	1	0	0	3
25–29	6	5	3	4	9	6	5	1	6	3	2	2	6
30–34	11	16	11	21	14	14	14	12	12	6	12	8	5
35–39	17	21	14	19	33	22	20	25	9	18	22	18	20
40–44	25	20	25	25	34	20	30	32	31	17	21	16	19
45–49	21	21	28	32	37	27	27	24	37	33	32	28	16
50–54	24	25	19	27	18	20	13	32	35	29	13	22	23
55–59	39	39	36	23	24	24	22	20	28	37	21	25	16
60–64	42	29	39	37	33	34	31	26	26	29	23	22	29
65–69	53	47	44	54	42	34	26	32	39	38	28	31	21
70–74	34	54	38	47	29	36	48	38	34	43	41	37	28
75–79	23	31	35	31	34	31	31	30	31	30	41	35	26
80–84	22	22	35	22	8	26	33	23	28	28	23	26	26
85+	22	16	18	24	23	34	22	23	25	21	22	30	31
Total	341	348	345	369	339	331	322	318	341	334	301	300	269

Note: Mortality data from 1986–1996 based on year of death and 1997 based on year of registration.

Source: AIHW Mortality Database.

Table A21: Age-specific and age-standardised death rates for cervical cancer by age, Australia, 1986–1998

Age group	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998
0–4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
5–9	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
10–14	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
15–19	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.0	0.0	0.2	0.0	0.0	0.0
20–24	0.3	0.3	0.0	0.2	0.1	0.4	0.0	0.0	0.0	0.1	0.0	0.0	0.4
25–29	0.9	0.7	0.4	0.6	1.3	0.9	0.7	0.1	0.9	0.4	0.3	0.3	0.8
30–34	1.7	2.5	1.7	3.1	2.0	2.0	1.9	1.6	1.6	0.8	1.7	1.1	0.7
35–39	2.7	3.4	2.2	2.9	5.0	3.3	3.0	3.6	1.3	2.5	3.0	2.4	2.7
40–44	5.1	3.7	4.4	4.2	5.5	3.1	4.7	4.9	4.7	2.5	3.1	2.3	2.7
45–49	5.1	5.0	6.4	7.0	7.7	5.4	5.0	4.2	6.2	5.4	5.0	4.4	2.5
50–54	6.7	6.8	5.0	6.9	4.5	4.8	3.1	7.4	7.7	6.1	2.6	4.1	4.0
55–59	10.5	10.6	9.9	6.4	6.7	6.7	6.0	5.3	7.3	9.4	5.2	6.0	3.7
60–64	11.4	7.9	10.5	10.0	8.9	9.2	8.5	7.2	7.3	8.1	6.4	6.1	7.8
65–69	17.4	14.9	13.4	15.7	12.0	9.7	7.4	9.0	11.0	10.7	7.9	8.8	6.0
70–74	12.9	20.2	14.2	17.7	10.7	12.8	16.4	12.5	10.7	13.3	12.5	11.3	8.5
75–79	12.0	15.6	17.0	14.4	15.4	13.7	13.5	13.0	13.6	12.9	16.8	13.7	9.7
80–84	18.5	17.8	27.1	16.4	5.7	17.9	21.8	14.5	16.7	16.2	13.0	14.5	14.5
85+	23.3	16.5	18.0	23.3	21.8	30.9	19.0	18.9	19.6	15.6	15.5	20.1	19.6
All ages													
AS Rate (A)	4.2	4.1	4.0	4.2	3.8	3.6	3.4	3.3	3.5	3.3	2.9	2.8	2.5
AS Rate (W)	3.3	3.2	3.1	3.3	3.0	2.7	2.6	2.6	2.7	2.6	2.2	2.1	1.9
Ages 20–69													
AS Rate (A)	4.9	4.5	4.3	4.6	4.6	3.8	3.4	3.6	3.9	3.6	2.9	2.8	2.6
AS Rate (W)	4.8	4.4	4.3	4.6	4.5	3.7	3.4	3.6	3.9	3.7	2.9	2.8	2.6

Notes: (1) Mortality data from 1993–1996 based on year of death and 1997 based on year of registration.

(2) Rates are expressed per 100,000 women and age-standardised to the Australian 1991 population (A) and the World Standard Population (W).

Source: AIHW Mortality Database.

Table A22: Number of deaths from cervical cancer by age, State and Territory, 1989–1992

Age group	NSW	Vic	Qld	WA	SA	Tas	ACT	NT	Australia
0–4	0	0	0	0	0	0	0	0	0
5–9	0	0	0	0	0	0	0	0	0
10–14	0	0	0	0	0	0	0	0	0
15–19	0	0	0	1	1	0	0	0	2
20–24	0	2	2	0	1	0	0	0	5
25–29	7	3	5	5	4	0	0	0	24
30–34	21	14	8	6	8	3	0	3	63
35–39	33	20	19	9	7	2	0	4	94
40–44	43	25	16	12	10	1	0	2	109
45–49	35	34	17	17	8	5	3	4	123
50–54	27	19	12	8	5	2	2	3	78
55–59	24	29	10	11	4	6	4	5	93
60–64	53	32	20	10	12	7	1	0	135
65–69	59	41	22	15	9	10	0	0	156
70–74	58	39	23	13	18	5	1	3	160
75–79	44	38	18	11	9	6	0	1	127
80–84	35	19	18	6	5	4	1	1	89
85+	34	17	26	11	12	3	0	0	103
Total	473	332	216	135	113	54	12	26	1,361

Note: Mortality data in this table are based on year of death.

Table A23: Age-specific and age-standardised death rates for cervical cancer by age, by State and Territory, 1989–1992

Age group	NSW	Vic	Qld	WA	SA	Tas	ACT	NT	Australia
0–4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
5–9	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
10–14	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
15–19	0.0	0.0	0.0	0.4	0.5	0.0	0.0	0.0	0.1
20–24	0.0	0.3	0.0	0.0	0.4	0.0	0.0	0.0	0.2
25–29	0.7	0.4	1.5	1.9	1.7	0.0	0.0	0.0	0.9
30–34	2.2	1.9	4.5	2.2	3.4	4.0	0.0	9.4	2.2
35–39	3.7	3.0	7.4	3.4	3.2	2.9	0.0	14.5	3.6
40–44	5.1	3.9	10.0	5.0	4.8	1.5	0.0	8.6	4.4
45–49	5.1	6.8	10.3	9.3	4.8	9.6	8.5	25.0	6.2
50–54	4.8	4.5	10.0	5.4	3.6	4.5	8.0	28.6	4.8
55–59	4.7	7.7	10.2	8.6	3.2	15.1	21.6	70.2	6.4
60–64	10.0	8.3	22.2	8.2	8.8	17.5	6.1	0.0	9.1
65–69	11.7	11.2	26.4	13.5	6.8	25.5	0.0	0.0	11.2
70–74	14.3	13.7	31.9	14.9	16.9	15.6	9.6	121.1	14.4
75–79	13.7	16.2	31.0	15.1	10.5	23.4	0.0	70.2	14.3
80–84	17.2	12.4	39.2	12.7	9.1	24.8	23.0	141.6	15.6
85+	22.2	14.1	49.5	31.2	28.5	26.3	0.0	0.0	23.7
All ages									
AS Rate (A)	3.7	3.5	3.6	4.3	3.4	5.4	2.7	15.4	3.7
95% CI	3.3–4.0	3.1–3.9	3.1–4.0	3.6–5.0	2.8–4.1	3.9–6.8	1.1–4.2	8.1–22.7	3.5–3.9
AS Rate (W)	2.8	2.8	2.7	3.4	2.7	4.2	2.3	11.5	2.9
95% CI	2.6–3.1	2.4–3.1	2.3–3.1	2.8–4.0	2.2–3.2	3.0–5.3	1.0–3.7	6.6–16.4	2.7–3.1
Ages 20–69									
AS Rate (A)	4.0	3.9	3.7	4.8	3.7	6.1	3.3	13.3	4.1
95% CI	3.6–4.5	3.4–4.5	3.0–4.3	3.8–5.8	2.8–4.5	4.1–8.1	1.2–5.4	7.4–19.3	3.8–4.4
AS Rate (W)	3.9	3.9	3.6	4.8	3.6	6.0	3.6	13.7	4.0
95% CI	3.6–4.5	3.4–4.5	3.0–4.3	3.8–5.8	2.8–4.5	4.1–8.1	1.0–5.5	7.0–19.6	3.8–4.4

Notes: (1) The age-standardised rates were averaged over 4 years to smooth annual variations that may occur in the smaller States and Territories.

(2) Deaths derived from place of usual residence and not place of death.

(3) Mortality data in this table are based on year of death.

(4) Rates are expressed per 100,000 women and age-standardised to the Australian 1991 population (A) and the World Standard Population.

Source: AIHW Mortality Database.

Table A24: Number of deaths from cervical cancer by age, State and Territory, 1995–1998

Age group	NSW	Vic	Qld	WA	SA	Tas	ACT	NT	Australia
0–4	0	0	0	0	0	0	0	0	0
5–9	0	0	0	0	0	0	0	0	0
10–14	0	0	0	0	0	0	0	0	0
15–19	1	0	0	0	0	0	0	0	1
20–24	0	2	2	0	0	0	0	0	4
25–29	0	3	5	2	1	0	1	0	12
30–34	8	8	7	7	2	1	0	0	33
35–39	28	15	13	11	5	3	0	2	77
40–44	35	14	12	6	3	2	3	1	76
45–49	43	22	14	11	5	4	4	5	108
50–54	37	17	16	8	2	2	3	0	85
55–59	33	15	20	12	9	6	2	1	98
60–64	35	22	20	6	8	9	1	2	103
65–69	45	33	17	12	5	4	2	2	120
70–74	57	38	27	8	12	6	2	1	151
75–79	47	29	31	5	11	6	1	1	131
80–84	33	30	14	9	14	1	1	0	102
85+	39	24	12	16	9	3	2	0	105
Total	441	272	210	113	86	47	22	15	1206

Note: Mortality data from 1993–1996 based on year of death and 1997 based on year of registration.
Source: AIHW Mortality Database.

Table A25: Age-specific and age-standardised death rates for cervical cancer by age, by State and Territory, 1995–1998

Age group	NSW	Vic	Qld	WA	SA	Tas	ACT	NT	Australia
0–4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
5–9	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
10–14	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
15–19	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
20–24	0.0	0.3	0.4	0.0	0.0	0.0	0.0	0.0	0.1
25–29	0.0	0.4	1.0	0.7	0.5	0.0	1.9	0.0	0.4
30–34	0.8	1.1	1.4	2.5	0.9	1.4	0.0	0.0	1.1
35–39	2.8	2.1	2.5	3.8	2.1	4.0	0.0	6.5	2.6
40–44	3.8	2.0	2.4	2.2	1.4	2.8	6.0	3.7	2.8
45–49	5.1	3.5	3.0	4.4	2.4	6.1	8.2	22.4	4.2
50–54	5.2	3.3	4.2	4.1	1.2	3.7	8.4	0.0	4.1
55–59	5.8	3.6	6.8	7.9	6.5	13.5	8.2	10.4	5.9
60–64	6.9	5.9	8.1	4.7	6.4	22.9	5.5	31.3	7.1
65–69	9.0	9.0	7.1	10.0	3.9	10.5	12.7	45.5	8.5
70–74	12.2	11.1	12.5	7.6	9.7	16.9	14.0	33.1	11.5
75–79	13.1	11.3	18.4	6.3	11.5	21.3	9.8	51.4	13.1
80–84	13.1	16.3	12.0	15.3	20.6	4.9	15.3	0.0	14.4
85+	19.0	15.4	12.7	32.3	15.9	19.3	41.9	0.0	18.1
All ages									
AS Rate (A)	3.0	2.5	2.9	3.0	2.2	4.3	3.8	7.8	2.9
95% CI	2.8–3.3	2.3–2.8	2.6–3.3	2.5–3.4	1.8–2.7	3.3–5.4	2.4–5.2	4.0–11.7	2.7–3.0
AS Rate (W)	2.3	1.9	2.2	2.3	1.6	3.5	3.0	6.2	2.2
95% CI	2.1–2.5	1.7–2.1	2.0–2.5	1.9–2.7	1.3–2.0	2.6–4.4	1.9–4.2	3.4–9.3	2.1–2.3
Ages 20–69									
AS Rate (A)	3.2	2.5	3.0	3.4	2.0	5.0	4.1	9.0	3.0
95% CI	2.9–3.5	2.2–2.9	2.6–3.4	2.7–4.0	1.5–2.6	3.6–6.5	2.4–6.1	4.7–13.9	2.8–3.2
AS Rate (W)	3.2	2.5	3.0	3.3	2.0	5.0	4.2	8.9	3.0
95% CI	2.8–3.5	2.1–2.8	2.6–3.4	2.7–3.9	1.5–2.6	3.6–6.6	2.4–6.1	4.6–13.8	2.8–3.2

Notes: (1) The age-standardised rates were averaged over 4 years to smooth annual variations that may occur in the smaller States and Territories.

(2) Deaths derived from place of usual residence and not place of death.

(3) Mortality data from 1995–1996 based on year of death and 1997 based on year of registration.

(4) Rates are expressed per 100,000 women and age-standardised to the Australian 1991 population (A) and the World Standard Population.

Source: AIHW Mortality Database.

Indicator 9: Mortality by location

Table A26: Number of deaths from cervical cancer by age and location, 1993–1995 to 1995–1997

Age group	Metropolitan			Rural			Remote		
	1993–95	1994–96	1995–97	1993–95	1994–96	1995–97	1993–95	1994–96	1995–97
0–4	0	0	0	0	0	0	0	0	0
5–9	0	0	0	0	0	0	0	0	0
10–14	0	0	0	0	0	0	0	0	0
15–19	1	1	1	0	0	0	0	0	0
20–24	0	1	1	0	0	0	0	0	0
25–29	8	8	3	2	2	3	1	0	0
30–34	16	21	19	12	9	9	1	1	0
35–39	38	39	43	11	9	13	3	2	1
40–44	58	42	35	22	26	21	3	2	1
45–49	67	70	62	19	24	23	6	6	7
50–54	65	55	44	26	20	16	4	3	2
55–59	57	59	57	21	22	24	4	3	1
60–64	54	51	44	23	20	27	4	5	3
65–69	66	66	66	35	35	28	6	5	5
70–74	85	86	84	30	32	36	2	2	3
75–79	64	73	75	26	27	29	1	1	1
80–84	55	57	52	23	20	22	1	0	2
85+	47	53	57	21	15	17	0	0	0
Total	681	682	643	271	261	268	36	30	26

Notes: (1) Deaths derived from place of usual residence and not place of death.

(2) Mortality data from 1983–1996 based on year of death and 1997 based on year of registration.

(3) The number of deaths are presented as 3-year rolling blocks of data.

(4) Years refer to calendar years.

Source: AIHW Mortality Database.

Table A27: Age-specific and age-standardised death rates for cervical cancer by age and location, 1993–1995 to 1995–1997

Age group	Metropolitan			Rural			Remote		
	1993–95	1994–96	1995–97	1993–95	1994–96	1995–97	1993–95	1994–96	1995–97
0–4	0	0	0	0	0	0	0	0	0
5–9	0	0	0	0	0	0	0	0	0
10–14	0	0	0	0	0	0	0	0	0
15–19	0.1	0.1	0.1	0	0	0	0	0	0
20–24	0	0.1	0.1	0	0	0	0	0	0
25–29	0.5	0.5	0.2	0.4	0.4	0.7	1.4	0	0
30–34	1.01	1.3	1.2	2.2	1.7	1.7	1.4	1.4	0
35–39	2.5	2.6	2.8	2.1	1.6	2.4	4.7	3.1	1.5
40–44	4.1	2.9	2.4	4.4	5.1	4.1	5.6	3.7	1.9
45–49	5.2	5.2	4.6	4.3	5.2	5.0	13.3	13.0	15.2
50–54	6.7	5.4	4.3	7.3	5.4	4.3	11.8	8.6	5.7
55–59	7.1	7.1	6.8	6.5	6.7	7.3	15.3	11.2	3.7
60–64	7.2	6.9	5.9	7.5	6.5	8.8	19.4	24.2	14.5
65–69	8.8	8.9	8.9	11.7	11.6	9.3	35.8	29.4	29.4
70–74	12.7	12.5	12.2	11.6	12.1	13.6	15.6	15.4	23.1
75–79	13.0	14.5	14.9	13.8	14.0	15.1	11.3	11.2	11.2
80–84	15.4	15.4	14	17.2	14.4	15.8	16.3	0.0	31.4
85+	16.9	18.2	19.6	20.8	14.0	15.9	0.0	0.0	0
All ages									
AS Rate (A)	3.2	3.1	2.9	3.4	3.3	3.3	6.2	5.2	4.7
95% CI	3.0–3.5	2.9–3.4	2.7–3.2	3.0–3.9	2.9–3.7	2.9–3.7	4.1–8.3	3.3–7.1	2.9–6.6
AS Rate (W)	2.5	2.4	2.2	2.7	2.5	2.6	5.2	4.4	3.7
95% CI	2.3–2.7	2.2–2.6	2.1–2.4	2.3–3.0	2.2–2.9	2.2–2.9	3.4–6.9	2.8–6.0	2.3–5.2
Ages 20–69									
AS Rate (A)	3.5	3.3	3.0	3.8	3.6	3.6	8.3	7.1	5.4
95% CI	3.2–3.8	3.0–3.6	2.7–3.3	3.2–4.3	3.1–4.2	3.0–4.1	5.4–11.3	4.3–9.9	2.9–7.7
AS Rate (W)	3.5	3.3	2.9	3.7	3.6	3.5	8.3	7.1	5.4
95% CI	3.2–3.9	3.0–3.6	2.7–3.3	3.2–4.3	3.1–4.2	3.0–4.1	5.4–11.3	4.4–9.9	3.0–7.7

Notes: (1) Deaths derived from place of usual residence and not place of death.

(2) Mortality data from 1983–1996 based on year of death and 1997 based on year of registration.

(3) Rates are expressed per 100,000 women and age-standardised to the Australian 1991 population (A) and the World Standard Population (W).

(4) The age-standardised rates are presented as 3-year rolling blocks of data.

(5) Years refer to calendar years.

Source: AIHW Mortality Database.

Indicator 10: Indigenous mortality

Table A28: Number of deaths from cervical cancer by age, by Indigenous and non-Indigenous status, SA, WA, NT, 1993 –1995 to 1995–1997

Age group	Indigenous			Non-Indigenous		
	1993–1995	1994–1996	1995–1997	1993–1995	1994–1996	1995–1997
0–4	0	0	0	0	0	0
5–9	0	0	0	0	0	0
10–14	0	0	0	0	0	0
15–19	0	0	0	1	1	1
20–24	0	0	0	1	1	1
25–29	0	0	0	10	11	7
30–34	0	0	0	30	30	26
35–39	0	0	0	52	49	58
40–44	1	1	1	80	69	54
45–49	5	4	7	94	102	93
50–54	3	0	0	96	77	64
55–59	1	1	2	85	86	83
60–64	2	3	2	81	78	74
65–69	2	3	3	109	105	97
70–74	1	2	3	115	118	121
75+	1	0	1	239	249	256
Total	16	14	19	993	976	935

Note: (1) Only South Australia, Western Australia and the Northern Territory have Indigenous death registration data considered to be of a publishable standard.

(2) Mortality data from 1993–1996 based on year of death and 1997 based on year of registration.

(3) Rates are expressed per 100,000 women and age standardised to the Australian 1991 population (A) and the World Standard Population (W).

(4) The number of deaths are presented as 3-year rolling blocks of data.

Source: AIHW Mortality Database.

Table A29: Age-specific and age-standardised death rates for cervical cancer by age, by Indigenous and non-Indigenous status, SA, WA, NT, 1993–1995 to 1995–1997

Age group	Indigenous			non-Indigenous		
	1993–1995	1994–1996	1995–1997	1993–1995	1994–1996	1995–1997
0–4	0.0	0.0	0.0	0.0	0.0	0.0
5–9	0.0	0.0	0.0	0.0	0.0	0.0
10–14	0.0	0.0	0.0	0.0	0.0	0.0
15–19	0.0	0.0	0.0	0.1	0.1	0.1
20–24	0.0	0.0	0.0	0.1	0.1	0.1
25–29	0.0	0.0	0.0	0.5	0.5	0.3
30–34	0.0	0.0	0.0	1.4	1.4	1.2
35–39	0.0	0.0	0.0	2.5	2.3	2.7
40–44	11.1	10.5	11.4	4.1	3.4	2.7
45–49	71.1	54.7	107.0	5.3	5.5	4.9
50–54	61.1	0.0	0.0	7.0	5.4	4.2
55–59	23.4	23.7	54.8	7.4	7.2	6.8
60–64	62.3	90.5	69.4	7.6	7.3	6.9
65–69	88.3	127.3	139.7	10.2	9.9	9.2
70–74	76.8	146.8	239.0	12.2	12.2	12.4
75+	53.1	0.0	58.3	15.2	15.3	15.2
All ages						
AS Rate (A)	19.8	18.2	27.6	3.4	3.2	3.0
95% CI	9.1–28.8	8.2–29.0	13.1–37.5	2.5–3.4	2.2–3.0	1.9–2.8
AS Rate (W)	16.7	15.2	22.2	2.6	2.5	2.3
95% CI	8.1–24.5	7.1–23.7	11.0–30.2	2.1–2.9	1.8–2.6	1.6–2.3
Ages 20–69						
AS Rate (A)	24.1	22.0	28.5	3.7	3.5	3.1
95% CI	11.0–37.2	9.0–35.1	12.7–41.1	3.2–4.5	2.8–4.1	2.5–3.7
AS Rate (W)	25.1	22.0	29.0	3.7	3.5	3.1
95% CI	10.7–37.5	9.3–34.8	12.7–41.1	3.1–4.5	2.8–4.1	2.5–3.7

Note: (1) Only South Australia, Western Australia and the Northern Territory have Indigenous death registration data considered to be of a publishable standard.

(2) Mortality data from 1993–1996 based on year of death and 1997 based on year of registration.

(3) Rates are expressed per 100,000 women and age standardised to the Australian 1991 population (A) and the World Standard Population (W).

(4) The age-standardised rates are presented as 3-year rolling blocks of data.

Source: AIHW Mortality Database.