



Australian Government

Australian Institute of Health and Welfare



infocus

australian
mesothelioma
registry

Mesothelioma in Australia 2020

Published November 2021

Australia has one of the highest measured incidence rates of mesothelioma in the world (Bray et al. 2017). Each year in Australia, between 700 and 800 people are diagnosed with the rare and aggressive cancer. In the 2018–19 financial year, the estimated health system expenditure for mesothelioma cases was \$27.4m (total cancers \$11.8 billion) (AIHW 2021).

Men were more likely to be diagnosed with mesothelioma than women across all age groups, and the number of cases diagnosed each year for both men and women has steadily increased over the past 40 years. There is no cure for mesothelioma. The main cause is exposure to asbestos—a material that has been banned in Australia since 2003.

This report presents the latest available statistics from the Australian Mesothelioma Registry (AMR), supplemented by data from the National Mortality Database (NMD), the Australian Cancer Database (ACD) and the National Death Index (NDI).

Main findings



642 cases of mesothelioma diagnosed in 2020 had been reported to the AMR as at 1 May 2021—the **median age at diagnosis was 75**.



Between 1988–1992 and 2013–2017, the age-adjusted **relative survival of people with mesothelioma has increased**, most notably 1-year relative survival.



In 2020, **696 deaths of people** with mesothelioma (from any cause of death including mesothelioma) were recorded on the AMR—a **mortality rate of 2.1 deaths per 100,000 population**.



More than **9 in 10** of the exposure assessment participants were assessed as having **possible or probable exposure** to asbestos.

The AMR collects information on new cases of mesothelioma diagnosed in Australia since 1 July 2010. The Registry's main goals are to better understand the relationship between asbestos exposure and mesothelioma, to assist in the development of policies to best deal with asbestos still in the environment, and to provide reliable information to policy makers and researchers. For more information on the AMR, see [*Mesothelioma in Australia 2020: methodology paper*](#).

AIHW

Stronger evidence,
better decisions,
improved health and welfare

What is mesothelioma?

Mesothelioma is a form of cancer in the mesothelium—the protective lining on the inside of body cavities and the outside of internal organs. In 2020, around 95% of cases reported to the AMR (for which tumour location information was available) were pleural mesothelioma—that is, the cancer occurs around the lungs. The other 5% of cases were mesothelioma in other areas of the body. More information on the diagnostic characteristics of cases of mesothelioma diagnosed in 2020 is available in Table A10 of [*Mesothelioma in Australia 2020—data tables*](#). Although mesothelioma most commonly occurs in the chest, it is not a lung cancer and requires different forms of treatment (Cancer Council 2019a).

Although mesothelioma is not curable, patients can undergo one or more different treatments to increase their survival period—including chemotherapy, radiotherapy, and surgery (or all three, which is referred to as ‘trimodality therapy’) (Cancer Council 2019b). However, in the more advanced stages of the disease, the focus is on improving quality of life and easing symptoms (Cancer Council 2019c).

The main cause of mesothelioma is exposure to asbestos—a group of naturally occurring fibrous silicate materials that are invisible to the naked eye and can be inhaled into the lungs (AMR 2017) where they do not readily break down.

Australia's use of asbestos


Australia's consumption of asbestos peaked at a total of around 700,000 metric tonnes during the period 1970–1979 (Soeberg 2016; Leigh et al. 2002). Australia mined and imported asbestos, which was primarily used in the construction and transport industries because it was durable and resistant to fire and chemicals (ASEA 2016). Asbestos-related regulatory controls have been significantly tightened over time—asbestos-containing materials (ACMs) have been banned in Australia since December 2003, and it is now illegal to make, use or import it from another country. However, a large amount of asbestos still remains in older structures and products, potentially exposing workers and/or the public to asbestos if relevant safety procedures that have been identified are not followed. The Commonwealth and state and territory work health and safety authorities have put in place strict asbestos-related regulatory measures to control asbestos exposure in workplaces (including safe management and removal of asbestos), to minimise asbestos-related diseases such as mesothelioma, asbestosis and lung cancer.

How have the number of cases changed over time?

The AMR is the most up-to-date source of national data on mesothelioma incidence (number of new cases) in Australia, because case notifications are fast-tracked from state and territory cancer registries to the AMR. However, most notifications are still received in the year after diagnosis because of the time it takes to make a definitive diagnosis, and the time between diagnosis and notification to cancer registries and to the AMR. For more information on AMR processes, see [*Mesothelioma in Australia 2020: methodology paper*](#).

Because not all cases of mesothelioma are reported to the AMR in the year that they are diagnosed, the number of cases recorded for each year continues to rise for all years dating back to 2011 (Figure 1). For example, of the 719 cases of mesothelioma diagnosed in 2013 and notified to the AMR, 47% of notifications were received in the year they were diagnosed (2013), 44% in 2014, and 9% were received from 2015 onwards.

Figure 1: Difference in number of mesothelioma cases between originally published data¹ and current report during 2011 to 2019




¹ Original published data refers to the number of cases notified at the time those data were initially published.

Sources: AMR 2012, 2013, 2014, 2015, 2016, 2017; AIHW 2018b; AIHW 2019; AIHW 2020; AIHW analysis of AMR data at 1 May 2021; Table A1 in [Mesothelioma in Australia 2020—data tables](#).

At 1 May 2021, 642 cases of mesothelioma diagnosed in 2020 had been reported to the AMR (Figure 2). Men have been consistently more likely to be diagnosed with mesothelioma than women—this is expected because the majority of cases are from exposure to asbestos in the type of environments in which men more commonly work, such as the mining and construction industries. After adjusting for differences in the age structure of the populations, current available data show that the rate for persons averaged around 2.7 cases per 100,000 people between 2011 and 2020; for men, it averaged 4.7 per 100,000, and for women, it averaged 0.9 per 100,000.

Figure 2: Number and age-standardised rate (per 100,000 population) of people diagnosed with mesothelioma, by year and sex, 2011 to 2020



Note: Rates have been age-standardised to the 2001 Australian Standard Population.


Source: AIHW analysis of AMR data at 1 May 2021; Table A2 in [Mesothelioma in Australia 2020—data tables](#).

Long-term trend shows a steady increase

Reporting using AMR data is only possible for the period since July 2010 as that is when the data collection began. However, longer-term trends in the number and incidence rates of mesothelioma cases can be described by combining data from the ACD for 1982–2010 and the AMR for 2011–2020.

Between 1982 and 2020, the number of new cases of mesothelioma reported annually steadily increased—from 135 to 499 for men and from 22 to 143 for women (Figure 3). To date, the highest overall number of cases (819) were those with a date of diagnosis in 2017. However, it is important to note that the apparent fall in cases in 2020 is likely due to delays in the AMR receiving notifications—the number of cases for 2020 is expected to rise in 2021 and subsequent years.


Figure 3: Number of people diagnosed with mesothelioma, by year and sex, 1982 to 2020



Sources: 1982–2010 (dotted line): AIHW 2018a; 2011–2020 (solid line): AIHW analysis of AMR data at 1 May 2021; Table A3 in [Mesothelioma in Australia 2020—data tables](#)

Between 1982 and 2003, the age-adjusted incidence rate rose for both men (2.3 to 5.9 cases per 100,000 population) and women (0.3 to 1.0 cases per 100,000 population) (Figure 4). The overall incidence rate rose from 1.2 to a peak of 3.2 cases per 100,000 population in 2003, and has since remained at around 2.8 cases per 100,000 population.

Figure 4: Age-standardised rate (per 100,000 population) of people diagnosed with mesothelioma, by year and sex, 1982 to 2020



Note: Rates have been age-standardised to the 2001 Australian Standard Population.


Sources: 1982–2010 (dotted line): AIHW 2018a; 2011–2020 (solid line): AIHW analysis of AMR data at 1 May 2021; Table A3 in [Mesothelioma in Australia 2020—data tables](#).

How does mesothelioma diagnosis vary by age and sex?

The age of people in the AMR who were diagnosed in 2020 ranged from 19 to 101 (Figure 5). The median age at diagnosis was 75, and a greater number of men than women were diagnosed across most age groups.

Mesothelioma has a long and highly varied latency period (Marinaccio et al. 2007; Shavelle et al. 2017), with symptoms typically appearing decades after a person has been exposed to asbestos. A study by Reid et al. (2014) that analysed the data of more than 22,000 people who had been exposed to asbestos found that for those with mesothelioma, the median time since first exposure to diagnosis was 38 years. A similar study by Olsen et al. (2011) based on data from people diagnosed in Western Australia between 1960 and 2008 reported a latency period of between 33 and 44 years. Most mesothelioma diagnoses are made later in life.


Figure 5: Number of people diagnosed with mesothelioma, by age group and sex, 2020



Source: AIHW analysis of AMR data at 1 May 2021; Table A4 in [Mesothelioma in Australia 2020—data tables](#).

In 2020, age-specific mesothelioma incidence rates (the number of new mesothelioma cases per 100,000 population in specific age groups) generally increased with increasing age for men up to age 80 to 84 (43.8 cases per 100,000 population) and for women aged 85 and over (8.7 cases per 100,000 population). Men consistently had higher rates than women across all age groups, with men aged 80 to 84 being almost 6 times as likely to be diagnosed as women (Figure 6).

Figure 6: Rate (per 100,000 population) diagnosed with mesothelioma, by age and sex, 2020



Source: AIHW analysis of AMR data at 1 May 2021; Table A4 in [Mesothelioma in Australia 2020—data tables](#)

How do rates vary by state and territory?

Because of the small number of people diagnosed in many age groups in some states and territories, data have been grouped for the years 2017–2020. This enables the data to be directly age standardised to remove the effect of differing age structures between the populations, which can affect incidence rates. During the period 2017–2020, age-standardised rates of mesothelioma ranged from 1.2 cases per 100,000 people in the Northern Territory to 3.5 cases per 100,000 people in Western Australia (Table 1) – likely due to extensive asbestos mining in the past (Cancer Council 2019a). Additional data are available for 2013–2016 in [Mesothelioma in Australia 2020—data tables](#).

Table 1: Number and age-standardised rate (per 100,000 population) of people diagnosed with mesothelioma, by sex and state/territory, 2017–2020

State of diagnosis	Males		Females		People	
	No.	Rate	No.	Rate	No.	Rate
NSW	754	4.0	170	0.8	924	2.3
Vic	532	3.7	134	0.8	666	2.1
Qld	500	4.4	134	1.0	634	2.6
WA	328	5.9	85	1.3	413	3.5
SA	193	4.3	56	1.1	249	2.5
Tas	57	3.8	7	0.4	64	2.0
ACT	24	3.2	5	0.5	29	1.7
NT	5	2.2	2	0.4	7	1.2
Australia	2,393	4.2	593	0.9	2,986	2.2

Note: Due to small counts in some states and territories, data have been grouped into the years 2017–2020, to enable rates to be directly age-standardised to the 2001 Australian Standard Population.

Source: AIHW analysis of AMR data at 1 May 2021.

How long do people live after diagnosis?

What is relative survival?

Relative survival is a measure of the survival of people with cancer compared with that of the general population, and is calculated by dividing the observed survival (of people with cancer) by expected survival (of the general population), where the numerator and denominator have been matched for age, sex and calendar year.

The population diagnosed with mesothelioma continues to change over time and is more commonly older in more recent years, averaging around 75 years of age at diagnosis between 2011 and 2020 (AMR data). Even though relative survival for people with mesothelioma is improving over time, the extent of improvement reflected in relative survival rates is diminished because an increasing proportion of people with mesothelioma are older in more recent years than in the past, and survival rates for older people tend to be lower.


By age-adjusting to a group with a consistent age structure, such as those people with mesothelioma in 2013–2017, age-adjusted relative survival rates account for the impact different age structure can have on comparisons over time in relative survival rates.

Note that results presented here are not comparable with the survival rates presented in previous AMR reports. For further information see [Mesothelioma in Australia 2020: methodology paper](#).

Due to its aggressive nature, mesothelioma has a very low survival rate, and this has shown little improvement over time (Faig et al. 2015). The condition is often diagnosed at advanced stages, because early symptoms can go unnoticed or be mistaken as symptoms for similar conditions or diseases (Asbestos Diseases Research Institute 2019; Cancer Council 2019a). Many factors can affect a person's chances of survival, including their age at diagnosis and overall health status, the type of mesothelioma they have, and level of exposure to asbestos (such as whether it was occupational or non-occupational and the duration of exposure) (Burgers & Damhuis 2004).

This section, which presents linked data from the ACD and NDI, shows that the average length of time that people survive after being diagnosed with mesothelioma is gradually improving over time. Figure 7 shows the 1, 3 and 5-year age-adjusted relative survival of people with mesothelioma from 1988–1992 to 2013–2017—all survival rates have increased over this period, most notably the 1-year survival. This improvement in survival may be due to factors such as earlier diagnosis and improved treatment (Musk et al. 2011).

Figure 7: 1, 3 and 5-year age-adjusted relative survival of people diagnosed with mesothelioma, 1988–1992 to 2013–2017



Notes:

1. Age-adjusted relative survival rates: use a single population (the base population) and apply the survival rates (by age) for the respective time periods to this population; the base population is the 2013–2017 relative survival reporting population diagnosed with mesothelioma (that is, the group diagnosed with mesothelioma and used to calculate the 2013–2017 mesothelioma relative survival rates). These age-adjusted relative survival rates are not directly comparable with survival rates for other sexes or cancers.
2. The error bars presented in this figure represent 95% confidence intervals. A 95% confidence interval (CI) implies that there is 95% confidence that the true value will be included in this interval.

Sources: AIHW Australian Cancer Database 2017; National Death Index; Table A5 in [Mesothelioma in Australia 2020—data tables](#).

Death rate remains stable

The AMR is linked to the NDI to determine how many people registered on the AMR had subsequently died. Date and cause of death information from state and territory cancer registries was used where NDI data was not available. In 2020, 696 deaths of people with mesothelioma (with any cause of death) were recorded on the AMR—a rate of 2.1 deaths per 100,000 population (Table 2). The number of recorded deaths is expected to increase as more information becomes available over time. Where cause of death information was available for 2012–2020, mesothelioma was recorded as the primary (or ‘underlying’) cause of more than 90.7% of deaths among people with mesothelioma each year.

Table 2: Number and age-standardised rate (per 100,000 population) of deaths among people with mesothelioma, by year and sex, 2012 to 2020

Year of death	Males		Females		People	
	No.	Rate	No.	Rate	No.	Rate
2012	492	4.3	93	0.6	585	2.3
2013	547	4.6	98	0.7	645	2.4
2014	582	4.8	118	0.8	700	2.6
2015	550	4.3	146	1.0	696	2.5
2016	577	4.6	120	0.8	697	2.4
2017	624	4.3	133	0.8	757	2.6
2018	606	4.3	135	0.8	741	2.4
2019	613	4.2	145	0.9	758	2.4
2020	540	3.6	156	0.9	696	2.1

Notes:


1. The total number of deaths for 2020 is preliminary and is expected to rise as more information becomes available.
2. Rates have been age-standardised to the 2001 Australian Standard Population.

Source: AMR data at 1 May 2021.

This report uses the NMD to report on long-term mortality trends. The number of mesothelioma deaths has increased from 416 in 1997 to 731 in 2019. Deaths from mesothelioma increased generally but with fluctuations over the period among both men and women. For men, the number of deaths in 2019 was 1.7 times higher than in 1997, while for women there were over twice as many deaths in 2019 compared to 1997 (Figure 8).

Although the number of deaths has increased over time, the age standardised rate of deaths has remained fairly stable. The most likely reason for this is Australia’s increasing and ageing population (whereby older Australians are accounting for an increasing proportion of the population) (ABS 2019).

Figure 8: Number of deaths from mesothelioma, by year and sex, 1997 to 2019




Notes:

1. Figures are presented by year of death for 1997–2018 and year of registration of death for 2019.
2. Deaths registered in 2015 and earlier are based on the final version of the Cause of Death Unit Record File (CODURF), deaths registered in 2016 are based on the revised version and deaths registered in 2017 and 2018 are based on the preliminary version. Revised and preliminary versions are subject to further revision by the Australian Bureau of Statistics (ABS).
3. The CODURF is provided to the AIHW by the Registries of Births, Deaths and Marriages and the National Coronial Information System (managed by the Victorian Department of Justice) and includes cause of death as coded by the ABS. The data are maintained by the AIHW in the National Mortality Database.

Sources: AIHW National Mortality Database; Table A6 in [Mesothelioma in Australia 2020—data tables](#).

From 1997 to 2019, age-standardised mesothelioma mortality rates fluctuated between 2.1 and 2.7 deaths per 100,000 population (Figure 9). Rates for men ranged from 4.0 deaths per 100,000 in 2006 to 5.0 deaths per 100,000 population in 2001 and 2004. Rates for women fluctuated around 0.8 deaths per 100,000 over the period.

Figure 9: Age-standardised rate (per 100,000 population) of deaths from mesothelioma, by year and sex, 1997 to 2019



Notes:

1. Figures are presented by year of death for 1997–2018 and year of registration of death for 2019.
2. Deaths registered in 2015 and earlier are based on the final version of the Cause of Death Unit Record File (CODURF), deaths registered in 2016 are based on the revised version and deaths registered in 2017 and 2018 are based on the preliminary version. Revised and preliminary versions are subject to further revision by the Australian Bureau of Statistics (ABS).
3. The CODURF is provided to the AIHW by the Registries of Births, Deaths and Marriages and the National Coronial Information System (managed by the Victorian Department of Justice) and includes cause of death as coded by the ABS. The data are maintained by the AIHW in the National Mortality Database.
4. Rates have been age-standardised to the 2001 Australian Standard Population.

Source: AIHW National Mortality Database; Table A6 in [Mesothelioma in Australia 2020—data tables](#).

Asbestos exposure among people with mesothelioma

Previous research has shown mesothelioma can be associated with occupational and non occupational exposure to asbestos. Historically, occupational exposure has been dominated by asbestos mining, manufacturing and use of asbestos-containing materials (enHealth 2013; Safe Work Australia 2014). Because mesothelioma typically develops a long time after exposure, the majority of current cases in this report are likely related to occupational exposure in workplaces that occurred before current occupational asbestos regulations and practices came into effect. The results here should not be interpreted as indicative of current risk in workplaces today.

In total, 1,253 people (1,000 men and 253 women) diagnosed with mesothelioma since 1 July 2010 consented to participate in the voluntary asbestos exposure assessments at 1 May 2021. Of these, 1,098 (867 men and 231 women) completed both the questionnaire and telephone interview components of the assessment.

Based on the jobs held by the participant during their working life, relevant job-specific questionnaire modules were allocated for the participant's telephone interview. For example, participants who have worked in jobs such as electrician, plumber and carpenter may be allocated the 'Trades' module. Participants could be allocated multiple job-specific modules for different jobs.

Participants' lifetime exposure in non-occupational settings (such as their home) was also evaluated with a non-occupational module. For the purposes of this assessment, potential exposures to asbestos were then classified according to the likelihood that they were above background levels of 0.0001 f/ml (fibres of asbestos per millilitre) (Brown 2001). Probability of exposure was assessed as either 'probable', 'possible' or 'unlikely', and level of exposure as either 'high', 'medium' or 'low'. For more information, see *Mesothelioma in Australia 2020: methodology paper*.

Nine in ten participants were assessed as having some exposure to asbestos

More than 9 in 10 (94% or 1,028) people were assessed as having possible or probable exposure to asbestos (Table 3). For men, this exposure most commonly occurred in their jobs. Occupational exposure to asbestos is typically associated with higher frequency, dose and duration of exposure than non-occupational exposure.

Of the 1,028 people for whom exposure was detected:

- 78% (646) of men provided information indicating possible or probable occupational exposure ('occupational exposure only' and 'both occupational and non-occupational exposure'), compared with 6.8% (14) of women (Table 3).
- 99% (204) of women provided information indicating non-occupational exposure ('non occupational exposure only' and 'both occupational and non-occupational exposure'), compared with 85% (699) of men (Table 3).

Table 3: Occupational and non-occupational exposure assessment, by sex, 2010–2020

Any exposure indicated	Men		Women		People	
	No.	%	No.	%	No.	%
Occupational exposure only	124	15.1	1	0.5	125	12.2
Non-occupational exposure only	177	21.5	191	93.2	368	35.8
Both occupational and non-occupational exposure	522	63.4	13	6.3	535	52.0
Total	823	100	205	100	1,028	100

Note: Of the 1,098 participants, 70 (6.4%) of participants (44 men and 26 women) were assessed as having neither occupational nor non-occupational exposure. Although it was not possible to identify asbestos exposure among these participants, this should not be taken to mean that these participants have never been exposed to asbestos; rather it means that no evidence of above background exposure was obtained by the exposure assessment methods used.

Source: AIHW analysis of AMR data at 1 May 2021, based on interviews completed among people who were diagnosed with mesothelioma between 1 July 2010–31 December 2020.

Table 4 shows the results from the asbestos exposure assessment among the 5 states and territories with the largest number of participants; New South Wales, Victoria, Queensland, Western Australia and South Australia. There were insufficient numbers of participants to present results for Northern Territory, Australian Capital Territory and Tasmania, so these were excluded.

Table 4 shows that the proportion of participants assessed as having:

- 'occupational exposure only' ranged from 11% to 13% across states
- 'non-occupational exposure only' ranged from 29% to 42% across states
- 'both occupational and non-occupational exposure' ranged from 45% to 60% across states.

As these percentages are based on small numbers of participants, drawing conclusions about apparent differences between states is not recommended.

Table 4: Occupational and non-occupational exposure assessment, by state, 2010–2020

Any exposure indicated	State									
	NSW		Vic		Qld		WA		SA	
	No.	%	No.	%	No.	%	No.	%	No.	%
Occupational exposure only	46	13.2	26	12.7	20	11.0	18	10.5	9	11.5
Non-occupational exposure only	123	35.3	87	42.4	53	29.3	57	33.3	33	42.3
Both occupational and non-occupational exposure	179	51.4	92	44.9	108	59.7	96	56.1	36	46.2
Total	348	100	205	100	181	100	171	100	78	100

Notes:

1. Of the 1,098 participants, 70 (6.4%) of participants (44 men and 26 women) were assessed as having neither occupational nor non-occupational exposure. Although it was not possible to identify asbestos exposure among these participants, this should not be taken to mean that these participants have never been exposed to asbestos; rather it means that no evidence of above background exposure was obtained by the exposure assessment methods used.

2. There were insufficient numbers of participants to present results for NT, ACT and Tasmania.


Source: AIHW analysis of AMR data at 1 May 2021, based on interviews completed among people who were diagnosed with mesothelioma between 1 July 2010–31 December 2020.

Occupational asbestos exposure

Figure 10 shows the estimated likelihood of exposure among jobs assessed using the three most commonly allocated job-specific interview modules. Based on the interview data collected, 81% of participants who received the ‘Trades’ module, 39% of participants who received the ‘Land transport’ module, and 73% of those who received the ‘Water transport’ module, were assessed as having had ‘probable’ exposure to asbestos for jobs in those categories.

Results are based on the responses people gave to each specific module - participants are typically assigned multiple job-specific modules for different jobs, so the sum of respondents assigned to each module does not equal the total number of participants. Because questionnaire modules are assigned only to jobs with some likelihood of exposure, the finding of exposure in a high proportion of the jobs that respondents were questioned about is to be expected.

Figure 10: Exposure assessment results by most commonly used job-specific modules, 2010–2020



Source: AIHW analysis of AMR data at 1 May 2021, based on interviews completed among people who were diagnosed with mesothelioma between 1 July 2010–31 December 2020; Table A7 in [Mesothelioma in Australia 2020—data tables](#).

Occupational exposure by job type


For participants who received the same module for more than one job, the results presented are based on a participant's highest exposed job in that job category. Because participants can have different exposure probabilities and/or levels for different jobs, the exposure estimate reported here is the maximum exposure likely in that job category for each individual. For example, if a participant had two trade-related jobs which were assessed as having a different probability and/or level of exposure, the higher of the two would be reported for that participant in that job category. Participants may be assigned different modules for different types of jobs, so the sum of respondents assigned to each module does not equal the total number of participants.

Jobs are coded according to the Australian and New Zealand Standard Classification of Occupations (ANZSCO).

Among participants who received the Trades module (Figure 11):

- 74% of people who had metal fitter/turner/toolmaker jobs were assessed as having had probable exposure in those jobs. For 73% of those with probable exposure, the level of exposure to asbestos was classified as 'high'.
- 85% of people who had plumber/gasfitter jobs, 84% of those who had electrical trade jobs, and 85% who had jobs in building trades, were assessed as having had probable exposure in those jobs. For the majority of these, the level of exposure was classified as 'high'.
- 60% of people who had telecommunications technician jobs and 35% who worked in engineer jobs were assessed as 'unlikely' to have been exposed in those jobs.

Figure 11: Occupational asbestos exposure by job title for the 'Trades' module, 2010–2020



Source: AIHW analysis of AMR data at 1 May 2021, based on interviews completed among people who were diagnosed with mesothelioma between 1 July 2010–31 December 2020; Table A8 in [Mesothelioma in Australia 2020—data tables](#).


Among participants who received the water transport module (Figure 12):

- 100% of people with marine engineer jobs were assessed as having had probable exposure, with the level of exposure estimated as 'high' for 79% of these. 72% of participants with shipwright/boat builder jobs were estimated to have had probable exposure.

Among participants who received the land transport module (Figure 12):

- 67% of people who had jobs in automotive maintenance, and 49% of those with driving jobs, were assessed as having had probable exposure—among those assessed as probably exposed in these jobs, the level of exposure was estimated as 'high' for most.

Figure 12: Occupational asbestos exposure by job title for the 'Water transport' and 'Land transport' modules, 2010–2020



Source: AIHW analysis of AMR data at 1 May 2021, based on interviews completed among people who were diagnosed with mesothelioma between 1 July 2010–31 December 2020; Table A8 in [Mesothelioma in Australia 2020—data tables](#).

Non-occupational asbestos exposure

All 1,098 participants completed the non-occupational questionnaire module. Of these, 903 (88%) were assessed as having had possible or probable exposure in non-occupational contexts. It was common to have indications of exposure in more than one non-occupational context, so a number of participants are counted in more than one category. Therefore, the sum of participants in each category does not add up to the total number of participants. The most common contexts in which non-occupational asbestos exposure was assessed as possible or probable were among those who reported ever having:

- undertaken major home renovations that involved asbestos products (including paid work) (42% assessed as possible exposure, 7.8% assessed as probable exposure)
- lived in a house undergoing renovations (38% assessed as possible exposure)
- serviced car brakes/clutch (excluding paid work) (30% assessed as probable exposure)
- lived in the same home as someone with a job where they were exposed to asbestos and who came home dusty (13% assessed as possible exposure, 6.4% assessed as probable exposure).
- lived in a house made of fibro that was built between 1947 and 1987 (11% assessed as probable exposure).

See Table A9 in [Mesothelioma in Australia 2020—data tables](#) for further information on sources of non-occupational exposure.

No asbestos exposure

For 70 participants (6.4% of all 1,098 participants—44 men and 26 women), there was no indication of asbestos exposure above background levels in either occupational or non-occupational contexts. This does not mean that these participants have never been exposed to asbestos; rather it means that no evidence of exposure above background levels was obtained using the exposure assessment methods.

What are the challenges in collecting data and reporting on mesothelioma?

Confirming a diagnosis of mesothelioma is often very challenging for a variety of reasons:

- Symptoms of mesothelioma are common to many other conditions, and mesothelioma cells can often look similar to cells of other cancers (Cancer Council 2019b).
- Because mesothelioma symptoms are not specific to the condition, diagnosis is often complicated. Diagnostic confirmation of mesothelioma generally involves a number of clinical investigations, including biopsies, radiology and clinical examinations conducted by a multidisciplinary team (van Zandwijk et al. 2013).
- Diagnostic and treatment practices for mesothelioma are not equally distributed across Australia (van Zandwijk et al. 2013).

If a mesothelioma diagnosis is uncertain for any reason, the AMR is not notified and the case remains unrecorded, until such a time as the diagnosis is confirmed. Other challenges in collecting data and reporting on mesothelioma include:

- Although state and territory cancer registries fast-track mesothelioma notifications, there is still a time lag between a person's diagnosis, their inclusion in the AMR data set and (if consent is given) when they are interviewed for the AMR's asbestos exposure collection. Reasons for this lag include the time it takes to make a definitive diagnosis, the time between diagnosis and notification to cancer registries and to the AMR. Case verification and recruitment processes also vary between state and territory cancer registries.
- The participation rate in the asbestos exposure assessment of the AMR is low, at around 15% of people with mesothelioma. This is partially due to people dying or being too unwell to participate.

Where do I go for more information?

More information on the AMR is available at www.mesothelioma-australia.com/home. The report Mesothelioma in Australia 2020 and previous annual reports are available at <www.mesothelioma-australia.com/publications-and-data/publications>. People diagnosed with mesothelioma can choose to self-notify by contacting the AMR via email at <amr@aihw.gov.au> or via the toll-free information line on 1800 378 861.

Glossary

age-adjusted rate: See age-standardised rate.

age standardisation: A way to remove the influence of age when comparing populations with different age structures. This is usually necessary because the rates of many diseases vary strongly with age (usually increasing with increasing age). The age structures of the different populations are converted to the same 'standard' structure and then the disease rates that would have occurred with that structure are calculated and compared. This report uses the direct method of age standardisation.

age standardised rate: A rate that results from removing the influence of age by converting the age structures of the different populations to the same 'standard' structure, providing a more valid way of comparing rates from populations with different age structures.

age-specific rate: The rate for a specific age-group. The numerator and denominator pertain to the same age group.

Australian Cancer Database (ACD): Contains data on all new cases of cancer diagnosed in Australia (except basal and squamous cell carcinomas of the skin) since 1982.

incidence: The number of new cases (of an illness or event, and so on) occurring during a given period, often expressed as a rate (number per population).

mortality: The number or rate of deaths in a population during a given time period.

National Death Index (NDI): A catalogue of death records used in data linkage for epidemiological studies.

National Mortality Database (NMD): Holds records for all deaths in Australia since 1964.

non-occupational exposures: Chemical, biological, psychosocial, physical and other factors from places other than the workplace that can potentially cause harm. Examples include contact with asbestos during private house renovations and living in the same home as someone with an asbestos-exposed occupation who came home dusty.

occupational exposures: Chemical, biological, psychosocial, physical and other factors in the workplace that can potentially cause harm.

relative survival: the probability of being alive for a given amount of time after diagnosis compared with the general population. A 5-year relative survival figure of 100% means that the cancer has no impact on the person's chance of still being alive 5 years after diagnosis, whereas a figure of 50% means that the cancer has halved that chance.

underlying cause of death: The disease or injury that initiated the sequence of events leading directly to death.

References

- ABS (Australian Bureau of Statistics) 2019. Australian Historical Population Statistics, 2016. Canberra: ABS. Viewed 1 May 2021, <https://www.abs.gov.au/AUSSTATS/abs@.nsf/mf/3105.0.65.001>.
- AIHW (Australian Institute of Health and Welfare) 2018a. Cancer data in Australia. Cat. no. CAN 122. Canberra: AIHW.
- AIHW 2018b. Mesothelioma in Australia 2017. Cat. no. CAN 121. Canberra: AIHW.
- AIHW 2019. Mesothelioma in Australia 2018. Cat. no. CAN 130. Canberra: AIHW.
- AIHW 2020. Mesothelioma in Australia 2019. Cat. no. CAN 134. Canberra: AIHW.
- AIHW 2021. Disease expenditure in Australia 2018–19. Canberra: AIHW. Viewed 10 September 2021, <https://www.aihw.gov.au/reports/health-welfare-expenditure/spending-on-disease-in-australia/data>.
- AMR (Australian Mesothelioma Registry) 2012. Australian Mesothelioma Registry 1st annual report: Mesothelioma in Australia 2011. NSW Health: Cancer Institute NSW.
- AMR 2013. Australian Mesothelioma Registry 2nd annual report: Mesothelioma in Australia 2012. Sydney: NSW Health—Cancer Institute NSW.
- AMR 2014. Australian Mesothelioma Registry 3rd annual report: Mesothelioma in Australia 2013. Sydney: NSW Health—Cancer Institute NSW.
- AMR 2015. Australian Mesothelioma Registry 4th annual report: Mesothelioma in Australia 2014. Sydney: NSW Health—Cancer Institute NSW.
- AMR 2016. Australian Mesothelioma Registry 5th annual report: Mesothelioma in Australia 2015. Sydney: NSW Health—Cancer Institute NSW.
- AMR 2017. Australian Mesothelioma Registry 6th annual report: Mesothelioma in Australia 2016. Sydney: NSW Health—Cancer Institute NSW.
- Asbestos Diseases Research Institute 2019. Mesothelioma. Sydney: Asbestos Diseases Research Institute. Viewed 5 May 2021, <https://adri.org.au/mesothelioma/>.
- ASEA (Asbestos Safety and Eradication Agency) 2016. Asbestos Safety and Eradication Agency Reports 2016. Asbestos: the International and Australian Contexts. Viewed 12 May 2021, https://www.asbestossafety.gov.au/sites/default/files/documents/2017-10/ASEA_Reports_Asbestos_International_Australian_contexts_ACC.pdf
- Bray F, Colombet M, Mery L, Piñeros M, Znaor A, Zanetti R et al. 2017. Cancer incidence in five continents, Vol. XI. Lyon: International Agency for Research on Cancer. Viewed 4 July 2021, <http://ci5.iarc.fr>.
- Burgers JA & Damhuis RA 2004. Prognostic factors in malignant mesothelioma. Lung Cancer 45 Supplement 1:S49–S54.
- Cancer Council 2019a. Mesothelioma. Sydney: Cancer Council. Viewed 16 April 2021, <https://www.cancer.org.au/about-cancer/types-of-cancer/mesothelioma.html>.
- Cancer Council 2019b. Trimodality therapy for pleural mesothelioma. Sydney: Cancer Council. Viewed 29 May 2020, <https://www.cancercouncil.com.au/pleural-mesothelioma/active-treatment/trimodality-therapy/>
- Cancer Council 2019c. Diagnosing mesothelioma. Sydney: Cancer Council. Viewed 29 May 2021, <https://www.cancervic.org.au/cancer-information/types-of-cancer/mesothelioma/diagnosing-mesothelioma.html>.

enHealth (Environmental Health Standing Committee) 2013. Asbestos: a guide for householders and the general public. Canberra: Australian Health Protection Principal Committee.

Faig J, Howard S, Levine EA, Casselman G, Hesdorffer M & Ohar JA 2015. Changing pattern in malignant mesothelioma survival. *Translational oncology* 8(1):35–39.

Leigh J, Davidson P, Hendrie L & Berry D 2002. Malignant mesothelioma in Australia, 1945–2000. *American Journal of Industrial Medicine* 41:188–201.

Marinaccio A, Binazzi A, Cauzillo G, Cavone D, De Zotti R, Mensi C et al. 2007. Analysis of latency time and its determinants in asbestos related malignant mesothelioma cases of the Italian register. *European Journal of Cancer* 43(18):2722–2728.

Musk AW, Olsen N, Alfonso H, Reid A, Mina R, Franklin P et al. 2011. Predicting survival in malignant mesothelioma. *European Respiratory Journal* 38(6):1420–1424.

Olsen NJ, Franklin PF, Reid, A, de Klerk NH, Threlfall TJ, Shilkin K et al. 2011. Increasing incidence of malignant mesothelioma after exposure to asbestos during home maintenance and renovation. *The Medical Journal of Australia* 195(5): 271–274.

Reid A, de Klerk NH, Magnani C, Ferrante D, Berry G, Musk AW et al. 2014. Mesothelioma risk after 40 years since first exposure to asbestos: a pooled analysis. *Thorax* 69:843–850.

Safe Work Australia 2014. Asbestos-related disease indicators. Canberra: Safe Work Australia. Viewed 12 June 2021, https://www.safeworkaustralia.gov.au/system/files/documents/1702/asbestos_related_disease_indicators_2014.pdf.

Shavelle R, Vavra-Musser K, Lee K & Brookes J 2017. Life Expectancy in Pleural and Peritoneal Mesothelioma. *Lung Cancer International*. doi: 10.1155/2017/2782590.

Soeberg MJ, Vallance DA, Keena V, Takahashi K & Leigh J. 2018. Australia's ongoing legacy of asbestos: significant challenges remain even after the complete banning of asbestos almost fifteen years ago. *International Journal of Environmental Research and Public Health* 15:384. doi:10.3390/ijerph15020384

van Zandwijk N, Clarke C, Henderson D, Musk AW, Fong K, Nowak A, et al. 2013. Guidelines for the diagnosis and treatment of malignant pleural mesothelioma. *Journal of Thoracic Disease* 5(6): E254–E307.

Acknowledgments

This report was prepared by Tom Watson and Sue Barker from the AMR at the AIHW. The AMR team would also like to acknowledge the valuable guidance and advice from Justin Harvey and Richard Juckes. The AMR Expert Advisory Group also provided valuable advice: Professor Tim Driscoll (Chair), Associate Professor Fraser Brims, Ms Elizabeth Chalker, Dr Claire Cooke Yarborough, Ms Justine Ross, Professor Malcolm Sim and Professor Ken Takahashi.

The AIHW acknowledges the data and assistance provided by all state and territory cancer registries, and thanks staff at the Monash Centre for Occupational and Environmental Health for undertaking and reporting on the asbestos exposure component of the AMR, and staff at the Social Research Centre for their work interviewing people diagnosed with mesothelioma.

The AMR is funded by Safe Work Australia.



© Australian Institute of Health and Welfare 2021 

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 3.0 (CC BY 3.0) licence (<http://creativecommons.org/licenses/by/3.0/au/>). You may distribute, remix and build upon this work. However, you must attribute the AIHW as the copyright holder of the work in compliance with our attribution policy available at www.aihw.gov.au/copyright/. The full terms and conditions of this licence are available at <http://creativecommons.org/licenses/by/3.0/au/>.

Suggested citation

Australian Institute of Health and Welfare 2021. Mesothelioma in Australia 2020. Cat. no. CAN 143. Canberra: AIHW.

ISBN 978-1-76054-921-3 (Online)
ISBN 978-1-76054-922-0 (Print)
DOI: 10.25816/2g6c-8q81

Any enquiries about copyright should be directed to the Australian Institute of Health and Welfare, GPO Box 570, Canberra ACT 2601,
Tel: (02) 6244 1000, Email: info@aihw.gov.au.



Stronger evidence,
better decisions,
improved health and welfare